

Greenup Analyses

Kara Dobson

February 26, 2021

COLLABORATORS: Phoebe Zarnetske, Moriah Young, Mark Hammond, Pat Bills

DATA INPUT: Data imported as csv files from shared Google drive L1 plant comp folder

PROJECT: warmXtrophic

The final_kbs and final_umbs dataframes contain data for greenup at each site.

“half_cover_date” is the date at which 50% of a species max cover was reached (per plot, per year)

“state” describes each treatment - warmed or ambient

```
# Clear all existing data
rm(list=ls())

#Load packages
library(tidyverse)
library(lme4)
library(olsrr)
library(predictmeans)
library(car)
library(fitdistrplus)
library(ggpubr)
library(rstatix)
library(vegan)

# Set working directory to Google Drive
# **** Update with the path to your Google drive on your computer
setwd("/Volumes/GoogleDrive/Shared drives/SpaCE_Lab_warmXtrophic/data/")

# Read in plant comp data
greenup <- read.csv("L1/greenup/final_greenup_L1.csv")
#str(greenup)

# changing scale of years
greenup$year[greenup$year == 2015] <- 1
greenup$year[greenup$year == 2016] <- 2
greenup$year[greenup$year == 2017] <- 3
greenup$year[greenup$year == 2018] <- 4
greenup$year[greenup$year == 2019] <- 5
greenup$year[greenup$year == 2020] <- 6


# create dataframes for kbs and umbs
final_kbs <- subset(greenup, site == "kbs")
```

```
final_umbs <- subset(greenup, site == "umbs")
```

Starting with KBS


First, checking for normality

```
hist(final_kbs$half_cover_date)
```



```
qqnorm(final_kbs$half_cover_date)
```

Normal Q-Q Plot




```
shapiro.test(final_kbs$half_cover_date)
```

```
##  
##  Shapiro-Wilk normality test  
##  
## data:  final_kbs$half_cover_date  
## W = 0.94273, p-value < 2.2e-16
```

```
hist(final_kbs$half_cover_date[final_kbs$state == "ambient"])
```


Histogram of final_kbs\$half_cover_date[final_kbs\$state == "ambient"]



final_kbs\$half_cover_date[final_kbs\$state == "ambient"]

```
hist(final_kbs$half_cover_date[final_kbs$state == "warmed"])
```

Histogram of final_kbs\$half_cover_date[final_kbs\$state == "warmed"]




final_kbs\$half_cover_date[final_kbs\$state == "warmed"]

Not normal, and previously attempted transformations don't help
Seeing what other distribution could fit

```
descdist(final_kbs$half_cover_date, discrete = FALSE)
```

Cullen and Frey graph




```
## summary statistics
## -----
## min: 59   max: 289
## median: 141
## mean: 152.8644
## estimated sd: 57.73216
## estimated skewness: 0.3763172
## estimated kurtosis: 2.058193
```

While uniform looks the closest, I'll try poisson


```
fit <- lm(half_cover_date~state, data = final_kbs)
residual <- fit$residuals
hist(residual, main="Raw residuals")
```

Raw residuals



```
pois <- glm(half_cover_date~state, data = final_kbs, family="poisson")  
hist(pois$residuals, main="Poisson glm residuals")
```

Poisson glm residuals



Below I try a few different generalized linear models with poisson distribution:

An interaction between state and year, plus insecticide as a fixed effect and species and plot as random effects

```
moda <- glmer(half_cover_date ~ state*year + insecticide + (1|species) + (1|plot),
              data=final_kbs, family = poisson)
```

```
## Warning in checkConv(attr(opt, "derivs"), opt$par, ctrl = control$checkConv, : Model is nearly unidentifiable:
## - Rescale variables?
```

```
summary(moda)
```

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
## Approximation) [glmerMod]
## Family: poisson ( log )
## Formula: half_cover_date ~ state * year + insecticide + (1 | species) +
## (1 | plot)
## Data: final_kbs
##
##          AIC          BIC    logLik deviance df.resid
## 42924.6   42963.9 -21455.3  42910.6      2021
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -9.0161 -2.8352 -0.6425  2.3763 14.4656
##
## Random effects:
## Groups Name Variance Std.Dev.
## species (Intercept) 0.069953 0.26449
## plot (Intercept) 0.001848 0.04299
## Number of obs: 2028, groups: species, 55; plot, 24
##
## Fixed effects:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)    5.020672   0.039631 126.686 < 2e-16 ***
## statewarmed     0.009389   0.019347   0.485  0.6275
## year           0.003600   0.001520   2.368  0.0179 *
## insecticideno_insects 0.028935   0.017936   1.613  0.1067
## statewarmed:year -0.009698   0.002116 -4.584 4.57e-06 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##              (Intr) sttwrm year  insct_
## statewarmed -0.243
## year        -0.130  0.249
## insctcdn_ns -0.227 -0.002  0.008
## statwrmd:yr  0.089 -0.375 -0.648  0.006
## convergence code: 0
## Model is nearly unidentifiable: very large eigenvalue
## - Rescale variables?
```

No interaction between state and year, but with state and insecticide as fixed effects and species and plot as random effects

```
modb <- glmer(half_cover_date ~ state + year + insecticide + (1|species) + (1|plot),
              data=final_kbs, family = poisson)
```

```
## Warning in checkConv(attr(opt, "derivs"), opt$par, ctrl = control$checkConv, : Model is nearly unidentifiable:
## - Rescale variables?
```

```
summary(modb)
```

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
## Approximation) [glmerMod]
## Family: poisson ( log )
## Formula: half_cover_date ~ state + year + insecticide + (1 | species) +
## (1 | plot)
## Data: final_kbs
##
##      AIC      BIC    logLik deviance df.resid
## 42943.6 42977.3 -21465.8 42931.6      2022
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -9.0054 -2.8448 -0.6575  2.3929 14.2130
##
## Random effects:
## Groups Name          Variance Std.Dev.
## species (Intercept) 0.069954 0.26449
## plot (Intercept)    0.001847 0.04297
## Number of obs: 2028, groups: species, 55; plot, 24
##
## Fixed effects:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)      5.0367370  0.0394718 127.603  <2e-16 ***
## statewarmed      -0.0238778  0.0179289  -1.332   0.183
## year             -0.0009129  0.0011579  -0.788   0.430
## insecticideno_insects 0.0294349  0.0179300   1.642   0.101
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##              (Intr) sttwrm year
## statewarmed -0.227
## year        -0.096  0.007
## insctcdn_ns -0.228  0.001  0.015
## convergence code: 0
## Model is nearly unidentifiable: very large eigenvalue
## - Rescale variables?
```


State and insecticide as fixed effects & year, species and plot as random effects

```
modc <- glmer(half_cover_date ~ state + insecticide + (1|year) + (1|species) + (1|plot),
              data=final_kbs, family = poisson)
summary(modc)
```

```
## Generalized linear mixed model fit by maximum likelihood (Laplace
##   Approximation) [glmerMod]
##   Family: poisson ( log )
## Formula: half_cover_date ~ state + insecticide + (1 | year) + (1 | species) +
##           (1 | plot)
##   Data: final_kbs
##
##           AIC          BIC    logLik deviance df.resid
##  41436.8   41470.5 -20712.4  41424.8      2022
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -9.3418 -2.8134 -0.5215  2.1783 13.1976
##
## Random effects:
##   Groups Name          Variance Std.Dev.
##   species (Intercept) 0.068408 0.26155
##   plot    (Intercept) 0.001820 0.04266
##   year    (Intercept) 0.005867 0.07659
## Number of obs: 2028, groups: species, 55; plot, 24; year, 6
##
## Fixed effects:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)      5.01945    0.04991 100.577 <2e-16 ***
## statewarmed      -0.02683    0.01781  -1.507  0.1319
## insecticideno_insects 0.03093    0.01781   1.737  0.0824 .
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##              (Intr) sttwrm
## statewarmed -0.178
## insctcdn_ns -0.178  0.001
```

Because no distributions seems to match well, I'll try a Friedman's test

```
#friedman_kbs <- final_kbs %>%  
# friedman_test(half_cover_date ~ state)
```

Error: Must extract column with a single valid subscript. x Subscript var can't be NA

Can't figure out what this means

If I include the blocks portion of the formula (from the documentation) I get this error

```
#friedman_kbs <- final_kbs %>%  
# friedman_test(half_cover_date ~ state | plot)
```

Error in friedman.test.default(c(141L, 202L, 122L, 101L, 127L, 120L, 197L, : not an unreplicated complete block design


Permanova?

```
knitr::opts_chunk$set(cache = TRUE)  
per1 <- adonis2(final_kbs$half_cover_date ~ state*year + insecticide, data = final_kbs)  
per1  
  
## Permutation test for adonis under reduced model  
## Terms added sequentially (first to last)  
## Permutation: free  
## Number of permutations: 999  
##  
## adonis2(formula = final_kbs$half_cover_date ~ state * year + insecticide, data = final_kbs)  
##           Df SumOfSqs      R2      F Pr(>F)  
## state      1    0.051 0.00074  1.5046  0.210  
## year      1    0.487 0.00698 14.2700  0.001 ***  
## insecticide 1    0.168 0.00241  4.9326  0.018 *  
## state:year  1    0.025 0.00036  0.7290  0.405  
## Residual   2023   69.058 0.98951  
## Total      2027   69.790 1.00000  
## ---  
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

UMBS


Checking for normality

```
hist(final_umbs$half_cover_date)
```



```
qqnorm(final_umbs$half_cover_date)
```

Normal Q-Q Plot




```
shapiro.test(final_umbs$half_cover_date)
```

```
##  
##  Shapiro-Wilk normality test  
##  
## data:  final_umbs$half_cover_date  
## W = 0.94754, p-value < 2.2e-16
```

```
hist(final_umbs$half_cover_date[final_kbs$state == "ambient"])
```


Histogram of final_umbs\$half_cover_date[final_kbs\$state == "ambier



final_umbs\$half_cover_date[final_kbs\$state == "ambient"]

```
hist(final_umbs$half_cover_date[final_kbs$state == "warmed"])
```

Histogram of final_umbs\$half_cover_date[final_kbs\$state == "warmed




final_umbs\$half_cover_date[final_kbs\$state == "warmed"]

These look pretty good

Trying log transformation

```
final_umbs$date_log <- log(final_umbs$half_cover_date)
hist(final_umbs$date_log)
```




```
shapiro.test(final_umbs$date_log)
```

```
##
##  Shapiro-Wilk normality test
##
## data:  final_umbs$date_log
## W = 0.97728, p-value = 6.765e-14
```

I think this looks good but shapiro-wilk is lower than 0.05

Trying inverse tranformation

```
final_umbs$date_inv <- 1/(final_umbs$half_cover_date)
hist(final_umbs$date_inv)
```



```
shapiro.test(final_umbs$date_inv)
```

```
##
##  Shapiro-Wilk normality test
##
## data:  final_umbs$date_inv
## W = 0.97993, p-value = 6.679e-13
```

This also looks good but is also still low for shapiro-wilk