

GEN 242: Linear Algebra

Chapter 8: Vectors/Matrix Differential Calculus

Solutions Guide

Instructor: Richard Bahin

Full Sail University

Table of Contents

Answers.....	5
Converting from Rectangular to Polar Coordinates	5
Converting from Polar to Rectangular Coordinates	5
Converting from Cylindrical to Rectangular Coordinates	5
Converting from Rectangular to Cylindrical Coordinates	6
Converting from Spherical to Rectangular Coordinates	6
Converting from Rectangular to Spherical Coordinates	6
Converting from Spherical to Cylindrical Coordinates.....	7
Converting from Cylindrical to Spherical Coordinates.....	7
Gradient of a Scalar Field	7
Curl of a Vector Field	7
Divergence of a Vector Field	8
Laplacian of a Scalar Field	8
Laplacian of a Vector Field	8
Derivative of a Vector with Respect to a Vector	8
Derivative of a Scalar with Respect to a Vector	9
Quadric Forms	9
Solutions.....	10
Rectangular to Polar Coordinates Conversion.....	10
Problem 1.....	10
Problem 2.....	11
Problem 3.....	12
Polar to Rectangular Coordinates Conversion.....	13
Problem 4.....	13
Problem 5.....	14
Problem 6.....	15
Cylindrical to Rectangular Coordinates Conversion	16
Problem 7.....	16
Problem 8.....	17
Problem 9.....	18
Rectangular to Cylindrical Coordinates Conversion	19
Problem 10.....	19
Problem 11.....	20
Problem 12.....	21

Spherical to Rectangular Coordinates Conversion	22
Problem 13	22
Problem 14	23
Problem 15	24
Rectangular to Spherical Coordinates Conversion	25
Problem 16	25
Problem 17	26
Problem 18	27
Spherical to Cylindrical Coordinates Conversion.....	28
Problem 19	28
Problem 20	29
Problem 21	30
Cylindrical to Spherical Coordinates Conversion.....	31
Problem 22	31
Problem 23	32
Problem 24	33
Gradient of a Scalar Field	34
Problem 25	34
Problem 26	34
Problem 27	35
Curl of a Vector Field	36
Problem 28	36
Problem 29	37
Problem 30	38
Computing the Divergence of a Vector Field.....	39
Problem 31	39
Problem 32	39
Problem 33	40
Computing the Laplacian of a Scalar Field.....	41
Problem 34	41
Problem 35	42
Problem 36	43
Computing the Laplacian of a Vector Field.....	44
Problem 37	44
Problem 38	45
Problem 39	46
Derivative of a Vector with Respect to a Vector	47
Problem 40	47
Problem 41	47
Problem 42	48
Derivative of a Scalar with Respect to a Vector (Jacobian)	49
Problem 43	49
Problem 44	49
Problem 45	50

Quadric Forms	51
Problem 46	51
Problem 47	52
Problem 48	53

Answers

Converting from Rectangular to Polar Coordinates

$$1. \vec{p} = (3, \sqrt{3})_{\text{rect}} \rightarrow \boxed{\vec{p} = \left(\frac{\pi}{6}, 2\sqrt{3}\right)_{\text{polar}}}$$

$$2. \vec{p} = (1, \sqrt{3})_{\text{rect}} \rightarrow \boxed{\vec{p} = \left(\frac{\pi}{3}, 2\right)_{\text{polar}}}$$

$$3. \vec{p} = (-1, -1)_{\text{rect}} \rightarrow \boxed{\vec{p} = \left(-\frac{3\pi}{4}, \sqrt{2}\right)_{\text{polar}}}$$

Converting from Polar to Rectangular Coordinates

$$4. \vec{p} = \left(\frac{7\pi}{4}, 3\sqrt{2}\right)_{\text{polar}} \rightarrow \boxed{\vec{p} = (3, -3)_{\text{rect}}}$$

$$5. \vec{p} = \left(\frac{\pi}{4}, 4\right)_{\text{polar}} \rightarrow \boxed{\vec{p} = (2\sqrt{2}, 2\sqrt{2})_{\text{rect}}}$$

$$6. \vec{p} = \left(\frac{\pi}{3}, 1\right)_{\text{polar}} \rightarrow \boxed{\vec{p} = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)_{\text{rect}}}$$

Converting from Cylindrical to Rectangular Coordinates

$$7. \vec{p} = \left(3, \frac{\pi}{2}, 1\right)_{\text{cyl}} \rightarrow \boxed{\vec{p} = (0, 3, 1)_{\text{rect}}}$$

$$8. \vec{p} = \left(4, \frac{\pi}{6}, 2\right)_{\text{cyl}} \rightarrow \boxed{\vec{p} = (2\sqrt{3}, 2, 2)_{\text{rect}}}$$

$$9. \vec{p} = \left(1, \frac{\pi}{4}, 5\right)_{\text{cyl}} \rightarrow \boxed{\vec{p} = \left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}, 5\right)_{\text{rect}}}$$

Converting from Rectangular to Cylindrical Coordinates

$$10. \vec{p} = (1, 1, 1)_{\text{rect}} \rightarrow \vec{p} = \left(\sqrt{2}, \frac{\pi}{4}, 1 \right)_{\text{cyl}}$$

$$11. \vec{p} = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}, 5 \right)_{\text{rect}} \rightarrow \vec{p} = \left(1, \frac{\pi}{3}, 5 \right)_{\text{cyl}}$$

$$12. \vec{p} = (\sqrt{2}, \sqrt{2}, 3)_{\text{rect}} \rightarrow \vec{p} = \left(2, \frac{\pi}{4}, 3 \right)_{\text{cyl}}$$

Converting from Spherical to Rectangular Coordinates

$$13. \vec{p} = \left(1, \frac{\pi}{4}, \pi \right)_{\text{sphere}} \rightarrow \vec{p} = (0, 0, -1)_{\text{rect}}$$

$$14. \vec{p} = \left(3, \frac{\pi}{3}, \frac{\pi}{4} \right)_{\text{sphere}} \rightarrow \vec{p} = \left(\frac{3\sqrt{6}}{4}, \frac{3\sqrt{6}}{4}, \frac{3\sqrt{2}}{2} \right)_{\text{rect}}$$

$$15. \vec{p} = \left(5, \frac{\pi}{2}, \pi \right)_{\text{sphere}} \rightarrow \vec{p} = (0, -5, 0)_{\text{rect}}$$

Converting from Rectangular to Spherical Coordinates

$$16. \vec{p} = (1, 1, \sqrt{2})_{\text{rect}} \rightarrow \vec{p} = \left(2, \frac{\pi}{4}, \frac{\pi}{4} \right)_{\text{sphere}}$$

$$17. \vec{p} = \left(\frac{\sqrt{3}}{4}, \frac{1}{2}, \frac{\sqrt{3}}{2} \right)_{\text{rect}} \rightarrow \vec{p} = \left(\frac{\sqrt{19}}{4}, 0.65, 0.86 \right)_{\text{sphere}}$$

$$18. \vec{p} = (1, 1, 0)_{\text{rect}} \rightarrow \vec{p} = \left(\sqrt{2}, \frac{\pi}{2}, \frac{\pi}{4} \right)_{\text{sphere}}$$

Converting from Spherical to Cylindrical Coordinates

$$19. \vec{p} = \left(4, \frac{\pi}{4}, \frac{\pi}{3}\right)_{\text{sphere}} \rightarrow \boxed{\vec{p} = \left(2\sqrt{2}, \frac{\pi}{3}, 2\sqrt{2}\right)_{\text{cyl}}}$$

$$20. \vec{p} = \left(2\sqrt{2}, \frac{\pi}{4}, \frac{\pi}{3}\right)_{\text{sphere}} \rightarrow \boxed{\vec{p} = \left(2, \frac{\pi}{3}, 2\right)_{\text{cyl}}}$$

$$21. \vec{p} = \left(\sqrt{2}, \frac{\pi}{2}, \frac{\pi}{4}\right)_{\text{sphere}} \rightarrow \boxed{\vec{p} = \left(\sqrt{2}, \frac{\pi}{4}, 0\right)_{\text{cyl}}}$$

Converting from Cylindrical to Spherical Coordinates

$$22. \vec{p} = \left(1, \frac{\pi}{2}, 1\right)_{\text{cyl}} \rightarrow \boxed{\vec{p} = \left(\sqrt{2}, \frac{\pi}{4}, \frac{\pi}{2}\right)_{\text{sphere}}}$$

$$23. \vec{p} = \left(\sqrt{6}, \frac{\pi}{4}, \sqrt{2}\right)_{\text{cyl}} \rightarrow \boxed{\vec{p} = \left(2\sqrt{2}, \frac{\pi}{3}, \frac{\pi}{2}\right)_{\text{sphere}}}$$

$$24. \vec{p} = \left(1, \frac{\pi}{4}, 5\right)_{\text{cyl}} \rightarrow \boxed{\vec{p} \approx \left(\sqrt{26}, 0.20, \frac{\pi}{4}\right)_{\text{sphere}}}$$

Gradient of a Scalar Field

$$25. f(x, y, z) = x^2y + xz + y^2 \rightarrow \boxed{\overrightarrow{\text{grad}}(f) = (2xy + z, x^2 + 2y, x)}$$

$$26. f(x, y, z) = x^2 + y^2 + z^2 + 2 \rightarrow \boxed{\overrightarrow{\text{grad}}(f) = (2x, 2y, 2z)}$$

$$27. f(x, y, z) = x + 3y + 5z + 2 \rightarrow \boxed{\overrightarrow{\text{grad}}(f) = (1, 3, 5)}$$

Curl of a Vector Field

$$28. \vec{u} = (x^2y)\hat{i} + (yz)\hat{j} - (z^2)\hat{k} \rightarrow \boxed{\text{curl}(\vec{u}) = -y\hat{i} - x^2\hat{k} = (-y, 0, -x^2)}$$

$$29. \vec{u} = (x^2)\hat{i} + (z^2)\hat{j} - (xy^3)\hat{k} \rightarrow \boxed{\text{curl}(\vec{u}) = -2xy^2\hat{i} + y^3\hat{j} = (-2xy^2, y^3, 0)}$$

$$30. \vec{u} = (x)\hat{i} + (z)\hat{j} - (x)\hat{k} \rightarrow \boxed{\text{curl}(\vec{u}) = -\hat{i} + \hat{j} = (-1, 1, 0)}$$

Divergence of a Vector Field

$$31. \vec{u} = x^2y\hat{i} + zy\hat{j} - z^2\hat{k} \rightarrow \boxed{\operatorname{div} \vec{u} = 2xy - z}$$

$$32. \vec{u} = x^2\hat{i} + z^2\hat{j} - xy^3\hat{k} \rightarrow \boxed{\operatorname{div} \vec{u} = 2x}$$

$$33. \vec{u} = x\hat{i} + z\hat{j} - x\hat{k} \rightarrow \boxed{\operatorname{div} \vec{u} = 1}$$

Laplacian of a Scalar Field

$$34. f(x, y, z) = x^2y + xz + y^2 \rightarrow \boxed{\vec{\nabla}^2 f = 2y + 2}$$

$$35. f(x, y, z) = x^2 + y^2 + z^2 + 2 \rightarrow \boxed{\vec{\nabla}^2 f = 6}$$

$$36. f(x, y, z) = zx + 3x^3y^2 + 2xz^2 \rightarrow \boxed{\vec{\nabla}^2 f = 18xy^2 + 6x^3 + 4x}$$

Laplacian of a Vector Field

$$37. \vec{u} = (3x^2y, zy^2, 3z^2) \rightarrow \boxed{\vec{\nabla}^2 \vec{u} = (6y, 2z, 6)}$$

$$38. \vec{u} = (x^2 + y, x + zy^2, z^2) \rightarrow \boxed{\vec{\nabla}^2 \vec{u} = (2, 2z, 2)}$$

39. Duplicate of #37.

Derivative of a Vector with Respect to a Vector

$$40. \vec{w} = \begin{bmatrix} w_1 \\ w_2 \end{bmatrix}, \begin{cases} w_1 = 2x + 3y \\ w_2 = 7x + 5y \end{cases}, \vec{u} = \begin{bmatrix} x \\ y \end{bmatrix} \rightarrow \boxed{\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} 2 & 3 \\ 7 & 5 \end{bmatrix}}$$

$$41. \vec{w} = \begin{bmatrix} w_1 \\ w_2 \end{bmatrix}, \begin{cases} w_1 = x - y + z \\ w_2 = x + 2y - z \end{cases}, \vec{u} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \rightarrow \boxed{\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} 1 & -1 & 1 \\ 1 & 2 & -1 \end{bmatrix}}$$

$$42. \vec{w} = \begin{bmatrix} w_1 \\ w_2 \\ w_3 \end{bmatrix}, \begin{cases} w_1 = xy + z \\ w_2 = x - y^2 + z \\ w_3 = 2x + y + xz \end{cases}, \vec{u} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \rightarrow \boxed{\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} y & x & 1 \\ 1 & -2y & 1 \\ 2 + z & 1 & x \end{bmatrix}}$$

Derivative of a Scalar with Respect to a Vector

$$43. s(\vec{x}) = (x_1 + 1)^2 + x_2^2 + (x_3 + 2)^2, \vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \rightarrow \frac{\partial s}{\partial \vec{x}} = \begin{bmatrix} 2x_1 + 2 \\ 2x_2 \\ 2x_3 + 4 \end{bmatrix}$$

$$44. f(\vec{x}) = x + y + z, \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \rightarrow \frac{\partial f}{\partial \vec{x}} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$45. g(\vec{x}) = x + xy + z^2, \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \rightarrow \frac{\partial g}{\partial \vec{x}} = \begin{bmatrix} 1 + y \\ x \\ 2z \end{bmatrix}$$

Quadric Forms

$$46. f(x, y) = x^2 + 6xy + 2y^2 \rightarrow \begin{matrix} f(x, y) = \vec{x}^T \cdot \begin{bmatrix} 1 & 3 \\ 3 & 2 \end{bmatrix} \cdot \vec{x} \\ \frac{\partial f(\vec{x})}{\partial \vec{x}} = \begin{bmatrix} 2x + 6y \\ 6x + 4y \end{bmatrix} \end{matrix}$$

$$47. f(x, y) = 5x^2 + 2xy + 2y^2 \rightarrow \begin{matrix} f(x, y) = \vec{x}^T \cdot \begin{bmatrix} 5 & 1 \\ 1 & 2 \end{bmatrix} \cdot \vec{x} \\ \frac{\partial f(\vec{x})}{\partial \vec{x}} = \begin{bmatrix} 10x + 2y \\ 2x + 4y \end{bmatrix} \end{matrix}$$

$$48. f(x, y, z) = 3x^2 + 8xy + 6xz + y^2 + 6yz + 3z^2 \rightarrow \begin{matrix} f(x, y, z) = \vec{x}^T \cdot \begin{bmatrix} 3 & 4 & 3 \\ 4 & 1 & 3 \\ 3 & 3 & 3 \end{bmatrix} \cdot \vec{x} \\ \frac{\partial f(\vec{x})}{\partial \vec{x}} = \begin{bmatrix} 6x + 8y + 6z \\ 8x + 2y + 6z \\ 6x + 6y + 6z \end{bmatrix} \end{matrix}$$

Solutions

Rectangular to Polar Coordinates Conversion

Problem 1

Change $\vec{p} = (3, \sqrt{3})$ from rectangular coordinates to polar coordinates.

$$\vec{p}_{\text{rect}} = (3, \sqrt{3}) = (x, y)$$

$$x = 3$$

$$y = \sqrt{3}$$

$$\vec{p}_{\text{polar}} = (\theta, r)$$

$$y = \sqrt{3} \rightarrow y \geq 0$$

$$\therefore \theta = \cos^{-1}\left(\frac{x}{r}\right)$$

$$\theta = \cos^{-1}\left(\frac{3}{2\sqrt{3}}\right)$$

$$\theta = \frac{\pi}{6}$$

$$r = \sqrt{x^2 + y^2}$$

$$r = \sqrt{(3)^2 + (\sqrt{3})^2}$$

$$r = \sqrt{9 + 3}$$

$$r = \sqrt{12} = 2\sqrt{3}$$

$$\boxed{\vec{p}_{\text{polar}} = \left(\frac{\pi}{6}, 2\sqrt{3}\right)}$$

Problem 2

Change $\vec{p} = (1, \sqrt{3})$ from rectangular coordinates to polar coordinates.

$$\vec{p}_{\text{rect}} = (1, \sqrt{3}) = (x, y)$$

$$x = 1$$

$$y = \sqrt{3}$$

$$\vec{p}_{\text{polar}} = (r, \theta)$$

$$r = \sqrt{x^2 + y^2}$$

$$y = \sqrt{3} \rightarrow y \geq 0$$

$$r = \sqrt{(1)^2 + (\sqrt{3})^2}$$

$$\theta = \cos^{-1} \left(\frac{x}{r} \right)$$

$$r = \sqrt{1 + 3}$$

$$\theta = \cos^{-1} \left(\frac{1}{2} \right)$$

$$r = \sqrt{4} = 2$$

$$\theta = \frac{\pi}{3}$$

$$\boxed{\vec{p}_{\text{polar}} = \left(\frac{\pi}{3}, 2 \right)}$$

Problem 3

Change $\vec{p} = (-1, -1)$ from rectangular coordinates to polar coordinates.

$$\vec{p}_{\text{rect}} = (-1, -1) = (x, y)$$

$$x = -1$$

$$y = -1$$

$$\vec{p}_{\text{polar}} = (\theta, r)$$

$$r = \sqrt{x^2 + y^2}$$

$$y = -1 \rightarrow y < 0$$

$$r = \sqrt{(-1)^2 + (-1)^2}$$

$$\theta = -\cos^{-1}\left(\frac{x}{r}\right)$$

$$r = \sqrt{1 + 1}$$

$$\theta = -\cos^{-1}\left(\frac{-1}{\sqrt{2}}\right)$$

$$r = \sqrt{2}$$

$$\theta = -\frac{3\pi}{4}$$

$$\boxed{\vec{p}_{\text{polar}} = \left(-\frac{3\pi}{4}, \sqrt{2}\right)}$$

Polar to Rectangular Coordinates Conversion

Problem 4

Change $\vec{p} = \left(\frac{7\pi}{4}, 3\sqrt{2}\right)$ from polar coordinates to rectangular coordinates.

$$\vec{p}_{\text{polar}} = \left(\frac{7\pi}{4}, 3\sqrt{2}\right) = (\theta, r)$$

$$r = 3\sqrt{2}$$

$$\theta = \frac{7\pi}{4}$$

$$\vec{p}_{\text{rect}} = (x, y)$$

$$x = r \cos(\theta)$$

$$y = 3\sqrt{2} \cdot \sin\left(\frac{7\pi}{4}\right)$$

$$x = 3\sqrt{2} \cdot \cos\left(\frac{7\pi}{4}\right)$$

$$y = 3\sqrt{2} \cdot \left(-\frac{\sqrt{2}}{2}\right)$$

$$x = 3\sqrt{2} \cdot \frac{\sqrt{2}}{2}$$

$$y = -3$$

$$x = 3$$

$$y = r \sin(\theta)$$

$$\boxed{\vec{p}_{\text{rect}} = (3, -3)}$$

Problem 5

Change $\vec{p} = \left(\frac{\pi}{4}, 4\right)$ from polar coordinates to rectangular coordinates.

$$\vec{p}_{\text{polar}} = \left(\frac{\pi}{4}, 4\right) = (\theta, r)$$

$$r = 4$$

$$\theta = \frac{\pi}{4}$$

$$\vec{p}_{\text{rect}} = (x, y)$$

$$x = r \cos(\theta)$$

$$y = 4 \cdot \sin\left(\frac{\pi}{4}\right)$$

$$x = 4 \cdot \cos\left(\frac{\pi}{4}\right)$$

$$y = 4 \cdot \frac{\sqrt{2}}{2}$$

$$x = 4 \cdot \frac{\sqrt{2}}{2}$$

$$y = 2\sqrt{2}$$

$$x = 2\sqrt{2}$$

$$y = r \sin(\theta)$$

$$\boxed{\vec{p}_{\text{rect}} = (2\sqrt{2}, 2\sqrt{2})}$$

Problem 6

Change $\vec{p} = \left(\frac{\pi}{3}, 1\right)$ from polar coordinates to rectangular coordinates.

$$\vec{p}_{\text{polar}} = \left(\frac{\pi}{3}, 1\right) = (\theta, r)$$

$$r = 1$$

$$\theta = \frac{\pi}{3}$$

$$\vec{p}_{\text{rect}} = (x, y)$$

$$x = r \cos(\theta)$$

$$y = r \sin(\theta)$$

$$x = 1 \cdot \cos\left(\frac{\pi}{3}\right)$$

$$y = 1 \cdot \sin\left(\frac{\pi}{3}\right)$$

$$x = 1 \cdot \frac{1}{2}$$

$$y = 1 \cdot \frac{\sqrt{3}}{2}$$

$$x = \frac{1}{2}$$

$$y = \frac{\sqrt{3}}{2}$$

$$\boxed{\vec{p}_{\text{rect}} = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)}$$

Cylindrical to Rectangular Coordinates Conversion

Problem 7

Change $\vec{p} = \left(3, \frac{\pi}{2}, 1\right)$ from cylindrical coordinates to rectangular coordinates.

$$\vec{p}_{\text{cyl}} = \left(3, \frac{\pi}{2}, 1\right) = (\rho, \phi, z)$$

$$\rho = 3$$

$$\phi = \frac{\pi}{2}$$

$$z = 1$$

$$\vec{p}_{\text{rect}} = (x, y, z)$$

$$x = \rho \cos(\phi)$$

$$y = \rho \sin(\phi)$$

$$z = z$$

$$x = 3 \cos\left(\frac{\pi}{2}\right)$$

$$y = 3 \sin\left(\frac{\pi}{2}\right)$$

$$z = 1$$

$$x = 3 \cdot 0$$

$$y = 3 \cdot 1$$

$$x = 0$$

$$y = 3$$

$$\boxed{\vec{p}_{\text{rect}} = (0, 3, 1)}$$

Problem 8

Change $\vec{p} = \left(4, \frac{\pi}{6}, 2\right)$ from cylindrical coordinates to rectangular coordinates.

$$\vec{p}_{\text{cyl}} = \left(4, \frac{\pi}{6}, 2\right) = (\rho, \phi, z)$$

$$\rho = 4$$

$$\phi = \frac{\pi}{6}$$

$$z = 2$$

$$\vec{p}_{\text{rect}} = (x, y, z)$$

$$x = \rho \cos(\phi)$$

$$y = \rho \sin(\phi)$$

$$z = z$$

$$x = 4 \cos\left(\frac{\pi}{6}\right)$$

$$y = 4 \sin\left(\frac{\pi}{6}\right)$$

$$z = 2$$

$$x = 4 \cdot \frac{\sqrt{3}}{2}$$

$$y = 4 \cdot \frac{1}{2}$$

$$x = 2\sqrt{3}$$

$$y = 2$$

$$\boxed{\vec{p}_{\text{rect}} = (2\sqrt{3}, 2, 2)}$$

Problem 9

Change $\vec{p} = \left(1, \frac{\pi}{4}, 5\right)$ from cylindrical coordinates to rectangular coordinates.

$$\vec{p}_{\text{cyl}} = \left(1, \frac{\pi}{4}, 5\right) = (\rho, \phi, z)$$

$$\rho = 1$$

$$\phi = \frac{\pi}{4}$$

$$z = 5$$

$$\vec{p}_{\text{rect}} = (x, y, z)$$

$$x = \rho \cos(\phi)$$

$$y = \rho \sin(\phi)$$

$$z = z$$

$$x = 1 \cos\left(\frac{\pi}{4}\right)$$

$$y = 1 \sin\left(\frac{\pi}{4}\right)$$

$$z = 5$$

$$x = 1 \cdot \frac{\sqrt{2}}{2}$$

$$y = 1 \cdot \frac{\sqrt{2}}{2}$$

$$x = \frac{\sqrt{2}}{2}$$

$$y = \frac{\sqrt{2}}{2}$$

$$\boxed{\vec{p}_{\text{rect}} = \left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}, 5\right)}$$

Rectangular to Cylindrical Coordinates Conversion

Problem 10

Change $\vec{p} = (1,1,1)$ from rectangular coordinates to cylindrical coordinates.

$$\vec{p}_{\text{rect}} = (1,1,1) = (x, y, z)$$

$$x = 1$$

$$y = 1$$

$$z = 1$$

$$\vec{p}_{\text{cyl}} = (\rho, \phi, z)$$

$$\rho = \sqrt{x^2 + y^2}$$

$$x = 1 \rightarrow x \geq 0$$

$$z = z$$

$$\rho = \sqrt{(1)^2 + (1)^2}$$

$$\phi = \sin^{-1}\left(\frac{y}{\rho}\right)$$

$$z = 1$$

$$\rho = \sqrt{1+1}$$

$$\phi = \sin^{-1}\left(\frac{1}{\sqrt{2}}\right)$$

$$\rho = \sqrt{2}$$

$$\phi = \frac{\pi}{4}$$

$$\boxed{\vec{p}_{\text{cyl}} = \left(\sqrt{2}, \frac{\pi}{4}, 1\right)}$$

Problem 11

Change $\vec{p} = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}, 5\right)$ from rectangular coordinates to cylindrical coordinates.

$$\vec{p}_{\text{rect}} = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}, 5\right) = (x, y, z)$$

$$x = \frac{1}{2}$$

$$y = \frac{\sqrt{3}}{2}$$

$$z = 5$$

$$\vec{p}_{\text{cyl}} = (\rho, \phi, z)$$

$$\rho = \sqrt{x^2 + y^2}$$

$$x = \frac{1}{2} \rightarrow x \geq 0$$

$$z = z$$

$$\rho = \sqrt{\left(\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2}$$

$$\phi = \sin^{-1}\left(\frac{y}{\rho}\right)$$

$$z = 5$$

$$\rho = \sqrt{\frac{1}{4} + \frac{3}{4}}$$

$$\phi = \sin^{-1}\left(\frac{\sqrt{3}/2}{1}\right)$$

$$\rho = \sqrt{1} = 1$$

$$\phi = \sin^{-1}\left(\frac{\sqrt{3}}{2}\right)$$

$$\phi = \frac{\pi}{3}$$

$$\boxed{\vec{p}_{\text{cyl}} = \left(1, \frac{\pi}{3}, 5\right)}$$

Problem 12

Change $\vec{p} = (\sqrt{2}, \sqrt{2}, 3)$ from rectangular coordinates to cylindrical coordinates.

$$\vec{p}_{\text{rect}} = (\sqrt{2}, \sqrt{2}, 3) = (x, y, z)$$

$$x = \sqrt{2}$$

$$y = \sqrt{2}$$

$$z = 3$$

$$\vec{p}_{\text{cyl}} = (\rho, \phi, z)$$

$$\rho = \sqrt{x^2 + y^2}$$

$$x = \sqrt{2} \rightarrow x \geq 0$$

$$z = z$$

$$\rho = \sqrt{(\sqrt{2})^2 + (\sqrt{2})^2}$$

$$\phi = \sin^{-1}\left(\frac{y}{\rho}\right)$$

$$z = 3$$

$$\rho = \sqrt{2 + 2}$$

$$\phi = \sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$$

$$\rho = \sqrt{4} = 2$$

$$\phi = \frac{\pi}{4}$$

$$\boxed{\vec{p}_{\text{cyl}} = \left(2, \frac{\pi}{4}, 3\right)}$$

Spherical to Rectangular Coordinates Conversion

Problem 13

Change $\vec{p} = \left(1, \frac{\pi}{4}, \pi\right)$ from spherical coordinates to rectangular coordinates.

$$\vec{p}_{\text{sphere}} = \left(1, \frac{\pi}{4}, \pi\right) = (r, \theta, \phi)$$

$$r = 1$$

$$\phi = \frac{\pi}{4}$$

$$\theta = \pi$$

$$\vec{p}_{\text{rect}} = (x, y, z)$$

$$x = r \sin(\theta) \cos(\phi)$$

$$y = r \sin(\theta) \sin(\phi)$$

$$z = r \cos(\theta)$$

$$x = 1 \sin(\pi) \cos\left(\frac{\pi}{4}\right)$$

$$y = 1 \sin(\pi) \sin\left(\frac{\pi}{4}\right)$$

$$z = 1 \cos(\pi)$$

$$x = 1 \cdot 0 \cdot \frac{\sqrt{2}}{2}$$

$$y = 1 \cdot 0 \cdot \frac{\sqrt{2}}{2}$$

$$z = 1 \cdot (-1)$$

$$z = -1$$

$$x = 0$$

$$y = 0$$

$$\boxed{\vec{p}_{\text{rect}} = (0, 0, -1)}$$

Problem 14

Change $\vec{p} = \left(3, \frac{\pi}{3}, \frac{\pi}{4}\right)$ from spherical coordinates to rectangular coordinates.

$$\vec{p}_{\text{sphere}} = \left(3, \frac{\pi}{3}, \frac{\pi}{4}\right) = (r, \theta, \phi)$$

$$r = 3$$

$$\theta = \frac{\pi}{3}$$

$$\phi = \frac{\pi}{4}$$

$$\vec{p}_{\text{rect}} = (x, y, z)$$

$$x = r \sin(\theta) \cos(\phi)$$

$$y = r \sin(\theta) \sin(\phi)$$

$$z = r \cos(\theta)$$

$$x = 3 \sin\left(\frac{\pi}{3}\right) \cos\left(\frac{\pi}{4}\right)$$

$$y = 3 \sin\left(\frac{\pi}{3}\right) \sin\left(\frac{\pi}{4}\right)$$

$$z = 3 \cos\left(\frac{\pi}{3}\right)$$

$$x = 3 \cdot \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2}$$

$$y = 3 \cdot \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2}$$

$$z = 3 \cdot \frac{1}{2}$$

$$x = \frac{3\sqrt{6}}{4}$$

$$y = \frac{3\sqrt{6}}{4}$$

$$z = \frac{3}{2}$$

$$\boxed{\vec{p}_{\text{rect}} = \left(\frac{3\sqrt{6}}{4}, \frac{3\sqrt{6}}{4}, \frac{3}{2}\right)}$$

Problem 15

Change $\vec{p} = \left(5, \frac{\pi}{2}, \pi\right)$ from spherical coordinates to rectangular coordinates.

$$\vec{p}_{\text{sphere}} = \left(5, \frac{\pi}{2}, \pi\right) = (r, \theta, \phi)$$

$$r = 5$$

$$\theta = \frac{\pi}{2}$$

$$\phi = \pi$$

$$\vec{p}_{\text{rect}} = (x, y, z)$$

$$x = r \sin(\theta) \cos(\phi)$$

$$y = r \sin(\theta) \sin(\phi)$$

$$z = r \cos(\theta)$$

$$x = 5 \sin\left(\frac{\pi}{2}\right) \cos(\pi)$$

$$y = 5 \sin\left(\frac{\pi}{2}\right) \sin(\pi)$$

$$z = 5 \cos\left(\frac{\pi}{2}\right)$$

$$x = 5 \cdot 1 \cdot 0$$

$$y = 5 \cdot 1 \cdot (-1)$$

$$z = 5 \cdot (0)$$

$$x = 0$$

$$y = -5$$

$$z = 0$$

$$\boxed{\vec{p}_{\text{rect}} = (0, -5, 0)}$$

Rectangular to Spherical Coordinates Conversion

Problem 16

Change $\vec{p} = (1, 1, \sqrt{2})$ from rectangular coordinates to spherical coordinates.

$$\vec{p}_{\text{rect}} = (1, 1, \sqrt{2}) = (x, y, z)$$

$$x = 1$$

$$y = 1$$

$$z = \sqrt{2}$$

$$\vec{p}_{\text{sphere}} = (r, \theta, \phi)$$

$$r = \sqrt{x^2 + y^2 + z^2}$$

$$\theta = \cos^{-1} \left(\frac{z}{r} \right)$$

$$\phi = \tan^{-1} \left(\frac{y}{x} \right)$$

$$r = \sqrt{(1)^2 + (1)^2 + (\sqrt{2})^2}$$

$$\theta = \cos^{-1} \left(\frac{\sqrt{2}}{2} \right)$$

$$\phi = \tan^{-1} \left(\frac{1}{1} \right)$$

$$r = \sqrt{1 + 1 + 2}$$

$$r = \sqrt{4}$$

$$\theta = \frac{\pi}{4}$$

$$\phi = \tan^{-1}(1)$$

$$r = 2$$

$$\phi = \frac{\pi}{4}$$

$$\boxed{\vec{p}_{\text{sphere}} = \left(2, \frac{\pi}{4}, \frac{\pi}{4} \right)}$$

Problem 17

Change $\vec{p} = \left(\frac{\sqrt{3}}{4}, \frac{1}{2}, \frac{\sqrt{3}}{2}\right)$ from rectangular coordinates to spherical coordinates.

$$\vec{p}_{\text{rect}} = \left(\frac{\sqrt{3}}{4}, \frac{1}{2}, \frac{\sqrt{3}}{2}\right) = (x, y, z)$$

$$x = \frac{\sqrt{3}}{4}$$

$$y = \frac{1}{2}$$

$$z = \frac{\sqrt{3}}{2}$$

$$\vec{p}_{\text{sphere}} = (r, \theta, \phi)$$

$$r = \sqrt{x^2 + y^2 + z^2}$$

$$\theta = \cos^{-1} \left(\frac{z}{r} \right)$$

$$\phi = \tan^{-1} \left(\frac{y}{x} \right)$$

$$r = \sqrt{\left(\frac{\sqrt{3}}{4}\right)^2 + \left(\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2}$$

$$\theta = \cos^{-1} \left(\frac{\frac{\sqrt{3}}{2}}{\frac{\sqrt{19}}{4}} \right)$$

$$\phi = \tan^{-1} \left(\frac{1/2}{\sqrt{3}/4} \right)$$

$$r = \sqrt{\frac{3}{16} + \frac{1}{4} + \frac{3}{4}}$$

$$\theta = \cos^{-1} \left(\frac{2\sqrt{57}}{19} \right)$$

$$\phi = \tan^{-1} \left(\frac{2\sqrt{3}}{3} \right)$$

$$r = \sqrt{\frac{3}{16} + \frac{4}{16} + \frac{12}{16}}$$

$$\theta \approx 0.65$$

$$\phi \approx 0.86$$

$$r = \sqrt{\frac{19}{16}}$$

$$r = \frac{\sqrt{19}}{4}$$

$$\boxed{\vec{p}_{\text{sphere}} \approx \left(\frac{\sqrt{19}}{4}, 0.65, 0.86\right)}$$

Problem 18

Change $\vec{p} = (1,1,0)$ from rectangular coordinates to spherical coordinates.

$$\vec{p}_{\text{rect}} = (1,1,0) = (x,y,z)$$

$$x = 1$$

$$y = 1$$

$$z = 0$$

$$\vec{p}_{\text{sphere}} = (r, \theta, \phi)$$

$$r = \sqrt{x^2 + y^2 + z^2}$$

$$\theta = \cos^{-1} \left(\frac{z}{r} \right)$$

$$\phi = \tan^{-1} \left(\frac{y}{x} \right)$$

$$r = \sqrt{(1)^2 + (1)^2 + (0)^2}$$

$$\theta = \cos^{-1} \left(\frac{0}{\sqrt{2}} \right)$$

$$\phi = \tan^{-1} \left(\frac{1}{1} \right)$$

$$r = \sqrt{1 + 1 + 0}$$

$$\theta = \cos^{-1}(0)$$

$$\phi = \tan^{-1}(1)$$

$$r = \sqrt{2}$$

$$\theta = \frac{\pi}{2}$$

$$\phi = \frac{\pi}{4}$$

$$\boxed{\vec{p}_{\text{sphere}} = \left(\sqrt{2}, \frac{\pi}{2}, \frac{\pi}{4} \right)}$$

Spherical to Cylindrical Coordinates Conversion

Problem 19

Change $\vec{p} = \left(4, \frac{\pi}{4}, \frac{\pi}{3}\right)$ from spherical coordinates to cylindrical coordinates.

$$\vec{p}_{\text{sphere}} = \left(4, \frac{\pi}{4}, \frac{\pi}{3}\right) = (r, \theta, \phi)$$

$$r = 4$$

$$\theta = \frac{\pi}{4}$$

$$\phi = \frac{\pi}{3}$$

$$p_{\text{cyl}} = (\rho, \phi, z)$$

$$\rho = r \sin(\theta)$$

$$\phi = \phi$$

$$z = r \cos(\theta)$$

$$\rho = (4) \sin\left(\frac{\pi}{4}\right)$$

$$\phi = \frac{\pi}{3}$$

$$z = (4) \cos\left(\frac{\pi}{4}\right)$$

$$\rho = (4) \left(\frac{\sqrt{2}}{2}\right)$$

$$z = (4) \left(\frac{\sqrt{2}}{2}\right)$$

$$\rho = 2\sqrt{2}$$

$$z = 2\sqrt{2}$$

$$\boxed{p_{\text{cyl}} = \left(2\sqrt{2}, \frac{\pi}{3}, 2\sqrt{2}\right)}$$

Problem 20

Change $\vec{p} = \left(2\sqrt{2}, \frac{\pi}{4}, \frac{\pi}{3}\right)$ from spherical coordinates to cylindrical coordinates.

$$\vec{p}_{\text{sphere}} = \left(2\sqrt{2}, \frac{\pi}{4}, \frac{\pi}{3}\right) = (r, \theta, \phi)$$

$$r = 2\sqrt{2}$$

$$\theta = \frac{\pi}{4}$$

$$\phi = \frac{\pi}{3}$$

$$p_{\text{cyl}} = (\rho, \phi, z)$$

$$\rho = r \sin(\theta)$$

$$\phi = \phi$$

$$z = r \cos(\theta)$$

$$\rho = (2\sqrt{2}) \sin\left(\frac{\pi}{4}\right)$$

$$\phi = \frac{\pi}{3}$$

$$z = (2\sqrt{2}) \cos\left(\frac{\pi}{4}\right)$$

$$\rho = (2\sqrt{2}) \left(\frac{\sqrt{2}}{2}\right)$$

$$z = (2\sqrt{2}) \left(\frac{\sqrt{2}}{2}\right)$$

$$\rho = 2$$

$$z = 2$$

$$\boxed{p_{\text{cyl}} = \left(2, \frac{\pi}{3}, 2\right)}$$

Problem 21

Change $\vec{p} = \left(\sqrt{2}, \frac{\pi}{2}, \frac{\pi}{4}\right)$ from spherical coordinates to cylindrical coordinates.

$$\vec{p}_{\text{sphere}} = \left(\sqrt{2}, \frac{\pi}{2}, \frac{\pi}{4}\right) = (r, \theta, \phi)$$

$$r = \sqrt{2}$$

$$\theta = \frac{\pi}{2}$$

$$\phi = \frac{\pi}{4}$$

$$p_{\text{cyl}} = (\rho, \phi, z)$$

$$\rho = r \sin(\theta)$$

$$\phi = \phi$$

$$z = r \cos(\theta)$$

$$\rho = (\sqrt{2}) \sin\left(\frac{\pi}{2}\right)$$

$$\phi = \frac{\pi}{4}$$

$$z = (\sqrt{2}) \cos\left(\frac{\pi}{2}\right)$$

$$\rho = (\sqrt{2})(1)$$

$$z = (2\sqrt{2})(0)$$

$$\rho = \sqrt{2}$$

$$z = 0$$

$$\boxed{p_{\text{cyl}} = \left(\sqrt{2}, \frac{\pi}{4}, 0\right)}$$

Cylindrical to Spherical Coordinates Conversion

Problem 22

Change $\vec{p} = \left(1, \frac{\pi}{2}, 1\right)$ from cylindrical coordinates to spherical coordinates.

$$\vec{p}_{\text{cyl}} = \left(1, \frac{\pi}{2}, 1\right) = (\rho, \phi, z)$$

$$\rho = 1$$

$$\phi = \frac{\pi}{2}$$

$$z = 1$$

$$\vec{p}_{\text{sphere}} = (r, \theta, \phi)$$

$$r = \sqrt{\rho^2 + z^2}$$

$$\theta = \tan^{-1}\left(\frac{\rho}{z}\right)$$

$$\phi = \phi$$

$$r = \sqrt{(1)^2 + (1)^2}$$

$$\theta = \tan^{-1}\left(\frac{1}{1}\right)$$

$$\phi = \frac{\pi}{2}$$

$$r = \sqrt{1 + 1}$$

$$\theta = \tan^{-1}(1)$$

$$r = \sqrt{2}$$

$$\theta = \frac{\pi}{4}$$

$$\boxed{\vec{p}_{\text{sphere}} = \left(\sqrt{2}, \frac{\pi}{4}, \frac{\pi}{2}\right)}$$

Problem 23

Change $\vec{p} = \left(\sqrt{6}, \frac{\pi}{4}, \sqrt{2}\right)$ from cylindrical coordinates to spherical coordinates.

$$\vec{p}_{\text{cyl}} = \left(\sqrt{6}, \frac{\pi}{4}, \sqrt{2}\right) = (\rho, \phi, z)$$

$$\rho = \sqrt{6}$$

$$\phi = \frac{\pi}{4}$$

$$z = \sqrt{2}$$

$$\vec{p}_{\text{sphere}} = (r, \theta, \phi)$$

$$r = \sqrt{\rho^2 + z^2}$$

$$\theta = \tan^{-1}\left(\frac{\rho}{z}\right)$$

$$\phi = \phi$$

$$r = \sqrt{(\sqrt{6})^2 + (\sqrt{2})^2}$$

$$\theta = \tan^{-1}\left(\frac{\sqrt{6}}{\sqrt{2}}\right)$$

$$\phi = \frac{\pi}{2}$$

$$r = \sqrt{6 + 2}$$

$$\theta = \tan^{-1}(\sqrt{3})$$

$$r = \sqrt{8} = 2\sqrt{2}$$

$$\theta = \frac{\pi}{3}$$

$$\boxed{\vec{p}_{\text{sphere}} = \left(2\sqrt{2}, \frac{\pi}{3}, \frac{\pi}{2}\right)}$$

Problem 24

Change $\vec{p} = \left(1, \frac{\pi}{4}, 5\right)$ from cylindrical coordinates to spherical coordinates.

$$\vec{p}_{\text{cyl}} = \left(1, \frac{\pi}{4}, 5\right) = (\rho, \phi, z)$$

$$\rho = 1$$

$$\phi = \frac{\pi}{4}$$

$$z = 5$$

$$\vec{p}_{\text{sphere}} = (r, \theta, \phi)$$

$$r = \sqrt{\rho^2 + z^2}$$

$$\theta = \tan^{-1}\left(\frac{\rho}{z}\right)$$

$$\phi = \phi$$

$$r = \sqrt{(1)^2 + (5)^2}$$

$$\theta = \tan^{-1}\left(\frac{1}{5}\right)$$

$$\phi = \frac{\pi}{4}$$

$$r = \sqrt{1 + 25}$$

$$\theta \approx 0.20$$

$$r = \sqrt{26}$$

$$\boxed{\vec{p}_{\text{sphere}} \approx \left(\sqrt{26}, 0.20, \frac{\pi}{4}\right)}$$

Gradient of a Scalar Field

Problem 25

Given the scalar field $f(x, y, z) = x^2y + xz + y^2$, calculate $\overrightarrow{\text{grad}}(f) = \vec{\nabla}f$.

$$\overrightarrow{\text{grad}}(f) = \vec{\nabla}f$$

$$\overrightarrow{\text{grad}}(f) = \frac{\partial f}{\partial x}\hat{i} + \frac{\partial f}{\partial y}\hat{j} + \frac{\partial f}{\partial z}\hat{k}$$

$$\overrightarrow{\text{grad}}(f) = \frac{\partial}{\partial x}(x^2y + xz + y^2) \cdot \hat{i} + \frac{\partial}{\partial y}(x^2y + xz + y^2) \cdot \hat{j} + \frac{\partial}{\partial z}(x^2y + xz + y^2) \cdot \hat{k}$$

$$\overrightarrow{\text{grad}}(f) = (2xy + z + 0) \cdot \hat{i} + (x^2 + 0 + 2y) \cdot \hat{j} + (0 + x + 0) \cdot \hat{k}$$

$$\overrightarrow{\text{grad}}(f) = (2xy + z)\hat{i} + (x^2 + 2y)\hat{j} + x\hat{k}$$

$$\boxed{\overrightarrow{\text{grad}}(f) = (2xy + z, x^2 + 2y, x)}$$

Problem 26

Given the scalar field $f(x, y, z) = x^2 + y^2 + z^2 + 2$, calculate $\overrightarrow{\text{grad}}(f) = \vec{\nabla}f$.

$$\overrightarrow{\text{grad}}(f) = \vec{\nabla}f$$

$$\overrightarrow{\text{grad}}(f) = \frac{\partial f}{\partial x}\hat{i} + \frac{\partial f}{\partial y}\hat{j} + \frac{\partial f}{\partial z}\hat{k}$$

$$\overrightarrow{\text{grad}}(f) = \frac{\partial}{\partial x}(x^2 + y^2 + z^2 + 2) \cdot \hat{i} + \frac{\partial}{\partial y}(x^2 + y^2 + z^2 + 2) \cdot \hat{j}$$

$$+ \frac{\partial}{\partial z}(x^2 + y^2 + z^2 + 2) \cdot \hat{k}$$

$$\overrightarrow{\text{grad}}(f) = (2x + 0 + 0 + 0) \cdot \hat{i} + (0 + 2y + 0 + 0) \cdot \hat{j} + (0 + 0 + 2z + 0) \cdot \hat{k}$$

$$\overrightarrow{\text{grad}}(f) = 2x\hat{i} + 2y\hat{j} + 2z\hat{k}$$

$$\boxed{\overrightarrow{\text{grad}}(f) = (2x, 2y, 2z)}$$

Problem 27

Given the scalar field $f(x, y, z) = x + 3y + 5z + 2$, calculate $\overrightarrow{\text{grad}}(f) = \vec{\nabla}f$.

$$\overrightarrow{\text{grad}}(f) = \vec{\nabla}f$$

$$\overrightarrow{\text{grad}}(f) = \frac{\partial f}{\partial x} \hat{i} + \frac{\partial f}{\partial y} \hat{j} + \frac{\partial f}{\partial z} \hat{k}$$

$$\begin{aligned} \overrightarrow{\text{grad}}(f) &= \frac{\partial}{\partial x}(x + 3y + 5z + 2) \cdot \hat{i} + \frac{\partial}{\partial y}(x + 3y + 5z + 2) \cdot \hat{j} \\ &\quad + \frac{\partial}{\partial z}(x + 3y + 5z + 2) \cdot \hat{k} \end{aligned}$$

$$\overrightarrow{\text{grad}}(f) = (1 + 0 + 0 + 0) \cdot \hat{i} + (0 + 3 + 0 + 0) \cdot \hat{j} + (0 + 0 + 5 + 0) \cdot \hat{k}$$

$$\overrightarrow{\text{grad}}(f) = \hat{i} + 3\hat{j} + 5\hat{k}$$

$$\boxed{\overrightarrow{\text{grad}}(f) = (1, 3, 5)}$$

Curl of a Vector Field

Problem 28

Given the vector field $\vec{u} = (u_x, u_y, u_z) = (x^2y)\hat{i} + (yz)\hat{j} - (z^2)\hat{k}$, calculate

$$\text{curl}(\vec{u}) = \vec{\nabla} \times \vec{u}.$$

$$\text{curl}(\vec{u}) = \vec{\nabla} \times \vec{u}$$

$$\text{curl}(\vec{u}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ u_x & u_y & u_z \end{vmatrix}$$

$$\text{curl}(\vec{u}) = \begin{vmatrix} \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ u_y & u_z \end{vmatrix} \hat{i} - \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial z} \\ u_x & u_z \end{vmatrix} \hat{j} + \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \\ u_x & u_y \end{vmatrix} \hat{k}$$

$$\text{curl}(\vec{u}) = \left(\frac{\partial u_z}{\partial y} - \frac{\partial u_y}{\partial z} \right) \hat{i} - \left(\frac{\partial u_z}{\partial x} - \frac{\partial u_x}{\partial z} \right) \hat{j} + \left(\frac{\partial u_y}{\partial x} - \frac{\partial u_x}{\partial y} \right) \hat{k}$$

$$\text{curl}(\vec{u}) = \left[\frac{\partial}{\partial y}(-z^2) - \frac{\partial}{\partial z}(yz) \right] \hat{i} - \left[\frac{\partial}{\partial x}(-z^2) - \frac{\partial}{\partial z}(x^2y) \right] \hat{j} + \left[\frac{\partial}{\partial x}(yz) - \frac{\partial}{\partial y}(x^2y) \right] \hat{k}$$

$$\text{curl}(\vec{u}) = (0 - y)\hat{i} - (0 - 0)\hat{j} + (0 - x^2)\hat{k}$$

$$\text{curl}(\vec{u}) = (-y)\hat{i} - (0)\hat{j} + (-x^2)\hat{k}$$

$$\boxed{\text{curl}(\vec{u}) = (-y, 0, -x^2)}$$

Problem 29

Given the vector field $\vec{u} = (u_x, u_y, u_z) = (x^2)\hat{i} + (z^2)\hat{j} - (xy^3)\hat{k}$, calculate

$$\text{curl}(\vec{u}) = \vec{\nabla} \times \vec{u}.$$

$$\vec{u} = (x^2)\hat{i} + (z^2)\hat{j} + (-xy^3)\hat{k}$$

$$\text{curl}(\vec{u}) = \vec{\nabla} \times \vec{u}$$

$$\text{curl}(\vec{u}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ u_x & u_y & u_z \end{vmatrix}$$

$$\text{curl}(\vec{u}) = \begin{vmatrix} \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ u_y & u_z \end{vmatrix} \hat{i} - \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial z} \\ u_x & u_z \end{vmatrix} \hat{j} + \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \\ u_x & u_y \end{vmatrix} \hat{k}$$

$$\text{curl}(\vec{u}) = \left(\frac{\partial u_z}{\partial y} - \frac{\partial u_y}{\partial z} \right) \hat{i} - \left(\frac{\partial u_z}{\partial x} - \frac{\partial u_x}{\partial z} \right) \hat{j} + \left(\frac{\partial u_y}{\partial x} - \frac{\partial u_x}{\partial y} \right) \hat{k}$$

$$\text{curl}(\vec{u}) = \left[\frac{\partial}{\partial y}(-xy^3) - \frac{\partial}{\partial z}(z^2) \right] \hat{i} - \left[\frac{\partial}{\partial x}(-xy^3) - \frac{\partial}{\partial z}(x^2) \right] \hat{j} + \left[\frac{\partial}{\partial x}(z^2) - \frac{\partial}{\partial y}(x^2) \right] \hat{k}$$

$$\text{curl}(\vec{u}) = (-2xy^2 - 0)\hat{i} - (-y^3 - 0)\hat{j} + (0 - 0)\hat{k}$$

$$\text{curl}(\vec{u}) = (-2xy^2)\hat{i} - (-y^3)\hat{j} + (0)\hat{k}$$

$$\text{curl}(\vec{u}) = (-2xy^2)\hat{i} + (y^3)\hat{j} + (0)\hat{k}$$

$$\boxed{\text{curl}(\vec{u}) = (-2xy^2, y^3, 0)}$$

Problem 30

Given the vector field $\vec{u} = (u_x, u_y, u_z) = (x)\hat{i} + (z)\hat{j} - (x)\hat{k}$, calculate $\text{curl}(\vec{u}) = \vec{\nabla} \times \vec{u}$.

$$\vec{u} = (x)\hat{i} + (z)\hat{j} + (-x)\hat{k}$$

$$\text{curl}(\vec{u}) = \vec{\nabla} \times \vec{u}$$

$$\text{curl}(\vec{u}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ u_x & u_y & u_z \end{vmatrix}$$

$$\text{curl}(\vec{u}) = \begin{vmatrix} \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ u_y & u_z \end{vmatrix} \hat{i} - \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial z} \\ u_x & u_z \end{vmatrix} \hat{j} + \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \\ u_x & u_y \end{vmatrix} \hat{k}$$

$$\text{curl}(\vec{u}) = \left(\frac{\partial}{\partial y} u_z - \frac{\partial}{\partial z} u_y \right) \hat{i} - \left(\frac{\partial}{\partial x} u_z - \frac{\partial}{\partial z} u_x \right) \hat{j} + \left(\frac{\partial}{\partial x} u_y - \frac{\partial}{\partial y} u_x \right) \hat{k}$$

$$\text{curl}(\vec{u}) = \left[\frac{\partial}{\partial y} (-x) - \frac{\partial}{\partial z} (z) \right] \hat{i} - \left[\frac{\partial}{\partial x} (-x) - \frac{\partial}{\partial z} (x) \right] \hat{j} + \left[\frac{\partial}{\partial x} (z) - \frac{\partial}{\partial y} (x) \right] \hat{k}$$

$$\text{curl}(\vec{u}) = (0 - 1)\hat{i} - (-1 - 0)\hat{j} + (0 - 0)\hat{k}$$

$$\text{curl}(\vec{u}) = (-1)\hat{i} - (-1)\hat{j} + (0)\hat{k}$$

$$\text{curl}(\vec{u}) = (-1)\hat{i} + (1)\hat{j} + (0)\hat{k}$$

$$\boxed{\text{curl}(\vec{u}) = (-1, 1, 0)}$$

Computing the Divergence of a Vector Field

Problem 31

Given the vector field $\vec{u} = (u_x, u_y, u_z) = x^2y\hat{i} + zy\hat{j} - z^2\hat{k}$, calculate $\text{div } \vec{u} = \vec{\nabla} \cdot \vec{u}$.

$$\vec{u} = (x^2y)\hat{i} + (zy)\hat{j} + (-z^2)\hat{k}$$

$$\text{div } \vec{u} = \vec{\nabla} \cdot \vec{u}$$

$$\text{div } \vec{u} = \frac{\partial u_x}{\partial x} + \frac{\partial u_y}{\partial y} + \frac{\partial u_z}{\partial z}$$

$$\text{div } \vec{u} = \frac{\partial}{\partial x}(x^2y) + \frac{\partial}{\partial y}(zy) + \frac{\partial}{\partial z}(-z^2)$$

$$\text{div } \vec{u} = 2xy + z + (-2z)$$

$$\boxed{\text{div } \vec{u} = 2xy - z}$$

Problem 32

Given the vector field $\vec{u} = (u_x, u_y, u_z) = x^2\hat{i} + z^2\hat{j} - xy^3\hat{k}$, calculate $\text{div } \vec{u} = \vec{\nabla} \cdot \vec{u}$.

$$\vec{u} = (x^2)\hat{i} + (z^2)\hat{j} + (-xy^3)\hat{k}$$

$$\text{div } \vec{u} = \vec{\nabla} \cdot \vec{u}$$

$$\text{div } \vec{u} = \frac{\partial u_x}{\partial x} + \frac{\partial u_y}{\partial y} + \frac{\partial u_z}{\partial z}$$

$$\text{div } \vec{u} = \frac{\partial}{\partial x}(x^2) + \frac{\partial}{\partial y}(z^2) + \frac{\partial}{\partial z}(-xy^3)$$

$$\text{div } \vec{u} = 2x + 0 + 0$$

$$\boxed{\text{div } \vec{u} = 2x}$$

Problem 33

Given the vector field $\vec{u} = (u_x, u_y, u_z) = x\hat{i} + z\hat{j} - x\hat{k}$, calculate $\text{div } \vec{u} = \vec{\nabla} \cdot \vec{u}$.

$$\vec{u} = (x)\hat{i} + (z)\hat{j} + (-x)\hat{k}$$

$$\text{div } \vec{u} = \vec{\nabla} \cdot \vec{u}$$

$$\text{div } \vec{u} = \frac{\partial u_x}{\partial x} + \frac{\partial u_y}{\partial y} + \frac{\partial u_z}{\partial z}$$

$$\text{div } \vec{u} = \frac{\partial}{\partial x}(x) + \frac{\partial}{\partial y}(z) + \frac{\partial}{\partial z}(-x)$$

$$\text{div } \vec{u} = 1 + 0 + 0$$

$$\boxed{\text{div } \vec{u} = 1}$$

Computing the Laplacian of a Scalar Field

Problem 34

Given the scalar field $f(x, y, z) = x^2y + xz + y^2$, compute the Laplacian

$$\vec{\nabla}^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}.$$

$$\vec{\nabla}^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right)$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} (f) \right]$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} (x^2y + xz + y^2) \right]$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} (2xy + z + 0)$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} (2xy + z)$$

$$\frac{\partial^2 f}{\partial x^2} = 2y + 0$$

$$\frac{\partial^2 f}{\partial x^2} = 2y$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} \left[\frac{\partial}{\partial y} (x^2y + xz + y^2) \right]$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} (x^2 + 0 + 2y)$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} (x^2 + 2y)$$

$$\frac{\partial^2 f}{\partial y^2} = 0 + 2$$

$$\frac{\partial^2 f}{\partial y^2} = 2$$

$$\frac{\partial^2 f}{\partial z^2} = \frac{\partial}{\partial z} \left[\frac{\partial}{\partial z} (x^2y + xz + y^2) \right]$$

$$\frac{\partial^2 f}{\partial z^2} = \frac{\partial}{\partial z} (0 + x + 0)$$

$$\frac{\partial^2 f}{\partial z^2} = \frac{\partial}{\partial z} (x)$$

$$\frac{\partial^2 f}{\partial z^2} = 0$$

$$\vec{\nabla}^2 f = 2y + 2 + 0$$

$$\boxed{\vec{\nabla}^2 f = 2y + 2}$$

Problem 35

Given the scalar field $f(x, y, z) = x^2 + y^2 + z^2 + 2$, compute the Laplacian

$$\vec{\nabla}^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}.$$

$$\vec{\nabla}^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} (x^2 + y^2 + z^2 + 2) \right]$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} (2x + 0 + 0 + 0)$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} (2x)$$

$$\frac{\partial^2 f}{\partial x^2} = 2$$

$$\frac{\partial^2 f}{\partial z^2} = \frac{\partial}{\partial z} \left[\frac{\partial}{\partial z} (x^2 + y^2 + z^2 + 2) \right]$$

$$\frac{\partial^2 f}{\partial z^2} = \frac{\partial}{\partial z} (0 + 0 + 2z + 0)$$

$$\frac{\partial^2 f}{\partial z^2} = \frac{\partial}{\partial z} (2z)$$

$$\frac{\partial^2 f}{\partial z^2} = 2$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} \left[\frac{\partial}{\partial y} (x^2 + y^2 + z^2 + 2) \right]$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} (0 + 2y + 0 + 0)$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} (2y)$$

$$\frac{\partial^2 f}{\partial y^2} = 2$$

$$\vec{\nabla}^2 f = 2 + 2 + 2$$

$$\boxed{\vec{\nabla}^2 f = 6}$$

Problem 36

Given the scalar field $f(x, y, z) = xz + 3x^3y^2 + 2xz^2$, compute the Laplacian

$$\bar{\nabla}^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}.$$

$$\bar{\nabla}^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} (xz + 3x^3y^2 + 2xz^2) \right]$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} (z + 9x^2y^2 + 2z^2)$$

$$\frac{\partial^2 f}{\partial x^2} = 0 + 18xy^2 + 0$$

$$\frac{\partial^2 f}{\partial x^2} = 18xy^2$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} \left[\frac{\partial}{\partial y} (xz + 3x^3y^2 + 2xz^2) \right]$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} (0 + 6x^3y + 0)$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} (6x^3y)$$

$$\frac{\partial^2 f}{\partial y^2} = 6x^3$$

$$\frac{\partial^2 f}{\partial z^2} = \frac{\partial}{\partial z} \left[\frac{\partial}{\partial z} (xz + 3x^3y^2 + 2xz^2) \right]$$

$$\frac{\partial^2 f}{\partial z^2} = \frac{\partial}{\partial z} (x + 0 + 4xz)$$

$$\frac{\partial^2 f}{\partial z^2} = \frac{\partial}{\partial z} (x + 4xz)$$

$$\frac{\partial^2 f}{\partial z^2} = 0 + 4x$$

$$\frac{\partial^2 f}{\partial z^2} = 4x$$

$$\boxed{\bar{\nabla}^2 f = 18xy^2 + 6x^3 + 4x}$$

Computing the Laplacian of a Vector Field

Problem 37

Given the vector field $\vec{u} = (u_x, u_y, u_z) = (3x^2y, y^2z, 3z^2)$, find the Laplacian of \vec{u} , $\vec{\nabla}^2\vec{u}$.

$$\vec{\nabla}^2\vec{u} = (\vec{\nabla}^2u_x)\hat{i} + (\vec{\nabla}^2u_y)\hat{j} + (\vec{\nabla}^2u_z)\hat{k}$$

$$\vec{\nabla}^2u_x = \frac{\partial^2u_x}{\partial x^2} + \frac{\partial^2u_x}{\partial y^2} + \frac{\partial^2u_x}{\partial z^2}$$

$$\vec{\nabla}^2u_x = \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} (3x^2y) \right] + \frac{\partial}{\partial y} \left[\frac{\partial}{\partial y} (3x^2y) \right] + \frac{\partial}{\partial z} \left[\frac{\partial}{\partial z} (3x^2y) \right]$$

$$\vec{\nabla}^2u_x = \frac{\partial}{\partial x} (6xy) + \frac{\partial}{\partial y} (3x^2) + \frac{\partial}{\partial z} (0)$$

$$\vec{\nabla}^2u_x = 6y + 0 + 0$$

$$\vec{\nabla}^2u_x = 6y$$

$$\vec{\nabla}^2u_y = \frac{\partial^2u_y}{\partial x^2} + \frac{\partial^2u_y}{\partial y^2} + \frac{\partial^2u_y}{\partial z^2}$$

$$\vec{\nabla}^2u_y = \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} (y^2z) \right] + \frac{\partial}{\partial y} \left[\frac{\partial}{\partial y} (y^2z) \right] + \frac{\partial}{\partial z} \left[\frac{\partial}{\partial z} (y^2z) \right]$$

$$\vec{\nabla}^2u_y = \frac{\partial}{\partial x} (0) + \frac{\partial}{\partial y} (2yz) + \frac{\partial}{\partial z} (y^2)$$

$$\vec{\nabla}^2u_y = 0 + 2z + 0$$

$$\vec{\nabla}^2u_y = 2z$$

$$\vec{\nabla}^2u_z = \frac{\partial^2u_z}{\partial x^2} + \frac{\partial^2u_z}{\partial y^2} + \frac{\partial^2u_z}{\partial z^2}$$

$$\vec{\nabla}^2u_z = \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} (3z^2) \right] + \frac{\partial}{\partial y} \left[\frac{\partial}{\partial y} (3z^2) \right] + \frac{\partial}{\partial z} \left[\frac{\partial}{\partial z} (3z^2) \right]$$

$$\vec{\nabla}^2u_z = \frac{\partial}{\partial x} (0) + \frac{\partial}{\partial y} (0) + \frac{\partial}{\partial z} (6z)$$

$$\vec{\nabla}^2u_z = 0 + 0 + 6$$

$$\vec{\nabla}^2u_z = 6$$

$$\vec{\nabla}^2\vec{u} = (6y)\hat{i} + (2z)\hat{j} + (6)\hat{k}$$

$$\boxed{\vec{\nabla}^2\vec{u} = (6y, 2z, 6)}$$

Problem 38

Given the vector field $\vec{u} = (u_x, u_y, u_z) = (x^2 + y, x + y^2z, z^2)$, find the Laplacian of \vec{u} , $\vec{\nabla}^2\vec{u}$.

$$\vec{\nabla}^2\vec{u} = (\vec{\nabla}^2u_x)\hat{i} + (\vec{\nabla}^2u_y)\hat{j} + (\vec{\nabla}^2u_z)\hat{k}$$

$$\vec{\nabla}^2u_x = \frac{\partial^2u_x}{\partial x^2} + \frac{\partial^2u_x}{\partial y^2} + \frac{\partial^2u_x}{\partial z^2}$$

$$\vec{\nabla}^2u_x = \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} (x^2 + y) \right] + \frac{\partial}{\partial y} \left[\frac{\partial}{\partial y} (x^2 + y) \right] + \frac{\partial}{\partial z} \left[\frac{\partial}{\partial z} (x^2 + y) \right]$$

$$\vec{\nabla}^2u_x = \frac{\partial}{\partial x} (2x + 0) + \frac{\partial}{\partial y} (0 + 1) + \frac{\partial}{\partial z} (0 + 0)$$

$$\vec{\nabla}^2u_x = \frac{\partial}{\partial x} (2x) + \frac{\partial}{\partial y} (1) + \frac{\partial}{\partial z} (0)$$

$$\vec{\nabla}^2u_x = 2 + 0 + 0$$

$$\vec{\nabla}^2u_x = 2$$

$$\vec{\nabla}^2u_y = \frac{\partial^2u_y}{\partial x^2} + \frac{\partial^2u_y}{\partial y^2} + \frac{\partial^2u_y}{\partial z^2}$$

$$\vec{\nabla}^2u_y = \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} (x + y^2z) \right] + \frac{\partial}{\partial y} \left[\frac{\partial}{\partial y} (x + y^2z) \right] + \frac{\partial}{\partial z} \left[\frac{\partial}{\partial z} (x + y^2z) \right]$$

$$\vec{\nabla}^2u_y = \frac{\partial}{\partial x} (1 + 0) + \frac{\partial}{\partial y} (0 + 2yz) + \frac{\partial}{\partial z} (0 + y^2)$$

$$\vec{\nabla}^2u_y = \frac{\partial}{\partial x} (1) + \frac{\partial}{\partial y} (2yz) + \frac{\partial}{\partial z} (y^2)$$

$$\vec{\nabla}^2u_y = 0 + 2z + 0$$

$$\vec{\nabla}^2u_y = 2z$$

$$\vec{\nabla}^2 u_z = \frac{\partial^2 u_z}{\partial x^2} + \frac{\partial^2 u_z}{\partial y^2} + \frac{\partial^2 u_z}{\partial z^2}$$

$$\vec{\nabla}^2 u_z = \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} (z^2) \right] + \frac{\partial}{\partial y} \left[\frac{\partial}{\partial y} (z^2) \right] + \frac{\partial}{\partial z} \left[\frac{\partial}{\partial z} (z^2) \right]$$

$$\vec{\nabla}^2 u_z = \frac{\partial}{\partial x} (0) + \frac{\partial}{\partial y} (0) + \frac{\partial}{\partial z} (2z)$$

$$\vec{\nabla}^2 u_z = 0 + 0 + 2$$

$$\vec{\nabla}^2 u_z = 2$$

$$\vec{\nabla}^2 \vec{u} = (2)\hat{i} + (2z)\hat{j} + (2)\hat{k}$$

$$\boxed{\vec{\nabla}^2 \vec{u} = (2, 2z, 2)}$$

Problem 39

Given the vector field $\vec{u}(u_x, u_y, u_z) = (3x^2y, y^2z, 3z^2)$, find the Laplacian of \vec{u} , $\vec{\nabla}^2 \vec{u}$.

[duplicate to #37]

Derivative of a Vector with Respect to a Vector

Problem 40

Given $\vec{w} = \begin{bmatrix} w_1 \\ w_2 \end{bmatrix}$, $\begin{cases} w_1 = 2x + 3y \\ w_2 = 7x + 5y \end{cases}$, $\vec{u} = \begin{bmatrix} x \\ y \end{bmatrix}$, calculate $\frac{\partial \vec{w}}{\partial \vec{u}}$.

$$\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} \frac{\partial w_1}{\partial u_1} & \frac{\partial w_1}{\partial u_2} \\ \frac{\partial w_2}{\partial u_1} & \frac{\partial w_2}{\partial u_2} \end{bmatrix}$$

$$\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} \frac{\partial}{\partial x}(2x + 3y) & \frac{\partial}{\partial y}(2x + 3y) \\ \frac{\partial}{\partial x}(7x + 5y) & \frac{\partial}{\partial y}(7x + 5y) \end{bmatrix}$$

$$\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} 2 + 0 & 0 + 3 \\ 7 + 0 & 0 + 5 \end{bmatrix}$$

$$\boxed{\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} 2 & 3 \\ 7 & 5 \end{bmatrix}}$$

Problem 41

Given $\vec{w} = \begin{bmatrix} w_1 \\ w_2 \end{bmatrix}$, $\begin{cases} w_1 = x - y + z \\ w_2 = x + 2y - z \end{cases}$, $\vec{u} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, calculate $\frac{\partial \vec{w}}{\partial \vec{u}}$.

$$\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} \frac{\partial w_1}{\partial u_1} & \frac{\partial w_1}{\partial u_2} & \frac{\partial w_1}{\partial u_3} \\ \frac{\partial w_2}{\partial u_1} & \frac{\partial w_2}{\partial u_2} & \frac{\partial w_2}{\partial u_3} \end{bmatrix}$$

$$\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} \frac{\partial}{\partial x}(x - y + z) & \frac{\partial}{\partial y}(x - y + z) & \frac{\partial}{\partial z}(x - y + z) \\ \frac{\partial}{\partial x}(x + 2y - z) & \frac{\partial}{\partial y}(x + 2y - z) & \frac{\partial}{\partial z}(x + 2y - z) \end{bmatrix}$$

$$\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} 1 - 0 + 0 & 0 - 1 + 0 & 0 - 0 + 1 \\ 1 + 0 - 0 & 0 + 2 - 0 & 0 + 0 - 1 \end{bmatrix}$$

$$\boxed{\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} 1 & -1 & 1 \\ 1 & 2 & -1 \end{bmatrix}}$$

Problem 42

Given $\vec{w} = \begin{bmatrix} w_1 \\ w_2 \\ w_3 \end{bmatrix}$, $\begin{cases} w_1 = xy + z \\ w_2 = x - y^2 + z \\ w_3 = 2x + y + xz \end{cases}$, $\vec{u} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, calculate $\frac{\partial \vec{w}}{\partial \vec{u}}$.

$$\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} \frac{\partial w_1}{\partial u_1} & \frac{\partial w_1}{\partial u_2} & \frac{\partial w_1}{\partial u_3} \\ \frac{\partial w_2}{\partial u_1} & \frac{\partial w_2}{\partial u_2} & \frac{\partial w_2}{\partial u_3} \\ \frac{\partial w_3}{\partial u_1} & \frac{\partial w_3}{\partial u_2} & \frac{\partial w_3}{\partial u_3} \end{bmatrix}$$

$$\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} \frac{\partial}{\partial x}(xy + z) & \frac{\partial}{\partial y}(xy + z) & \frac{\partial}{\partial z}(xy + z) \\ \frac{\partial}{\partial x}(x - y^2 + z) & \frac{\partial}{\partial y}(x - y^2 + z) & \frac{\partial}{\partial z}(x - y^2 + z) \\ \frac{\partial}{\partial x}(2x + y + xz) & \frac{\partial}{\partial y}(2x + y + xz) & \frac{\partial}{\partial z}(2x + y + xz) \end{bmatrix}$$

$$\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} y + 0 & x + 0 & 0 + 1 \\ 1 - 0 + 0 & 0 - 2y + 0 & 0 - 0 + 1 \\ 2 + 0 + z & 0 + 1 + 0 & 0 + 0 + x \end{bmatrix}$$

$$\frac{\partial \vec{w}}{\partial \vec{u}} = \begin{bmatrix} y & x & 1 \\ 1 & -2y & 1 \\ 2 + z & 1 & x \end{bmatrix}$$

Derivative of a Scalar with Respect to a Vector (Jacobian)

Problem 43

Given $s = s(\vec{x}) = (x_1 + 1)^2 + x_2^2 + (x_3 + 2)^2$, $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$, calculate $\frac{\partial s}{\partial \vec{x}}$.

$$\frac{\partial s}{\partial \vec{x}} = \left[\frac{\partial s}{\partial x_1} \quad \frac{\partial s}{\partial x_2} \quad \frac{\partial s}{\partial x_3} \right]$$

$$\frac{\partial s}{\partial \vec{x}} = \left[\frac{\partial}{\partial x_1} [(x_1 + 1)^2 + x_2^2 + (x_3 + 2)^2] \quad \frac{\partial}{\partial x_2} [(x_1 + 1)^2 + x_2^2 + (x_3 + 2)^2] \quad \frac{\partial}{\partial x_3} [(x_1 + 1)^2 + x_2^2 + (x_3 + 2)^2] \right]$$

$$\frac{\partial s}{\partial \vec{x}} = [2(x_1 + 1) + 0 + 0 \quad 0 + 2x_2 + 0 \quad 0 + 0 + 2(x_3 + 2)]$$

$$\boxed{\frac{\partial s}{\partial \vec{x}} = [2x_1 + 2 \quad 2x_2 \quad 2x_3 + 4]}$$

Problem 44

Given $f = f(\vec{x}) = x + y + z$, $\vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, calculate $\frac{\partial f}{\partial \vec{x}}$.

$$\frac{\partial f}{\partial \vec{x}} = \left[\frac{\partial f}{\partial x} \quad \frac{\partial f}{\partial y} \quad \frac{\partial f}{\partial z} \right]$$

$$\frac{\partial f}{\partial \vec{x}} = \left[\frac{\partial}{\partial x} (x + y + z) \quad \frac{\partial}{\partial y} (x + y + z) \quad \frac{\partial}{\partial z} (x + y + z) \right]$$

$$\frac{\partial f}{\partial \vec{x}} = [1 + 0 + 0 \quad 0 + 1 + 0 \quad 0 + 0 + 1]$$

$$\boxed{\frac{\partial f}{\partial \vec{x}} = [1 \quad 1 \quad 1]}$$

Problem 45

Given $g = g(\vec{x}) = x + xy + z^2$, $\vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, calculate $\frac{\partial g}{\partial \vec{x}}$.

$$\frac{\partial g}{\partial \vec{x}} = \left[\frac{\partial g}{\partial x} \quad \frac{\partial g}{\partial y} \quad \frac{\partial g}{\partial z} \right]$$

$$\frac{\partial g}{\partial \vec{x}} = \left[\frac{\partial}{\partial x}(x + xy + z^2) \quad \frac{\partial}{\partial y}(x + xy + z^2) \quad \frac{\partial}{\partial z}(x + xy + z^2) \right]$$

$$\frac{\partial g}{\partial \vec{x}} = [1 + y + 0 \quad 0 + x + 0 \quad 0 + 0 + 2z]$$

$$\boxed{\frac{\partial g}{\partial \vec{x}} = [1 + y \quad x \quad 2z]}$$

Quadratic Forms

Problem 46

Given the quadric form $f(x, y) = x^2 + 6xy + 2y^2$:

- a. Express in matrix form, $f(\vec{x}) = \vec{x}^T \cdot A \cdot \vec{x}$.

$$f(x, y) = a_{11}x^2 + a_{12}xy + a_{21}xy + a_{22}y^2$$

$$f(x, y) = (1)x^2 + \left(\frac{6}{2}\right)xy + \left(\frac{6}{2}\right)xy + (2)y^2$$

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 3 \\ 3 & 2 \end{bmatrix}$$

$$f(x, y) = \vec{x}^T \cdot A \cdot \vec{x}$$

$$\boxed{f(x, y) = \vec{x}^T \cdot \begin{bmatrix} 1 & 3 \\ 3 & 2 \end{bmatrix} \cdot \vec{x}}$$

- b. Calculate $\frac{\partial f(\vec{x})}{\partial \vec{x}} = \frac{\partial}{\partial \vec{x}} (\vec{x}^T A \vec{x})$.

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = \frac{\partial}{\partial \vec{x}} (\vec{x}^T A \vec{x})$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = 2\vec{x}^T A^T$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = 2 \begin{bmatrix} x \\ y \end{bmatrix}^T \begin{bmatrix} 1 & 3 \\ 3 & 2 \end{bmatrix}^T$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = 2 \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 3 & 2 \end{bmatrix}$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = \begin{bmatrix} (2)(x)(1) + (2)(y)(3) \\ (2)(x)(3) + (2)(y)(2) \end{bmatrix}$$

$$\boxed{\frac{\partial f(\vec{x})}{\partial \vec{x}} = \begin{bmatrix} 2x + 6y \\ 6x + 4y \end{bmatrix}}$$

Problem 47

Given the quadric form $f(x, y) = 5x^2 + 2xy + 2y^2$:

- a. Express in matrix form, $f(\vec{x}) = \vec{x}^T \cdot A \cdot \vec{x}$.

$$f(x, y) = a_{11}x^2 + a_{12}xy + a_{21}xy + a_{22}y^2$$

$$f(x, y) = (5)x^2 + \left(\frac{2}{2}\right)xy + \left(\frac{2}{2}\right)xy + (2)y^2$$

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

$$A = \begin{bmatrix} 5 & 1 \\ 1 & 2 \end{bmatrix}$$

$$f(x, y) = \vec{x}^T \cdot A \cdot \vec{x}$$

$$\boxed{f(x, y) = \vec{x}^T \cdot \begin{bmatrix} 5 & 1 \\ 1 & 2 \end{bmatrix} \cdot \vec{x}}$$

- b. Calculate $\frac{\partial f(\vec{x})}{\partial \vec{x}} = \frac{\partial}{\partial \vec{x}} (\vec{x}^T A \vec{x})$.

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = \frac{\partial}{\partial \vec{x}} (\vec{x}^T A \vec{x})$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = 2\vec{x}^T A^T$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = 2 \begin{bmatrix} x \\ y \end{bmatrix}^T \begin{bmatrix} 5 & 1 \\ 1 & 2 \end{bmatrix}^T$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = 2 \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 5 & 1 \\ 1 & 2 \end{bmatrix}$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = \begin{bmatrix} (2)(x)(5) + (2)(y)(1) \\ (2)(x)(1) + (2)(y)(2) \end{bmatrix}$$

$$\boxed{\frac{\partial f(\vec{x})}{\partial \vec{x}} = \begin{bmatrix} 10x + 2y \\ 2x + 4y \end{bmatrix}}$$

Problem 48

Given the quadric form $f(x, y, z) = 3x^2 + 8xy + 6xz + y^2 + 6yz + 3z^2$:

- a. Express in matrix form, $f(\vec{x}) = \vec{x}^T \cdot A \cdot \vec{x}$.

$$f(x, y, z) = a_{11}x^2 + a_{12}xy + a_{21}xy + a_{13}xz + a_{31}xz + a_{22}y^2 + a_{23}yz + a_{32}yz + a_{33}z^2$$

$$f(x, y, z) = (3)x^2 + \left(\frac{8}{2}\right)xy + \left(\frac{8}{2}\right)xy + \left(\frac{6}{2}\right)xz + \left(\frac{6}{2}\right)xz + (1)y^2 + \left(\frac{6}{2}\right)yz + \left(\frac{6}{2}\right)yz + (3)z^2$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$A = \begin{bmatrix} 3 & 4 & 3 \\ 4 & 1 & 3 \\ 3 & 3 & 3 \end{bmatrix}$$

$$f(x, y, z) = \vec{x}^T \cdot A \cdot \vec{x}$$

$$f(x, y, z) = \vec{x}^T \cdot \begin{bmatrix} 3 & 4 & 3 \\ 4 & 1 & 3 \\ 3 & 3 & 3 \end{bmatrix} \cdot \vec{x}$$

- b. Calculate $\frac{\partial f(\vec{x})}{\partial \vec{x}} = \frac{\partial}{\partial \vec{x}}(\vec{x}^T A \vec{x})$.

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = \frac{\partial}{\partial \vec{x}}(\vec{x}^T A \vec{x})$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = 2\vec{x}^T A^T$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = 2 \begin{bmatrix} x \\ y \\ z \end{bmatrix}^T \begin{bmatrix} 3 & 4 & 3 \\ 4 & 1 & 3 \\ 3 & 3 & 3 \end{bmatrix}^T$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = 2 \begin{bmatrix} x & y & z \end{bmatrix} \begin{bmatrix} 3 & 4 & 3 \\ 4 & 1 & 3 \\ 3 & 3 & 3 \end{bmatrix}$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = \begin{bmatrix} (2)(x)(3) + (2)(y)(4) + (2)(z)(3) \\ (2)(x)(4) + (2)(y)(1) + (2)(z)(3) \\ (2)(x)(3) + (2)(y)(3) + (2)(z)(3) \end{bmatrix}$$

$$\frac{\partial f(\vec{x})}{\partial \vec{x}} = \begin{bmatrix} 6x + 8y + 6z \\ 8x + 2y + 6z \\ 6x + 6y + 6z \end{bmatrix}$$

END