

# STRAIGHTENING LAWS IN ALGEBRAIC COMPLEXITY

ANAKIN DEY

ABSTRACT.

## CONTENTS

|                                                   |    |
|---------------------------------------------------|----|
| 1. Introduction                                   | 3  |
| 1.1. Lower Bounds and Polynomial Identity Testing | 3  |
| 1.2. Geometric Complexity Theory                  | 3  |
| 1.3. Straightening Laws                           | 3  |
| 2. Preliminaries on Algebraic Complexity          | 5  |
| 2.1. Computational Models                         | 5  |
| 2.2. Complexity Classes                           | 7  |
| 2.3. Basic Structural Results                     | 7  |
| 2.4. Border Complexity                            | 9  |
| 3. Preliminaries on Straightening Laws            | 10 |
| 3.1. Determinantal Ideals                         | 10 |
| 3.2. Pfaffian Ideals                              | 12 |
| 3.3. Hankel Ideals                                | 15 |
| 3.4. Symmetric Determinantal Ideals               | 15 |
| 3.5. Permanental Ideals                           | 15 |
| 4. Prior Work                                     | 15 |
| 4.1. Generalities on Complexity in Ideals         | 15 |
| 5. Current Work on Ideals with Straightening Laws | 16 |
| 5.1. Approximative Results                        | 17 |

|                                            |    |
|--------------------------------------------|----|
| 5.2. Debordering                           | 17 |
| 5.3. Proof Overview for Determinantal Case | 18 |
| 6. Future Work and Open Problems           | 22 |
| 6.1. Symmetric Determinantal Ideals        | 22 |
| 6.2. Permanent Ideals                      | 22 |
| 6.3. Hankel and Toeplitz Ideals            | 22 |
| 6.4. Complexity of Straightening           | 22 |
| References                                 | 22 |

## 1. INTRODUCTION

⟨⟨ **Note about how alot of material here is based on [DG25].** ⟩⟩

Many problems in combinatorics, geometry, and computer science can be expressed in terms of the computation of some polynomials. For example, finding a perfect matching in a graph is equivalent to determining if it's *Tutte Matrix* has determinant identically zero [Tut47]. With this, we see that it is important to study the complexity of computing polynomials in general, and specifically computing special polynomials such as the determinant or permanent polynomials. Algebraic complexity theory is the study of the complexity of such computations.

⟨⟨ **Talk a bit about the history of the field starting with Valient** ⟩⟩

**1.1. Lower Bounds and Polynomial Identity Testing.** ⟨⟨ **Talk about Nisan-Wigderson, Kabanets-Impagliazzo** ⟩⟩

**1.2. Geometric Complexity Theory.** ⟨⟨ **Talk about Mulmuley-Sohoni** ⟩⟩

**1.3. Straightening Laws.** One of the most classical examples of standard monomial theory and straightening laws comes from the coordinate ring of the Grassmannian, whose relations are (*quadratic*)-*Plücker relations*. Many good references on this topic can be found [Ful96; LB18].

Let  $V$  be an  $n$ -dimensional vector space over a field  $\mathbb{F}$ . Let  $\{e_1, \dots, e_n\}$  be a basis for  $V$ . We will notate by  $\Lambda(n, d)$  the set of all ordered tuples  $(i_1 < \dots < i_d)$  where each  $i_k \in [n]$ .

**Definition 1.1 (Grassmannian):** Let  $\text{Gr}(n, d) = \{W \subseteq V \mid \dim_{\mathbb{F}}(W) = d\}$  be the  $d$ -dimensional *Grassmannian*.

One can give this the structure of an algebraic variety via the following map.

**Definition 1.2 (Plücker Embedding, Plücker Coordinates):** For  $U \in \text{Gr}(n, d)$ , let  $U$  have basis  $\{u_1, \dots, u_d\}$ . Define a map  $\text{Gr}(n, d) \rightarrow \mathbb{P}(\bigwedge^d V)$  given by  $U \mapsto [u_1 \wedge \dots \wedge u_d]$ . This map is called the *Plücker embedding*.

The set  $\{e_{\bar{i}} = e_{i_1} \wedge \dots \wedge e_{i_d} \mid \bar{i} \in \Lambda(n, d)\}$  gives a basis for  $\bigwedge^d V$ . Let  $\{p_{\bar{j}}\}_{\bar{j} \in \Lambda(n, d)}$  denote the basis of  $(\bigwedge^d V)^*$  dual to  $\{e_{\bar{i}}\}_{\bar{i} \in \Lambda(n, d)}$ . Then for each  $U \in \text{Gr}(n, d)$ , we can represent  $U$  via an  $n \times d$  matrix of rank  $d$  via the basis  $\{u_1, \dots, u_d\}$ . Then  $p_{\bar{j}}(U)$  is the  $d \times d$  determinant of the minor defined by rows  $\bar{j} \in \Lambda(n, d)$ . The coordinates  $p_{\bar{j}}$  are known as *Plücker coordinates*. We will often identify the Plücker coordinate  $p_{j_1, \dots, j_d}$  with the associated determinant.

**Definition 1.3 (Plücker Relations):** For each  $(i_1, \dots, i_{d-1}) \in \Lambda(n, d-1)$  and  $(j_1, \dots, j_{d+1}) \in \Lambda(n, d+1)$ , define the following quadratic polynomial in the Plücker coordinates:

$$\sum_{i=1}^{d+1} p_{i_1, \dots, i_{d-1}, j_i} \cdot p_{j_1, \dots, \widehat{j_i}, \dots, j_{d+1}}.$$

The ideal  $\mathfrak{q}$  of these quadratic polynomials is the ideal of *Plücker relations*.

Note that the ideal  $\mathfrak{q}$  of quadratic relations in the polynomial ring  $\mathbb{F}[p_{\bar{j}}]$  is a prime ideal [Ful96, §8.4, Proposition 2].

**Proposition 1.4 ([LB18, Theorem 12.1.3]):** The zero set of  $\mathfrak{q}$  is the Grassmannian  $\text{Gr}(n, d)$ .

**Definition 1.5 (Plücker / Bracket Algebra):** By Proposition 1.4, the coordinate ring  $\mathbb{F}[\text{Gr}(n, d)]$  of  $\text{Gr}(n, d)$  is given by

$$\mathbb{F}[p_{\bar{i}} \mid \bar{i} \in \Lambda(n, d)] / \mathfrak{q}.$$

In fact, one can also show that this coordinate ring is the invariant subring  $\mathbb{F}[x_{i,j} \mid 1 \leq i \leq n, 1 \leq j \leq d]^{\text{SL}_d(\mathbb{F})}$ . This algebra is known as the *Plücker* or *bracket algebra*.

The fact that the invariant subring  $\mathbb{F}[x_{i,j}]^{\text{SL}_d(\mathbb{F})}$  is generated by the  $d \times d$  minors is known as the *First Fundamental Theorem of Invariant Theory*. That the relations between the minors are all generated by the Plücker relations is known as the *Second Fundamental Theorem of Invariant Theory*.

We can represent the product of  $k$  minors using a  $d \times k$  tableau. One standard reference for this theory is [Stu08].

**Definition 1.6 (Tableau, Tableaux Ordering, Standard Monomials):** We can order the minors  $p_{i_1, \dots, i_d} \prec p_{j_1, \dots, j_d}$  if there is some  $1 \leq m \leq d$  such that  $i_k = j_k$  for  $1 \leq k \leq m-1$  and  $i_m < j_m$ . This gives a total order on the Plücker coordinates  $p_{\bar{j}}$  which induces a degree reverse lexicographic monomial ordering on the bracket algebra  $\mathbb{F}[p_{\bar{j}}]$ . This ordering is known as the *tableaux ordering* as we can express a product of  $k$  coordinates  $\bar{i}^1 \preceq \dots \preceq \bar{i}^k$  as a tableau

$$\begin{bmatrix} i_i^1 & \dots & i_d^1 \\ i_i^2 & \dots & i_d^2 \\ & \vdots & \\ i_i^k & \dots & i_d^k \end{bmatrix}.$$

We will say that such a monomial is *standard* if the columns are sorted (recall the rows are already sorted by convention). Otherwise, such a monomial is *nonstandard*.

This *standard monomial* theory is originally due to Hodge [Hod43]. However, to express the product of standard monomials in terms of other standard monomials, we use a *straightening* whose formulation is due to Young [You28].

**Definition 1.7 (van der Waerden Syzygies, Straightening Syzygies):** For  $\bar{\lambda} \in \Lambda(n, d)$ , let  $\bar{\lambda}^*$  be the *complement* tuple in  $\Lambda(n, n-d)$  consisting of the  $n-d$  elements which do not appear in  $\bar{\lambda}$ . The *sign* of  $\bar{\lambda}$  and  $\bar{\lambda}^*$ , denoted by  $\text{sgn}(\bar{\lambda}, \bar{\lambda}^*)$ , is the sign of the permutation  $\sigma \in \mathfrak{S}_n$  given in one-line notation by  $\lambda_1 \cdots \lambda_d \lambda_1^* \cdots \lambda_{n-d}^*$ .

For  $s \in [d]$ ,  $\bar{\alpha} \in \Lambda(n, s-1)$ ,  $\bar{\beta} \in \Lambda(n, d+1)$ , and  $\bar{\gamma} \in \Lambda(n, d-s)$ , the *van der Waerden syzygy*  $\left[\left[\bar{\alpha}\bar{\beta}\bar{\gamma}\right]\right]$  is the following quadratic polynomial in  $\mathbb{F}[p_{\bar{i}} \mid \bar{i} \in \Lambda(n, d)]$ :

$$\left[\left[\bar{\alpha}\bar{\beta}\bar{\gamma}\right]\right] := \sum_{\bar{\tau} \in \Lambda(d+1, s)} \text{sgn}(\bar{\tau}, \bar{\tau}^*) \cdot p_{\alpha_1, \dots, \alpha_s, \beta_{\tau_1^*}, \dots, \beta_{\tau_{d+1-s}^*}} \cdot p_{\beta_{\tau_1}, \dots, \beta_{\tau_s}, \gamma_1, \dots, \gamma_{d-s}}.$$

Such a van der Waerden syzygy is a *straightening syzygy* if  $\alpha_{s-1} < \beta_{s+1}$  and  $\beta_s < \gamma_s$ . Note that if we take  $s = 1$ , we get the Plücker relations.

**Theorem 1.8 ([Stu08, Theorem 3.1.7], [Rai]):** The set of straightening syzygies is a Gröbner basis for the ideal of quadratic Plücker relations  $\mathfrak{q}$  under the tableaux order. A monomial is standard if and only if it does not lie in the initial ideal  $\text{LM}_{\leq} \mathfrak{q}$ . Thus, the set of standard monomials forms an  $\mathbb{F}$ -vector space basis for  $\mathbb{F}[\text{Gr}(n, d)]$ . If we declare each  $p_{\bar{i}}$  to have degree one, then the standard tableau of shape  $d \times k$  give a basis for the  $k$ -th graded component of  $\mathbb{F}[\text{Gr}(n, d)]$ .

**Example 1.9:** ⟨ Example straightening calculations will go here ⟩

## 2. PRELIMINARIES ON ALGEBRAIC COMPLEXITY

### 2.1. Computational Models.

**Definition 2.1 (Algebraic Circuits):** An *algebraic circuit* over a field  $\mathbb{F}$  is a directed acyclic graph whose *input nodes*, nodes of in-degree zero, are labeled either by variables  $x_1, \dots, x_n$  or constants from  $\mathbb{F}$ . Nodes of in-degree greater than zero are labeled by  $\Sigma$  or  $\Pi$  which are *addition gates* or *multiplication gates* respectively. Each edge or *wire* is also labeled by a constant from  $\mathbb{F}$ .

Computation is defined inductively node by node. Input nodes compute their label. Every  $\Sigma$  gate computes the sum of their children multiplied by the constants along each wire, and computation at  $\Pi$  gates is defined similarly. Nodes of out-degree zero are *outputs*. We say that an algebraic circuit *computes*  $\{f_1(\bar{x}), \dots, f_k(\bar{x})\}$  if these are the polynomials computed by all the outputs.

The *size* of a circuit is the total number of edges. We will denote by  $\text{size}(f_1, \dots, f_k)$  the size of the smallest circuit over  $\mathbb{F}$  computing  $\{f_1(\bar{x}), \dots, f_k(\bar{x})\}$ . Some sources also define the size to be the number of vertices. These are polynomially related as a directed simple graph with  $v$  vertices has  $O(v^2)$  edges. It's *depth* is the length of the longest input-to-output path. The *fan-in* and *fan-out* of a node are in-degree and out-degree respectively. The *fan-in* and *fan-out* of the circuit is the maximum

fan-in and fan-out respectively over all nodes in the circuit. The *degree* or *syntactic degree* of a circuit is the maximum degree of a polynomial computed by any of the nodes of the circuit.

**Example 2.2:** Let  $f(x) = \sum_{i=0}^d a_i x^i \in \mathbb{F}[x]$  be a univariate polynomial.

Then *Horner's Rule* gives a recurrence for efficiently computing  $f(x)$ :

$$p_0(x) = a_0, \quad p_k(x) = a_{d-k} + x \cdot p_{k-1}(x).$$

This recurrence can be formulated as an algebraic circuit of size  $O(d)$  and depth  $O(d)$ . In fact, this uses exactly  $n$  additions and  $n$  multiplications, which is optimal. The proof of optimality of the number of additions for computing a generic univariate polynomial  $f(x)$  in this manner was given by [Ost54]. The corresponding proof of optimality for multiplications was given by [Pan66].

**Definition 2.3 (Algebraic Formulas):** An *algebraic formula* over a field  $\mathbb{F}$  is an algebraic circuit whose underlying graph is a tree.

**Definition 2.4 (Oracle Circuits and Formulas):** For a fixed polynomial  $f(\bar{x}) \in \mathbb{F}[\bar{x}]$ , an *f-oracle circuit* is an algebraic circuit defined in the same manner as Definition 2.1, however instead of just  $\Sigma$  and  $\Pi$  gates we now additionally allow  $f(\bar{x})$  gates. Note that such gates may have some inputs set as wires and some inputs set as fixed constants from  $\mathbb{F}$ . We also define an *f-oracle formula* as an *f-oracle circuit* whose underlying graph is a tree.

**Definition 2.5 (Algebraic Branching Programs):** An *algebraic branching program (ABP)* over a field  $\mathbb{F}$  is a directed acyclic graph  $A$  such that:

- The nodes  $V$  of  $A$  are partitioned into non-empty subsets  $V = V_0 \sqcup \dots \sqcup V_d$  for some integer  $d \geq 0$  and we call  $V_i$  the  $i$ -th layer. Layer  $V_0$  has exactly one node called the *source*  $s$  and layer  $V_d$  has exactly one node called the *sink*  $t$ .
- Each edge in  $A$  goes from layer  $V_{i-1}$  to  $V_i$  for some  $i \in [d]$ .
- Each edge  $e$  in  $A$  is labeled with a polynomial  $\gamma_e \in \mathbb{F}[\bar{x}]$  of degree at most one.

For a path  $\bar{e} = (e_1, \dots, e_d)$  from  $s$  to  $t$ , where each edge  $e_i$  goes from layer  $i-1$  to layer  $i$ , define the polynomial  $\gamma_{\bar{e}} := \prod_{i=1}^d \gamma_{e_i}$ . Then  $A$  computes the polynomial  $\sum_{\text{path } \bar{e} \text{ from } s \text{ to } t} \gamma_{\bar{e}}$ .

The *size* of an ABP  $A$  is the number of edges in  $A$ . Some sources also define the size to be the number of vertices. These are polynomially related as a directed simple graph with  $v$  vertices has  $O(v^2)$  edges. The *width* of  $A$  is the maximum number of vertices in any layer of  $A$ . The *depth* of  $A$  is the number of layers.

**Remark 2.6:** Definition 2.5 is really the definition of a *layered* ABP, where each edge only goes from one layer to the next without skipping any layers. However, one can convert a non-layered ABP to a layered ABP with only a quadratic blowup in size.

**Definition 2.7 (Determinantal Complexity):** For a polynomial  $f(\bar{x})$  over a field  $\mathbb{F}$ , the *determinantal complexity* of  $f(\bar{x})$ , denoted  $\text{dc}(f)$ , is the smallest  $m$  such that there exists an  $m \times m$  matrix  $M$  with entries in  $\mathbb{F}[\bar{x}]_{\leq 1}$  such that  $\det_m(M) = f(\bar{x})$ .

**Lemma 2.8 ([Val79; Bür24]):** Suppose that  $f(\bar{x})$  can be computed by an algebraic branching program of size  $m$ . Then  $\text{dc}(f) \leq m$ .

## 2.2. Complexity Classes.

**Definition 2.9 (VP):**

**Definition 2.10 (VF):**

**Definition 2.11 (VBP):**

**Definition 2.12 (VNP):**

## 2.3. Basic Structural Results.

**Lemma 2.13:** ⟨⟨ Depth reduction, conversions from circuits to formulas / ABPs and vice versa. ⟩⟩

**Proposition 2.14:** ⟨⟨ Inclusions ⟩⟩  $\text{VF} \subseteq \text{VBP} \subseteq \text{VP}$

**Lemma 2.15:** ⟨⟨ Elimination of Division ⟩⟩

**Definition 2.16:** For  $f \in \mathbb{F}[\bar{x}, t]$  and an integer  $i \in \mathbb{Z}_{\geq 0}$ , denote by  $\text{coeff}_{t^i}(f)$  the coefficient of  $t^i$  in  $f$  when  $f$  is viewed as a univariate polynomial in  $t$  over the ring  $\mathbb{F}[\bar{x}]$ .

**Lemma 2.17 ([DG25, Lemma 2.29]):** Let  $g \in \mathbb{F}[\bar{x}, t]$ , and let  $f \in \mathbb{F}[\bar{x}, t]$  be a degree- $d$  polynomial such that for each  $\alpha \in \mathbb{F}$ , the polynomial  $f(\bar{x}, \alpha)$  is computed by a  $g$ -oracle circuit  $C_\alpha$  of size at most  $s$  and depth at most  $\Delta$ . Assume that  $|\mathbb{F}| \geq d + 1$ . Then, for every  $0 \leq i \leq d$ , there exists a  $g$ -oracle circuit  $D_i$  of size  $O(ds)$  and depth  $\Delta + 1$  that computes  $\text{coeff}_{t^i}(f)$ , with its top gate being an addition gate and bottom  $\Delta$  layers consisting of  $d + 1$  copies of  $g$ -oracle circuits of the form  $C_\alpha$ . Moreover, if the top gate of each  $C_\alpha$  is an addition gate, then the depth bound can be improved to  $\Delta$ .

**Proof:** By definition,

$$f = \sum_{i=0}^d \text{coeff}_{t^i}(f) t^i.$$

Let  $\alpha_1, \dots, \alpha_{d+1}$  be  $d+1$  distinct elements in  $\mathbb{F}$ . Then we have that

$$\begin{pmatrix} \alpha_1^d & \alpha_1^{d-1} & \cdots & \alpha_1 & 1 \\ \alpha_2^d & \alpha_2^{d-1} & \cdots & \alpha_2 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \alpha_d^d & \alpha_d^{d-1} & \cdots & \alpha_d & 1 \\ \alpha_{d+1}^d & \alpha_{d+1}^{d-1} & \cdots & \alpha_{d+1} & 1 \end{pmatrix} \begin{pmatrix} \text{coeff}_{t^d}(f) \\ \text{coeff}_{t^{d-1}}(f) \\ \vdots \\ \text{coeff}_t(f) \\ \text{coeff}_1(f) \end{pmatrix} = \begin{pmatrix} f(\bar{x}, \alpha_1) \\ f(\bar{x}, \alpha_2) \\ \vdots \\ f(\bar{x}, \alpha_d) \\ f(\bar{x}, \alpha_{d+1}) \end{pmatrix}$$

The  $(d+1) \times (d+1)$  matrix  $(\alpha_i^{d+1-j}) \in \mathbb{F}^{(d+1) \times (d+1)}$  is a Vandermonde matrix. Its determinant is  $\prod_{i < j} (\alpha_j - \alpha_i)$ . As all the  $\alpha_i$  are distinct, this determinant is nonzero so that the matrix is invertible. Thus, there exists a  $(d+1) \times (d+1)$  matrix  $(z_{i,j}) \in \mathbb{F}^{(d+1) \times (d+1)}$  such that

$$\begin{pmatrix} \text{coeff}_{t^d}(f) \\ \text{coeff}_{t^{d-1}}(f) \\ \vdots \\ \text{coeff}_t(f) \\ \text{coeff}_1(f) \end{pmatrix} = \begin{pmatrix} z_{1,1} & z_{1,2} & \cdots & z_{1,d} & z_{1,d+1} \\ z_{2,1} & z_{2,2} & \cdots & z_{2,d} & z_{2,d+1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ z_{d,1} & z_{d,2} & \cdots & z_{d,d} & z_{d,d+1} \\ z_{d+1,1} & z_{d+1,2} & \cdots & z_{d+1,d} & z_{d+1,d+1} \end{pmatrix} \begin{pmatrix} f(\bar{x}, \alpha_1) \\ f(\bar{x}, \alpha_2) \\ \vdots \\ f(\bar{x}, \alpha_d) \\ f(\bar{x}, \alpha_{d+1}) \end{pmatrix}$$

Fix  $0 \leq i \leq d$ . We now have an explicit expression for  $\text{coeff}_{t^i}(f)$ :

$$(1) \quad \text{coeff}_{t^i}(f) = \sum_{j=1}^{d+1} z_{d+1-i,j} f(\bar{x}, \alpha_j)$$

With Equation (1), we can now describe the circuit for  $\text{coeff}_{t^i}(f)$ . The scalars  $\alpha_1, \dots, \alpha_{d+1}$  are fixed a priori, and hence so are the  $z_{i,j}$  for  $i, j \in [d+1]$ . Recall that scalar multiplications along wires are free. For each  $j \in [d+1]$ , computing  $f(\bar{x}, \alpha_j)$  requires a single  $g$ -oracle circuit  $C_{\alpha_j}$  of size at most  $s$  and depth  $\Delta$ . Hence, evaluating all  $d+1$  such terms requires total size  $O(ds)$ . The top-level summation uses one addition gate and  $d+1$  scalar multiplications by the  $z_{d+1-i,j}$  along  $d+1$  wires. Therefore, Equation (1) yields a  $g$ -oracle circuit  $D_i$  of size  $O(ds + d) = O(ds)$  and depth  $\Delta + 1$  computing  $\text{coeff}_{t^i}(f)$ . Moreover, if the top gate of each  $C_{\alpha_j}$  is an addition gate, then the top two layers of  $D_i$  consist entirely of additions and can be merged to reduce the depth bound to  $\Delta$ .  $\square$

**Remark 2.18:** The condition  $|\mathbb{F}| \geq d+1$  in Lemma 2.17 is required for interpolation. We note that if  $g$  is given not as a black box but as a white-box circuit, then its homogeneous components can be extracted in a gate-by-gate fashion [Str73], which requires no lower bound on  $|\mathbb{F}|$ .

**Lemma 2.19 ([KS01, Lemma 4]):** Let  $C$  be any collection of distinct linear forms in variables  $z_1, \dots, z_\ell$  with coefficients in the range  $\{0, \dots, K\}$  for some integer  $K \in \mathbb{Z}_{\geq 0}$ . Let  $\varepsilon > 0$ . Let  $z_1, \dots, z_\ell$  be independently and uniformly chosen from  $\{0, \dots, M\}$  at random, where  $M \geq K\ell/\varepsilon$ . Then, with probability at least  $1 - \varepsilon$ , there is a unique form in  $C$  of minimum value at  $z_1, \dots, z_\ell$ .

A simple rephrasing of Lemma 2.19 allows us to isolate a term of a multivariate polynomial  $f(y_1, \dots, y_\ell)$  under a random monomial substitution  $y_i \mapsto w^{z_i}$ :

**Corollary 2.20 ([DG25, Corollary 2.27]):** Let  $K \in \mathbb{Z}_{\geq 0}$ . Let  $A$  be any ring, and let  $f(y_1, \dots, y_\ell) \in A[\overline{y}]$  be a polynomial whose individual degree in each variable is at most  $K$ , i.e.,  $\deg_{y_i}(f) \leq K$  for all  $i \in [\ell]$ . Fix  $\varepsilon > 0$ . Choose  $z_1, \dots, z_\ell$  independently and uniformly at random from  $\{0, \dots, M\}$ , where  $M \geq K\ell/\varepsilon$ . Let  $w$  be a new indeterminate, and define  $\phi: A[y_1, \dots, y_\ell] \rightarrow A[w]$  to be the ring homomorphism sending  $y_i \mapsto w^{z_i}$  for each  $i \in [\ell]$ . Then, with probability at least  $1 - \varepsilon$ , there exists a unique monomial  $m = y_1^{e_1} \cdots y_\ell^{e_\ell}$  among all monomials of  $f$  such that  $\phi(m) = w^{e_1 z_1 + \dots + e_\ell z_\ell}$  attains the minimum degree in  $w$ .

## 2.4. Border Complexity.

**Definition 2.21 (Border Computation):** We say that  $g \in \mathbb{F}[x_1, \dots, x_n]$  is *border computed* by a circuit  $C$  over  $\mathbb{F}(\varepsilon)$  if  $C$  computes some polynomial  $h \in \mathbb{F}(\varepsilon)[x_1, \dots, x_n]$ , and  $h - g \in \varepsilon \cdot \mathbb{F}[[\varepsilon]][x_1, \dots, x_n]$ , where  $\mathbb{F}[[\varepsilon]]$  denotes the ring of formal power series in  $\varepsilon$ . We will denote by  $\overline{\text{size}}(f)$  the size of the smallest circuit over  $\mathbb{F}(\varepsilon)$  border computing  $f(\overline{x})$ .

In the settings where  $\mathbb{F}[\varepsilon][x_1, \dots, x_n]$  can be equipped with the Euclidean topology in a natural way, such as  $\mathbb{F} = \mathbb{C}$  or  $\mathbb{R}$ , this would imply that  $h \rightarrow g$  as  $\varepsilon \mapsto 0$ . However, this does not mean that we could just substitute  $\varepsilon$  by zero in  $C$  and obtain a circuit exactly computing  $g$  since  $C$  may use scalars in  $\mathbb{F}(\varepsilon)$ , such as  $1/\varepsilon$ , that are not well-defined at  $\varepsilon = 0$ .

**Example 2.22 ([DL25]):** In order to interpolate the coefficients of  $f(\overline{x}, t) = \sum_{i=0}^d \text{coeff}_{t^i}(f) \cdot t^i$  over  $\mathbb{F}$  as in Lemma 2.17, one needs that  $|\mathbb{F}| \geq d + 1$ . However, we can border compute  $\text{coeff}_{t^r}(f)$  if  $|\mathbb{F}| \geq r + 1$ .

Suppose  $\mathbb{F}$  has distinct elements  $\{\alpha_0, \dots, \alpha_r\}$ . Then, there exists elements  $\beta_0, \dots, \beta_r \in \mathbb{F}$  such that

$$\text{coeff}_{t^r}(f) = \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^r} \left( \sum_{i=0}^r \beta_i \cdot f(\overline{x}, \varepsilon \alpha_i) \right).$$

Indeed let  $g(\overline{x}, t) = \sum_{i=0}^r \text{coeff}_{t^i}(f) \cdot t^i$  and  $h(\overline{x}, t) = f(\overline{x}, t) - g(\overline{x}, t)$ . Then standard interpolation on  $\varepsilon \alpha_i$  gives us that there exists field constants  $\beta_0, \dots, \beta_r$  such that  $\varepsilon^r \text{coeff}_{t^r}(f) = \sum_{i=0}^r \beta_i \cdot g(\varepsilon \alpha_i)$ .

Thus, we have that

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^r} \left( \sum_{i=0}^r \beta_i \cdot f(\bar{x}, \varepsilon \alpha_i) \right) \\
&= \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^r} \left( \sum_{i=0}^r \beta_i \cdot g(\bar{x}, \varepsilon \alpha_i) \right) + \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^r} \left( \sum_{i=0}^r \beta_i \cdot h(\bar{x}, \varepsilon \alpha_i) \right) \\
&= \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^r} \left( \sum_{i=0}^r \beta_i \cdot g(\bar{x}, \varepsilon \alpha_i) \right) + 0 = \text{coeff}_{t^r}(f).
\end{aligned}$$

**Example 2.23:** ⟨⟨ Bordering Waring Rank ⟩⟩

⟨⟨ Define border complexity classes ⟩⟩

**Example 2.24:** ⟨⟨ Add some of Pranjali's examples of separations ⟩⟩

### 3. PRELIMINARIES ON STRAIGHTENING LAWS

**3.1. Determinantal Ideals.** Let  $\mathbb{F}$  be a field. Let  $X = (x_{i,j})_{\substack{1 \leq i \leq n \\ 1 \leq j \leq m}}$  be a  $n \times m$  matrix of variables. We will denote by  $\mathbb{F}[X]$  the polynomial ring  $\mathbb{F}[x_{i,j} \mid 1 \leq i \leq n, 1 \leq j \leq m]$ . We can natural associate a product of minors of  $X$  to a pair of Young tableau where the minors are described by the entries of the Young tableau.

**Definition 3.1 (Bitableau, bideterminant):** Let  $\sigma = (\sigma_1 \geq \dots \geq \sigma_k)$  be a partition. An  $(n, m)$ -bitableau  $(S \mid T)$  of shape  $\sigma$  is a pair of two Young tableau of shape  $\sigma$  with where the entries of  $S$  come from  $[n]$  and the entries of  $T$  come from  $[m]$ . When  $n$  and  $m$  are clear from context, we just call  $(S \mid T)$  a *bitableau*. For each  $i \in [\widehat{\sigma}_1]$ , write the  $i$ -th row of  $S$  and the  $i$ -th row of  $T$  as  $S_i = (S(i, 1), \dots, S(i, \sigma_i))$  and  $T_i = (T(i, 1), \dots, T(i, \sigma_i))$  respectively. The  $\sigma_i \times \sigma_i$  minor of  $X$  defined by  $S_i$  and  $T_i$  is the matrix

$$(x_{S(i,a), T(i,b)})_{1 \leq a, b \leq \sigma_i} = \begin{pmatrix} x_{S(i,1), T(i,1)} & x_{S(i,1), T(i,2)} & \cdots & x_{S(i,1), T(i, \sigma_i)} \\ x_{S(i,2), T(i,1)} & x_{S(i,2), T(i,2)} & \cdots & x_{S(i,2), T(i, \sigma_i)} \\ \vdots & \vdots & \ddots & \vdots \\ x_{S(i, \sigma_i), T(i,1)} & x_{S(i, \sigma_i), T(i,2)} & \cdots & x_{S(i, \sigma_i), T(i, \sigma_i)} \end{pmatrix}.$$

Then the associated *bideterminant*  $(S \mid T)(X)$  is the product of all such minors over  $i \in [\widehat{\sigma}_1]$ :

$$(S \mid T)(X) := \prod_{i=1}^{\widehat{\sigma}_1} \det_{\sigma_i} (x_{S(i,a), T(i,b)})_{1 \leq a, b \leq \sigma_i}.$$

An  $n \times n$  determinant is homogeneous of degree  $n$  so that  $(S \mid T)(X)$  is a homogeneous polynomial of degree  $|\sigma|$ .

**Example 3.2:** If  $\sigma = (4, 2, 1)$  and  $(S | T)$  is the bitableau

$$(S | T) = \left( \begin{array}{|c|c|c|c|} \hline 1 & 2 & 4 & 5 \\ \hline 3 & 6 & & \\ \hline 4 & & & \\ \hline \end{array} \middle| \begin{array}{|c|c|c|c|} \hline 1 & 3 & 5 & 6 \\ \hline 2 & 7 & & \\ \hline 3 & & & \\ \hline \end{array} \right),$$

then the bideterminant  $(S | T)(X)$  is given by

$$\begin{pmatrix} x_{1,1} & x_{1,3} & x_{1,5} & x_{1,6} \\ x_{2,1} & x_{2,3} & x_{2,5} & x_{2,6} \\ x_{4,1} & x_{4,3} & x_{4,5} & x_{4,6} \\ x_{5,1} & x_{5,3} & x_{5,5} & x_{5,6} \end{pmatrix} \begin{pmatrix} x_{3,2} & x_{3,7} \\ x_{6,2} & x_{6,7} \end{pmatrix} (x_{4,3}).$$

The following lemma shows that every monomial in  $\mathbb{F}[X]$  can be expressed as a bideterminant, and hence the bideterminants span  $\mathbb{F}[X]$  as an  $\mathbb{F}$ -vector space. The proof is immediate from the definition.

**Lemma 3.3:** A degree  $d$  monomial  $\prod_{i=1}^d x_{r_i, c_i}$  is given by a bideterminant:

$$\left( \begin{array}{|c|} \hline r_1 \\ \hline r_2 \\ \hline \vdots \\ \hline r_d \\ \hline \end{array} \middle| \begin{array}{|c|} \hline c_1 \\ \hline c_2 \\ \hline \vdots \\ \hline c_d \\ \hline \end{array} \right) (X) = \prod_{i=1}^d x_{r_i, c_i}.$$

Thus, the bideterminants span  $\mathbb{F}[X]$ .

While the bideterminants span  $\mathbb{F}[X]$ , there is a specific subset of bideterminants, the *standard* bideterminants, which form an  $\mathbb{F}$ -basis of  $\mathbb{F}[X]$ .

**Definition 3.4 (Standard bitableau, standard bideterminant):** We call a bitableau  $(S | T)$  and its associated bideterminant  $(S | T)(X)$  *standard* if  $S$  and  $T$  are both conjugate semistandard Young tableaux.

The next theorem is known in the literature as the *straightening law*.

**Theorem 3.5 ([DRS74, §8, Theorem 3], [dCEP80, Corollary 2.3]):** Let  $(S | T)(X)$  be a bideterminant of shape  $\sigma$ . Then  $(S | T)(X)$  can be uniquely expressed as a linear combination

$$(S | T)(X) = \sum_{(A|B)} c_{A,B} (A | B)(X),$$

where  $(A | B)(X)$  is a standard bideterminant of shape  $\tau \geq_{\text{lex}} \sigma$ , and  $c_{A,B} \in \mathbb{Z}$  when  $\text{char}(\mathbb{F}) = 0$ , while  $c_{A,B} \in \mathbb{Z}/p\mathbb{Z}$  when  $\text{char}(\mathbb{F}) = p > 0$ .

The fact that bideterminants are homogeneous polynomials, together with Lemma 3.3 and Theorems 3.5, imply the following corollary.

**Corollary 3.6:** The standard bideterminants form an  $\mathbb{F}[X]$ -basis of  $\mathbb{F}[X]$ . In particular, since bideterminants are homogeneous with respect to total degree, the degree- $d$  component of  $\mathbb{F}[X]$  has as basis the standard bideterminants of shape  $\sigma$  such that  $|\sigma| = d$ .

Let  $I_{n,m,r}^{\det}$  be the ideal of  $\mathbb{F}[X]$  generated by the  $r \times r$  minors of  $X$ . Note that this ideal is a  $\mathbb{F}$ -subspace of  $\mathbb{F}[X]$ . We have the following crucial statement on the  $\mathbb{F}$ -basis of this ideal.

**Proposition 3.7 ([BC03, Corollary 1.7]):** Polynomials  $f \in I_{n,m,r}^{\det}$  are supported by standard bideterminants of shape  $\sigma$  such that  $\sigma_1 \geq r$ ; that is, by those whose first row has length at least  $r$ .

The notion of the total degree of a polynomial is too coarse for differentiating terms in the expansion of a polynomial  $f \in I_{n,m,r}^{\det}$  with respect to the standard basis. Thus, we describe a grading of the polynomial ring  $\mathbb{F}[X]$  that is better suited for this purpose.

**Definition 3.8:** The polynomial ring  $\mathbb{F}[X]$  has a natural  $(\mathbb{N}^n \oplus \mathbb{N}^m)$ -grading. Let  $\bar{e}_i \in \mathbb{N}^n$  be the standard basis vector with a 1 in position  $i$  and 0's in the other positions. Define  $\bar{e}_i \in \mathbb{N}^m$  the same way. Then we assign  $x_{i,j}$  to have degree  $\bar{e}_i \oplus \bar{e}_j$ . We call this assignment the *multidegree*. We say a polynomial  $f \in \mathbb{F}[X]$  is *multihomogeneous* of multidegree  $(s_1\bar{e}_1 + \cdots + s_n\bar{e}_n) \oplus (t_1\bar{e}_1 + \cdots + t_m\bar{e}_m)$  if every monomial in  $f$  has multidegree  $(s_1\bar{e}_1 + \cdots + s_n\bar{e}_n) \oplus (t_1\bar{e}_1 + \cdots + t_m\bar{e}_m)$ .

The following lemma gives the multidegree of bideterminants. In particular, the multidegree of canonical and anti-canonical bideterminants will allow us to extract terms of a particular shape.

**Lemma 3.9:** Fix a natural number  $n \in \mathbb{Z}_{\geq 0}$ . Let  $\sigma = (\sigma_1 \geq \cdots \geq \sigma_d)$  be a partition such that the length  $\sigma_i$  of each row is at most  $n$ . Let  $S, T$  be two conjugate semistandard Young tableaux. Then a bideterminant  $(S | T)(X)$  is multihomogeneous of multidegree  $(s_1\bar{e}_1 + \cdots + s_n\bar{e}_n) \oplus (t_1\bar{e}_1 + \cdots + t_n\bar{e}_n)$ , where  $s_i$  is equal to the number of  $i$ 's that appear in  $S$  and  $t_j$  is equal to the number of  $j$ 's that appear in  $T$ . In particular,  $(K_\sigma | K_\sigma)(X)$  is multihomogeneous of multidegree  $(\hat{\sigma}_1\bar{e}_1 + \cdots + \hat{\sigma}_n\bar{e}_1) \oplus (\hat{\sigma}_1\bar{e}_1 + \cdots + \hat{\sigma}_n\bar{e}_1)$ . Moreover, for distinct partitions  $\sigma \neq \tau$ , the multidegrees of  $(K_\sigma | K_\sigma)(X)$  and  $(K_\tau | K_\tau)(X)$  are distinct, and the same holds for the multidegrees of  $(\bar{K}_\sigma | \bar{K}_\sigma)(X)$  and  $(\bar{K}_\tau | \bar{K}_\tau)(X)$ .

**Proof:** The multidegree of a standard bideterminant  $(S | T)(X)$  is immediate from Definition 3.1. To see that for distinct partitions  $\sigma \neq \tau$  the multidegrees of  $(K_\sigma | K_\sigma)(X)$  and  $(K_\tau | K_\tau)(X)$  are distinct, we make the following observation. The number of 1's in  $K_\sigma$  is the coefficient of  $\bar{e}_1$  in the multidegree, the number of 2's in  $K_\sigma$  is the coefficient of  $\bar{e}_2$ , and so on. This allows us to get the values  $\hat{\sigma}_1, \dots, \hat{\sigma}_n$  which in turn completely determines  $\sigma$ .  $\square$

**3.2. Pfaffian Ideals.** We now introduce notation for Pfaffian ideals. Let  $\mathbb{F}$  be a field. Let  $x_{i,j}$  be indeterminates for  $1 \leq i < j \leq 2n$  and let  $x_{j,i} := -x_{i,j}$  for  $1 \leq j < i \leq 2n$  and  $x_{i,i} = 0$  for  $1 \leq i \leq 2n$ .

Thus,  $X = (x_{i,j})_{1 \leq i,j \leq 2n}$  is a skew-symmetric  $2n \times 2n$  matrix of variables. For notational consistency with the prior subsections, we denote the polynomial ring  $\mathbb{F}[x_{i,j} \mid 1 \leq i < j \leq 2n]$  by  $\mathbb{F}[X]$ .

Recall that for a skew-symmetric matrix  $X$ , the determinant when viewed as a polynomial in the entries of the skew-symmetric matrix is a perfect square of some other polynomial. The square root of  $\det_{2n}(X)$  is called the *Pfaffian* of  $X$ . For the Pfaffian of a  $2n \times 2n$  matrix  $M \in \mathbb{F}^{2n \times 2n}$ , we use the notation  $\text{pfa}ff_{2n}$ . The Pfaffian has a more intrinsic definition given as follows:

$$(2) \quad \text{pfa}ff_{2n}(X) = \frac{1}{2^n n!} \sum_{\sigma \in \mathfrak{S}_{2n}} \text{sgn}(\sigma) \prod_{i=1}^n x_{\sigma(2i-1), \sigma(2i)}.$$

Just like the determinant, this is a polynomial with coefficients  $\pm 1$  so that it is well defined over any field. To see this, note that every monomial in Equation (2) appears exactly  $2^n n!$  times. We also have that if  $m$  is odd, then  $\det_m(X) = 0$  when  $X$  is a  $m \times m$  skew-symmetric matrix so that  $\text{pfa}ff_m(X) = 0$ . Thus, we only consider Pfaffians on matrices of even order.

Pfaffians are deeply connected to determinants, as indicated by the following lemma:

**Lemma 3.10 ([AF22, Lemma 2.27]):** Let  $X$  be a  $2n \times 2n$  skew-symmetric matrix and let  $E$  be an arbitrary  $2n \times 2n$  matrix. Then  $EXE^\top$  is also skew-symmetric and  $\text{pfa}ff_{2n}(EXE^\top) = \det_{2n}(E) \text{pfa}ff_{2n}(X)$ .

Since the Pfaffian is only well-defined for skew-symmetric matrices, we cannot consider all possible row and column combinations when looking at Pfaffians defined on minors of a matrix.

**Definition 3.11 (Principal bitableau, bipfaffian):** A bitableau  $(S \mid T)$  is *principal* if  $S = T$ , i.e., if the submatrix of  $X$  with rows given by  $S$  and columns given by  $T$  is principal. In this case, we denote  $[T] := (T \mid T)$ . Since  $m \times m$  determinants of skew-symmetric matrices are zero for odd  $m$ , we will assume all partitions have only even row lengths in this case. In particular,  $|\sigma|$  will always be divisible by 2.

Let  $\sigma$  be a partition such that the length of each row of  $\sigma$  is even, and let  $T$  be a Young tableau of shape  $\sigma$ . Writing the rows as  $T_i = (T(i, 1), \dots, T(i, \sigma_i))$ , the submatrix of  $X$  with rows and columns given by  $T_i$  is skew-symmetric as well and thus has a well-defined *principal pfaffian*. The associated *bipfaffian*<sup>\*</sup>  $[T](X)$  is the product of all such Pfaffians defined by the rows of  $T$ :

$$[T](X) := \prod_{i=1}^{\hat{\sigma}_1} \text{pfa}ff(x_{T(i,a), T(i,b)})_{1 \leq a, b \leq \sigma_i}.$$

Note that the total degree  $\deg([T](X))$  is equal to  $|\sigma|/2$ .

---

<sup>\*</sup>[AF22] call this a *standard monomial*. We deviate from this as standard monomial is a more general term which also refers to standard bideterminants and other analogues in standard monomial theory.

The following lemma shows that every monomial in  $\mathbb{F}[X]$  can be expressed as a bipfaffian and hence the bipfaffians span  $\mathbb{F}[X]$ . The proof is immediate from the definition.

**Lemma 3.12:** A degree  $d$  monomial  $\prod_{i=1}^d x_{r_i, c_i}$  where for each  $1 \leq i \leq d$ ,  $r_i < c_i$  is given by a bipfaffian:

$$\left[ \begin{array}{cc} r_1 & c_1 \\ r_2 & c_2 \\ \vdots & \vdots \\ r_d & c_d \end{array} \right] (X) = \prod_{i=1}^d x_{r_i, c_i}.$$

Analogous to the classical determinantal case, a specific subset of bipfaffians form an  $\mathbb{F}$ -basis of  $\mathbb{F}[X]$ .

**Definition 3.13 (Standard bipfaffian):** A bipfaffian  $[S](X)$  is *standard* if  $S$  is a conjugate semistandard Young tableau.

As in the determinantal case, the Pfaffian case also admits a straightening law.

**Theorem 3.14** ([dCP76, Theorem 6.5], [HT92, §5]): Let  $S$  be a Young tableau of shape  $\sigma$ . Then  $[S](X)$  can be uniquely expressed as a linear combination

$$[S](X) = \sum_A c_A [A](X),$$

where  $[A](X)$  is a standard bipfaffian of shape  $\tau \geq_{\text{lex}} \sigma$ , and  $c_A \in \mathbb{Z}$  when  $\text{char}(\mathbb{F}) = 0$ , while  $c_A \in \mathbb{Z}/p\mathbb{Z}$  when  $\text{char}(\mathbb{F}) = p > 0$ .

Lemma 3.12 and Theorems 3.14 together imply the following corollary.

**Corollary 3.15:** The standard bipfaffians form an  $\mathbb{F}$ -basis of  $\mathbb{F}[X]$ . In particular, since bipfaffians are homogeneous with respect to total degree, the degree- $d$  component of  $\mathbb{F}[X]$  has as basis the standard bipfaffians of shape  $\sigma$  such that  $|\sigma| = 2d$ . Note that if  $|\sigma| = 2d$ , then the number of rows  $\widehat{\sigma}_1$  of  $\sigma$  is at most  $d$  as each non-empty row of  $\sigma$  has length at least 2.

Let  $I_{2n, 2r}^{\text{pfaff}}$  be the ideal of  $\mathbb{F}[X]$  generated by the  $2r \times 2r$  principal bipfaffians of  $X$ . We have the following crucial statement.

**Proposition 3.16** ([HT92, §5]): Polynomials  $f \in I_{2n, 2r}^{\text{pfaff}}$  are supported by standard bipfaffians of shape  $\sigma$  such that  $\sigma_1 \geq 2r$ .

Next, we describe a grading of the polynomial ring  $\mathbb{F}[X]$  that is useful for the proof.

**Definition 3.17:** The polynomial ring  $\mathbb{F}[X]$  has a natural  $\mathbb{N}^{2n}$ -grading. Let  $\bar{e}_i \in \mathbb{N}^{2n}$  be the standard basis vector with a 1 in position  $i$  and 0's in the other positions. Then we assign  $x_{i,j}$  to have degree  $\bar{e}_i + \bar{e}_j$ . We call this assignment the *multidegree*. We say a polynomial  $f \in \mathbb{F}[X]$  is *multihomogeneous* of multidegree  $s_1 \bar{e}_1 + \cdots + s_{2n} \bar{e}_{2n}$  if every monomial in  $f$  has multidegree  $s_1 \bar{e}_1 + \cdots + s_{2n} \bar{e}_{2n}$ .

The following lemma gives the multidegree of bipfaffians. It follows immediately from the definition and is proven similarly to Lemma 3.9 and thus we omit the proof.

**Lemma 3.18:** Fix a natural number  $n \in \mathbb{Z}_{\geq 0}$ . Let  $\sigma = (\sigma_1 \geq \dots \geq \sigma_k)$  be a partition such that the length of each row is at most  $2n$ . Let  $S$  be a conjugate semistandard Young tableau. Then a bipfaffian  $[S](X)$  is multihomogeneous of multidegree  $s_1 \bar{e}_1 + \dots + s_{2n} \bar{e}_{2n}$ , where  $s_i$  is equal to the number of  $i$ 's that appear in  $S$ . In particular,  $[K_\sigma](X)$  is multihomogeneous of multidegree  $\hat{\sigma}_1 \bar{e}_1 + \dots + \hat{\sigma}_{2n} \bar{e}_{2n}$ . Moreover, for distinct partitions  $\sigma \neq \tau$ , the multidegrees of  $[K_\sigma](X)$  and  $[K_\tau](X)$  are distinct, and the same holds for the multidegrees of  $[\bar{K}_\sigma](X)$  and  $[\bar{K}_\tau](X)$ .

### 3.3. Hankel Ideals.

### 3.4. Symmetric Determinantal Ideals. ⟨⟨ Only if time ⟩⟩

### 3.5. Permanent Ideals. ⟨⟨ Only if time ⟩⟩

## 4. PRIOR WORK

**4.1. Generalities on Complexity in Ideals.** Roughly speaking, the complexity of an ideal  $I$  under a given algebraic model (such as algebraic circuits or algebraic branching programs) is defined as the minimum complexity, in that model, of any nonzero polynomial  $f \in I$  [Gro20]. Proving lower bounds for ideals is harder than for individual polynomials, since one must establish a bound for every nonzero  $f \in I$  rather than just a single polynomial. This motivates the search for a property of the following form: if an arbitrary nonzero  $f \in I$  is efficiently computable in a given model, then some polynomial from a distinguished set  $S = \{g_1, \dots, g_k\}$  (often a natural generating set of  $I$  or some related set with desirable properties) is also efficiently computable in the given model. In this way, proving lower bounds for arbitrary nonzero  $f \in I$  reduces to proving them for  $g_1, \dots, g_k$ . We call such a property a *closure result*, by analogy with closure under factorization in the principal ideal case. In short, closure results reduce the task of showing hardness for all polynomials in  $I$  to proving hardness for a small set of distinguished polynomials.

**Definition 4.1 (Hitting Set Generator (HSG)):**

**Definition 4.2 (Polynomial Identity Testing (PIT)):**

Closure results for ideals are useful for constructing hitting set generators, which are the algebraic-complexity analogs of pseudorandom generators and can be used to derandomize polynomial identity testing (PIT). The idea is as follows: for  $s < n$ , we seek a hitting set generator  $G: \mathbb{F}^s \rightarrow \mathbb{F}^n$  against a family  $\mathcal{C}$  of  $n$ -variate polynomials. This means that  $G$  has the property that for any nonzero polynomial  $f \in \mathcal{C}$ , we have  $f \circ G \neq 0$ . Here, one should think of  $\mathcal{C}$  as a family of “low-complexity” polynomials. Let  $I$  be the ideal of polynomials  $f$  satisfying  $f \circ G = 0$ , also known as the *vanishing ideal* of  $G$  [HMM24].

To show that  $G$  has the desired hitting set property, it suffices to show that every nonzero  $f \in I$  has sufficiently high complexity so that no such  $f$  can belong to  $\mathcal{C}$ . With a closure result for  $I$ , this reduces to proving lower bounds for a distinguished set of polynomials  $\{g_1, \dots, g_k\}$  for  $I$ . This can be viewed as a particular way of implementing the “hardness versus randomness” paradigm of [NW94; KI04] in the setting of algebraic complexity.

The general question we want to study is of the following form:

**Question 4.3:** Say that  $I = \langle g_1, \dots, g_k \rangle$ . Let  $0 \neq f \in I$ . How does the complexity of  $f$  compare to the complexity of the generators  $g_1, \dots, g_k$ .

4.1.1. *Principal Ideals.* Let us consider the special case of Question 4.3 where  $I = \langle g \rangle$  is a *principal ideal*. Then our question is now the following:

**Question 4.4:** Let  $0 \neq f \in \langle g \rangle$ . Does  $g$  have a small  $f$ -oracle circuit?

The affirmative was actually conjectured by Bürgisser:

**Conjecture 4.5 ([Bür00, Conjecture 8.3]):** If  $g$  is a factor of  $f \neq 0$ , then  $\text{size}(g) \leq \text{poly}(\text{size}(f), \deg(g))$ .

There has been much progress on Conjecture 4.5. Factoring in algebraic complexity is well studied, see the survey of Bhargav, Dwivedi, and Saxena for an overview of recent progress [BDS26]. For example, it is known that for fields of characteristic zero that  $\text{size}(g) \leq \text{poly}(\text{size}(f), \deg(f))$  [Kal87; Bür00]. Compare the use of  $\deg(f)$  here to  $\deg(g)$  in Conjecture 4.5. Thus, the most interesting regime to study here is the situation where the degree of the factor  $g$  is significantly smaller than the degree of the whole polynomial  $f$ . This result is only known in the setting of border computation over fields of characteristic zero:

**Theorem 4.6 ([Bür04, Theorem 1.3]):** Let  $\mathbb{F}$  be a field of characteristic zero. Let  $f(\bar{x}) \neq 0$  be an  $n$ -variate polynomial. If  $g(\bar{x})$  is a factor of  $f(\bar{x})$ , then  $\overline{\text{size}}(g) \leq \text{poly}(\text{size}(f), \deg(g), n)$ .

Bürgisser’s result can actually be formulated in the setting of oracle complexity, so that we get a  $\text{poly}(\deg(g), n)$  size  $f$ -oracle circuit which approximately computes  $g(\bar{x})$ .

**<< TODO: Read and understand Kayal’s work on annihilator polynomials and how it connects to HSGs >>**

4.1.2. *PIT vs Factoring.*

4.1.3. *More Than One Generator.*

## 5. CURRENT WORK ON IDEALS WITH STRAIGHTENING LAWS

**<< TODO: motivate Grochow’s conjecture >>**

**Conjecture 5.1** ([Gro20, Conjecture 6.3]): Let  $X$  be an  $n \times n$  matrix of indeterminates and let  $I_n$  be the ideal generated by the  $\frac{n}{2} \times \frac{n}{2}$  minors of  $X$ . Then for every nonzero polynomial  $f \in I_n$ , there is a constant-depth<sup>†</sup> algebraic circuit of size  $\text{poly}(n)$  with  $f$ -oracle gates that computes the  $m \times m$  determinant for some  $m = n^{\Theta(1)}$ .

**5.1. Approximative Results.** Analogous to Bürgisser’s theorem [Bür04] that the Factor Conjecture holds with respect to border complexity, Andrews and Forbes [AF22] proved that Grochow’s conjecture holds with respect to border complexity. Formally, they proved the following theorem:

**Theorem 5.2** ([AF22, Theorem 1.1]): Let  $\mathbb{F}$  be a field of characteristic zero. Let  $X = (x_{i,j})_{\substack{1 \leq i \leq n \\ 1 \leq j \leq m}}$  be an  $n \times m$  matrix of variables over  $\mathbb{F}$  and let  $I_{n,m,r}^{\det} \subseteq \mathbb{F}[X] = \mathbb{F}[x_{i,j}]$  be the ideal generated by the  $r \times r$  minors of  $X$ . Let  $f \in I_{n,m,r}^{\det}$  be a nonzero polynomial. Then there is a depth-three  $f$ -oracle circuit of size  $O(n^2 m^2)$  that approximately computes the  $t \times t$  determinant for  $t = \Theta(r^{1/3})$ .

**Theorem 5.3** ([AF22, Theorem 1.1]): Let  $\mathbb{F}$  be a field of characteristic zero. Let  $X = (x_{i,j})_{1 \leq i,j \leq 2n}$  be a  $2n \times 2n$  matrix of variables over  $\mathbb{F}$  and let  $I_{2n,2r}^{\text{pfaff}} \subseteq \mathbb{F}[X] = \mathbb{F}[x_{i,j}]$  be the ideal generated by the Pfaffians of the  $2r \times 2r$  principal submatrices of  $X$ . Let  $f \in I_{2n,2r}^{\text{pfaff}}$  be a nonzero polynomial. Then there is a depth-three  $f$ -oracle circuit of size  $O(n^4)$  that approximately computes the  $t \times t$  Pfaffian for  $t = \Theta(r^{1/3})$ .

**5.2. Debordering.** In [DG25], we deborder the results of [AF22]. This is done via a novel application of the Isolation Lemma: Lemma 2.19 and Corollary 2.20.

**Theorem 5.4 (Informal version of [DG25, Theorem 3.18, Corollary 3.19]):** Let  $X = (x_{i,j})_{\substack{1 \leq i \leq n \\ 1 \leq j \leq m}}$  be an  $n \times m$  matrix of variables over a field  $\mathbb{F}$ . Let  $I_{n,m,r}^{\det} \subseteq \mathbb{F}[X] = \mathbb{F}[x_{i,j} \mid 1 \leq i \leq n, 1 \leq j \leq m]$  be the ideal generated by the  $r \times r$  minors of  $X$ . Let  $f \in I_{n,m,r}^{\det}$  be a nonzero polynomial. Assume that the characteristic of  $\mathbb{F}$  is either zero or greater than  $\deg(f)$ . Also assume that the size of  $\mathbb{F}$  is sufficiently large.<sup>‡</sup> Then there exists a depth-three  $f$ -oracle circuit of size  $\text{poly}(n, m, \deg(f))$  that computes the  $t \times t$  determinant for  $t = \Theta(r^{1/3})$ .

**Theorem 5.5 (Informal version of [DG25, Theorem 4.10, Corollary 4.11]):** Let  $X = (x_{i,j})_{1 \leq i < j \leq 2n}$  be a  $2n \times 2n$  skew-symmetric matrix of variables over a field  $\mathbb{F}$  so that  $x_{j,i} = -x_{i,j}$  for  $1 \leq j < i \leq n$  and  $x_{i,i} = 0$  for  $1 \leq i \leq 2n$ . Let  $I_{2n,2r}^{\text{pfaff}} \subseteq \mathbb{F}[X] = \mathbb{F}[x_{i,j} \mid 1 \leq i < j \leq 2n]$  be the ideal generated by the  $2r \times 2r$  principal Pfaffians of  $X$ . Let  $f \in I_{2n,2r}^{\text{pfaff}}$  be a nonzero polynomial. Assume that the characteristic of  $\mathbb{F}$  is either zero or greater than  $\deg(f)$ . Also assume that the size of  $\mathbb{F}$  is sufficiently large. Then

<sup>†</sup>The original statement of the conjecture does not specify constant-depth algebraic circuits. However, it is already known that the  $m \times m$  determinant can be computed by algebraic circuits of size  $\text{poly}(m)$ .

<sup>‡</sup>Specifically, we require  $|\mathbb{F}|$  to be bounded below by some polynomial in  $n, m$ , and  $\deg(f)$ . Such assumptions are standard in the literature (for instance, in work on polynomial factorization), since one can often achieve them by passing to a suitable field extension.

there exists a depth-three  $f$ -oracle circuit of size  $\text{poly}(n, \deg(f))$  that computes the  $t \times t$  Pfaffian of a skew-symmetric matrix for  $t = \Theta(r^{1/3})$ .

### 5.3. Proof Overview for Determinantal Case.

**5.3.1. Linear Operators.** Our application of the determinantal straightening law (Corollary 3.6), together with the isolation lemma (Corollary 2.20) and extraction of coefficients (Lemma 2.17), allows us to isolate a canonical bitableau from the expansion of nonzero  $f \in I_{n,m,r}^{\det}$  in the basis of standard bideterminants. In particular, we will show that this can be done via a polynomial-sized depth-three  $f$ -oracle circuit using the isolation lemma (Corollary 2.20) as well as coefficient extraction (Lemma 2.17). This is in contrast to the proof of [AF22], where they use a Kronecker-type substitution in order to isolate terms. To do this, we will give a more careful analysis of the terms that arise when applying the substitution operators to the expression of a polynomial  $f(X) \in I_{n,m,r}^{\det}$ .

Throughout, fix a field  $\mathbb{F}$ . Let  $n, m$  be natural numbers, and for  $1 \leq i \leq n$  and  $1 \leq j \leq m$ , let  $x_{i,j}$  be indeterminates. Throughout this section, let  $X = (x_{i,j})$  be an  $n \times m$  matrix of variables. Let  $1 \leq r \leq \min(n, m)$  and define  $I_{n,m,r}^{\det}$  to be the ideal in  $\mathbb{F}[X]$  generated by the  $r \times r$  minors of  $X$ .

Our first step will be to establish a series of linear transformations that send a conjugate semistandard bitableau to the anti-canonical conjugate semistandard bitableau of the same shape.

**Definition 5.6 (Substitution operator):** Let  $i < j$ . Define the operator  $\text{Sub}_{i \rightarrow j}$  which takes as input a tableau  $S$  and returns the tableau  $S'$  which is formed by taking each row of  $S$  which has an  $i$  but not a  $j$ , replacing that  $i$  with  $j$ , and then sorting the row in increasing order. We also define  $h_i^j(S)$  to be the number of times  $i$  is replaced by  $j$  after applying  $\text{Sub}_{i \rightarrow j}$  to  $S$ .

**Definition 5.7 (Canonical/anti-canonical tableau):** Fix a natural number  $n$ . Let  $\sigma = (\sigma_1 \geq \dots \geq \sigma_k > 0)$  be a partition such that the length  $\sigma_i$  of each row is at most  $n$ . Define  $K_{\sigma,n}$  to be the tableau of shape  $\sigma$  where row  $i$  has entries  $1, 2, \dots, \sigma_i$ . Similarly, define  $\bar{K}_{\sigma,n}$  to be the tableau of shape  $\sigma$  where row  $i$  has entries  $(n - \sigma_i + 1, n - \sigma_i + 2, \dots, n)$ . We call  $K_{\sigma,n}$  and  $\bar{K}_{\sigma,n}$  the *canonical* and *anti-canonical* tableau respectively.<sup>§</sup> Note that  $K_{\sigma,n}$  and  $\bar{K}_{\sigma,n}$  are both conjugate semistandard. When  $n$  is clear from context, we will just write  $K_{\sigma}$  and  $\bar{K}_{\sigma}$ .

**Example 5.8:** If  $\sigma = (5, 4, 2, 1)$  and  $n = 7$ , then  $K_{\sigma}$  and  $\bar{K}_{\sigma}$  are given in Figure 1.

<sup>§</sup>In some sources, such as [BV88], these are referred to as the *initial* and *final* tableau.

|   |   |   |   |   |  |  |  |  |  |   |   |   |   |   |  |  |  |  |  |
|---|---|---|---|---|--|--|--|--|--|---|---|---|---|---|--|--|--|--|--|
| 1 | 2 | 3 | 4 | 5 |  |  |  |  |  | 3 | 4 | 5 | 6 | 7 |  |  |  |  |  |
| 1 | 2 | 3 | 4 |   |  |  |  |  |  | 4 | 5 | 6 | 7 |   |  |  |  |  |  |
| 1 | 2 |   |   |   |  |  |  |  |  | 6 | 7 |   |   |   |  |  |  |  |  |
| 1 |   |   |   |   |  |  |  |  |  | 7 |   |   |   |   |  |  |  |  |  |

FIGURE 1. The canonical and anti-canonical tableau for  $\sigma = (5, 4, 2, 1)$  and  $n = 7$ .

We now study how these substitution operators act on bideterminants  $(S | T)(X)$ . From properties of the determinant, we see that multiplying the matrix of variables  $X$  by a elementary matrix  $E_{i,j}$  corresponds to applying substitution operators on  $S$  or  $T$ :

**Lemma 5.9:** Let  $\lambda$  be a new indeterminate. Let  $(S | T)$  be a bitableau, not necessarily standard, of shape  $\sigma$ . For  $0 \leq h \leq h_i^j(S) - 1$ , let  $\mathcal{C}_{i \rightarrow j}^h(S)$  be the set of tableaux of shape  $\sigma$  obtained by changing  $i$  to  $j$  at exactly  $h$  rows of  $S$  which contain  $i$  but not  $j$  and reordering those rows to be increasing. Then we have that

$$(3) \quad (S | T)(E_{i,j}(\lambda)X) = \pm \lambda^{h_i^j(S)} (\text{Sub}_{i \rightarrow j}(S) | T)(X) + \sum_{h=0}^{h_i^j(S)-1} \lambda^h \sum_{S' \in \mathcal{C}_{i \rightarrow j}^h(S)} \pm (S' | T)(X).$$

Similarly, we also have that

$$(4) \quad (S | T)(X E_{i,j}(\lambda)^\top) = \pm \lambda^{h_i^j(T)} (S | \text{Sub}_{i \rightarrow j}(T))(X) + \sum_{h=0}^{h_i^j(T)-1} \lambda^h \sum_{T' \in \mathcal{C}_{i \rightarrow j}^h(T)} \pm (S | T')(X).$$

**Definition 5.10:** In addition to the variables  $x_{i,j}$ , where  $i \in [n]$  and  $j \in [m]$ , we introduce new sets of indeterminates  $\Lambda = \{\lambda_{i,j} \mid 1 \leq i < j \leq n\}$  and  $\Xi = \{\xi_{i,j} \mid 1 \leq i < j \leq m\}$ . We have that  $|\Lambda| = \binom{n}{2} = O(n^2)$  and  $|\Xi| = \binom{m}{2} = O(m^2)$ . The purpose of these new indeterminates is that they will allow us to keep track of the effects of substitution operators as we isolate a canonical bitableau.

Define  $M \in \mathbb{F}[\Lambda]^{n \times n}$  and  $N \in \mathbb{F}[\Xi]^{m \times m}$  by

$$M = E_{1,2}(\lambda_{1,2}) \cdots E_{1,n}(\lambda_{1,n}) E_{2,3}(\lambda_{2,3}) \cdots E_{2,n}(\lambda_{2,n}) \\ \cdots E_{n-2,n-1}(\lambda_{n-2,n-1}) E_{n-2,n}(\lambda_{n-2,n}) E_{n-1,n}(\lambda_{n-1,n})$$

and

$$N = E_{m-1,m}(\xi_{m-1,m})^\top E_{m-2,m}(\xi_{m-2,m})^\top E_{m-2,m-1}(\xi_{m-2,m-1})^\top \\ \cdots E_{2,m}(\xi_{2,m})^\top \cdots E_{2,3}(\xi_{2,3})^\top E_{1,m}(\xi_{1,m})^\top \cdots E_{1,2}(\xi_{1,2})^\top.$$

For an integer  $k > 0$ , let  $J_k \in \mathbb{F}^{k \times k}$  be the  $k \times k$  matrix with 1's along the anti-diagonal from the bottom left to the top right and 0's elsewhere. Note that  $\det_k(J_k) = (-1)^{\binom{k}{2}} = \pm 1$ . The substitution  $X \mapsto J_n X J_m$  has the effect of replacing row  $i$  with row  $n - i + 1$  and column  $j$  with column  $m - j + 1$ . This turns an anti-canonical bitableau  $(\bar{K}_\sigma \mid \bar{K}_\sigma)$  into a canonical one  $(K_\sigma \mid K_\sigma)$ .

### 5.3.2. Isolation of a Single Term.

**Corollary 5.11:** Let  $f(X) \in I_{n,m,r}^{\det}$  be a nonzero polynomial of degree  $d$ . Then  $f(MJ_n X J_m N)$  can be expressed as a sum

$$(5) \quad f(MJ_n X J_m N) = \sum_{k \in A} \hat{c}_k \Lambda^{\bar{e}_k} \Xi^{\bar{f}_k} \cdot (K_{\sigma_k} \mid K_{\sigma_k})(X) + \sum_{\ell \in B} \hat{c}_\ell \Lambda^{\bar{e}_\ell} \Xi^{\bar{f}_\ell} \cdot (S_\ell \mid T_\ell)(X),$$

such that the following hold:

- (1)  $A$  is a nonempty finite set and  $B$  is finite set disjoint from  $A$ .
- (2) All terms in Equation (5) have the form  $c \Lambda^{\bar{e}} \Xi^{\bar{f}} \cdot (S \mid T)(X)$ , where  $c \in \mathbb{F}^\times$ ,  $\bar{e} \in \{0, 1, \dots, d\}^{\binom{n}{2}}$ ,  $\bar{f} \in \{0, 1, \dots, d\}^{\binom{m}{2}}$ , and  $(S \mid T)(X)$  is a bideterminant in  $I_{n,m,r}^{\det}$  of degree  $d$ . In particular, if  $S$  and  $T$  have shape  $\sigma$ , then  $|\sigma| \leq d$ .
- (3) The triples  $(\bar{e}_k, \bar{f}_k, \sigma_k)$  are distinct, where  $k$  ranges over  $A$ .

Next, we introduce  $n$  variables  $Y = \{y_1, \dots, y_n\}$  and  $Z = \{z_1, \dots, z_m\}$ . Define the diagonal matrices

$$D = \text{diag}(y_1, \dots, y_n) \in \mathbb{F}[Y]^{n \times n} \quad \text{and} \quad D' = \text{diag}(z_1, \dots, z_m) \in \mathbb{F}[Z]^{m \times m}.$$

**Lemma 5.12:** Let  $(S \mid T)(X)$  be a bideterminant of degree at most  $d$ . Then

$$(S \mid T)(DXD') = Y^{\bar{s}} Z^{\bar{t}} \cdot (S \mid T)(X),$$

where  $\bar{s} = (s_1, \dots, s_n) \in \{0, 1, \dots, d\}^n$ ,  $\bar{t} = (t_1, \dots, t_m) \in \{0, 1, \dots, d\}^m$ ,  $s_i$  is the number of times  $i$  appears in  $S$ ,  $t_j$  is the number of times  $j$  appears in  $T$ , i.e.,  $(s_1 e_1 + \dots + s_n e_n) \oplus (t_1 e_1 + \dots + t_m e_m)$  is the multidegree of  $(S \mid T)(X)$  with respect to the grading defined in Definition 3.8.

We will often simply say that  $(S \mid T)(X)$  in Lemma 5.12 has multidegree  $(\bar{s}, \bar{t})$  rather than writing it as  $(s_1 e_1 + \dots + s_n e_n) \oplus (t_1 e_1 + \dots + t_m e_m)$ .

Next, we introduce another variable  $v$ , and modify  $D$  and  $D'$  as follows:

$$\tilde{D} = \text{diag}(y_1 v, y_2 v^2, \dots, y_n v^n) \in \mathbb{F}[Y, v]^{n \times n} \quad \text{and} \quad \tilde{D}' = \text{diag}(z_1 v, z_2 v^2, \dots, z_m v^m) \in \mathbb{F}[Z, v]^{m \times m}.$$

**Lemma 5.13:** Let  $(S \mid T)(X)$  be a bideterminant. Then

$$(S \mid T)(\tilde{D}X\tilde{D}') = Y^{\bar{s}} Z^{\bar{t}} v^{|\bar{s}|+|\bar{t}|} \cdot (S \mid T)(X),$$

where  $(\bar{s}, \bar{t})$  is the multidegree of  $(S \mid T)(X)$ .

**Lemma 5.14:** Let  $f(X) \in I_{n,m,r}^{\det}$  be a nonzero polynomial of degree  $d$ . Let  $g = f(MJ_n \tilde{D}X \tilde{D}'J_m N) \in \mathbb{F}[X, \Lambda, \Xi, Y, Z, v]$ . View  $g$  as a univariate polynomial in  $v$  with coefficients in  $\mathbb{F}[X, \Lambda, \Xi, Y, Z]$ , and write  $g = \sum_i \text{coeff}_{v^i}(g) v^i$ , where  $\text{coeff}_{v^i}(g) \in \mathbb{F}[X, \Lambda, \Xi, Y, Z]$  denotes the coefficient of  $v^i$  in  $g$ . Choose the smallest integer  $d_{\min}$  such that  $\text{coeff}_{v^{d_{\min}}}(g) \neq 0$ . Then  $d_{\min} = 2|K_\sigma|$  for some shape  $\sigma$  with  $|\sigma| \leq d$  and we may write  $\text{coeff}_{v^{d_{\min}}}(g)$  as a finite sum

$$(6) \quad \text{coeff}_{v^{d_{\min}}}(g) = \sum_{k \in I} c_k \Lambda^{\bar{e}_k} \Xi^{\bar{f}_k} Y^{\bar{s}_k} Z^{\bar{t}_k} \cdot (K_{\sigma_k} \mid K_{\sigma_k})(X),$$

such that the following hold:

- (1) For each  $k \in I$ ,  $c_k \in \mathbb{F}^\times$ ,  $(\sigma_k)_1 \geq r$ , and  $|\sigma_k| \leq d$ .
- (2) The tuples  $(\bar{e}_k, \bar{f}_k, \bar{s}_k, \bar{t}_k)$  are distinct, where  $k$  ranges over  $I$ .

Lemma 5.14 helps us separate a collection of terms solely containing canonical bideterminants. The next lemma further applies the isolation lemma to single out one term from this collection. **⟨ I will better motivate isolation lemma here. ⟩**

**Lemma 5.15:** Let  $f(X) \in I_{n,m,r}^{\det}$  be a nonzero polynomial of degree  $d$ . Let  $g = f(MJ_n \tilde{D}X \tilde{D}'J_m N) \in \mathbb{F}[X, \Lambda, \Xi, Y, Z, v]$ . Then there exist integers  $z_t = O(d(n^2 + m^2))$  for each variable  $t \in \Lambda \sqcup \Xi \sqcup Y \sqcup Z$ , and an integer  $z_v = O(d^2(n^2 + m^2))$  such that for the two variable substitution maps

$$\phi : t \mapsto w^{z_t} \quad \text{for } t \in \Lambda \sqcup \Xi \sqcup Y \sqcup Z$$

and

$$\psi : v \mapsto w^{z_v},$$

we have that  $h := (\psi \circ \phi)(g) \in \mathbb{F}[X, w]$  has  $\deg_w(h) = O(d^3(n^3 + m^3))$ , and that  $\text{coeff}_{w^i}(h) = c \cdot (K_\sigma \mid K_\sigma)(X)$  for some integer  $i \leq \deg_w(h)$ , where  $c \in \mathbb{F}^\times$ ,  $\sigma_1 \geq r$  and  $|\sigma| \leq d$ .

Lemma 5.14 helps us separate a collection of terms solely containing canonical bideterminants. The next lemma further applies the isolation lemma to single out one term from this collection.

**Theorem 5.16:** Let  $f(X) \in I_{n,m,r}^{\det}$  be a nonzero polynomial of degree  $d$ . Assume  $|\mathbb{F}| \geq c_0 d^3(n^3 + m^3)$ , where  $c_0 > 0$  is a large enough constant. Then, there exists a depth-three  $f$ -oracle circuit of size  $O(n^2 m^2 d^3(n^3 + m^3)) = \text{poly}(n, m, d)$  computing  $(K_\sigma \mid K_\sigma)(X)$ , where  $\sigma_1 \geq r$  and  $|\sigma| \leq d$ . Furthermore, the top gate of this circuit is an addition gate, and the bottom layer consists of  $O(nmd^3(n^3 + m^3))$  addition gates. The total number of gates and wires, excluding those between the bottom addition gates and the input gates, is  $O(nmd^3(n^3 + m^3))$ .

### 5.3.3. Projection to Determinant.

## 6. FUTURE WORK AND OPEN PROBLEMS

## 6.1. Symmetric Determinantal Ideals.

6.2. Permanent Ideals.  $\langle\langle$  if time  $\rangle\rangle$ 

## 6.3. Hankel and Toeplitz Ideals.

6.4. Complexity of Straightening.  $\langle\langle$  if time  $\rangle\rangle$ 

## REFERENCES

- [AF22] Robert Andrews and Michael A. Forbes. “Ideals, Determinants, and Straightening: Proving and Using Lower Bounds for Polynomial Ideals”. In: *Proceedings of the 54th Annual ACM SIGACT Symposium on Theory of Computing*. June 9, 2022, pp. 389–402. DOI: [10.1145/3519935.3520025](https://doi.org/10.1145/3519935.3520025). arXiv: [2112.00792 \[cs\]](https://arxiv.org/abs/2112.00792). URL: <http://arxiv.org/abs/2112.00792> (visited on 01/19/2026).
- [BC03] Winfried Bruns and Aldo Conca. “Gröbner Bases and Determinantal Ideals”. In: *Commutative Algebra, Singularities and Computer Algebra*. Springer Netherlands, 2003, pp. 9–66. DOI: [10.1007/978-94-007-1092-4\\_2](https://doi.org/10.1007/978-94-007-1092-4_2).
- [BDS26] C. S. Bhargav, Prateek Dwivedi, and Nitin Saxena. *A Primer on the Closure of Algebraic Complexity Classes under Factoring*. Version 2. June 9, 2026. DOI: [10.48550/arXiv.2506.19604](https://doi.org/10.48550/arXiv.2506.19604). arXiv: [2506.19604 \[cs.CC\]](https://arxiv.org/abs/2506.19604). Pre-published.
- [Bür00] Peter Bürgisser. *Completeness and Reduction in Algebraic Complexity Theory*. Springer Berlin Heidelberg, 2000. ISBN: 978-3-662-04179-6. DOI: [10.1007/978-3-662-04179-6](https://doi.org/10.1007/978-3-662-04179-6).
- [Bür04] Peter Bürgisser. “The Complexity of Factors of Multivariate Polynomials”. In: *Foundations of Computational Mathematics* 4.4 (Nov. 2004), pp. 369–396. ISSN: 1615-3375, 1615-3383. DOI: [10.1007/s10208-002-0059-5](https://doi.org/10.1007/s10208-002-0059-5). URL: <http://link.springer.com/10.1007/s10208-002-0059-5> (visited on 01/19/2026).
- [Bür24] Peter Bürgisser. *Completeness Classes in Algebraic Complexity Theory*. June 10, 2024. DOI: [10.48550/arXiv.2406.06217](https://doi.org/10.48550/arXiv.2406.06217). arXiv: [2406.06217 \[cs\]](https://arxiv.org/abs/2406.06217). URL: <http://arxiv.org/abs/2406.06217> (visited on 10/23/2025). Pre-published.
- [BV88] Winfried Bruns and Udo Vetter. *Determinantal Rings*. Springer, 1988. DOI: [10.1007/BFb0080378](https://doi.org/10.1007/BFb0080378).
- [dCEP80] Corrado de Concini, David Eisenbud, and Claudio Procesi. “Young Diagrams and Determinantal Varieties”. In: *Inventiones mathematicae* 56.2 (1980), pp. 129–165. DOI: [10.1007/BF01392548](https://doi.org/10.1007/BF01392548).

- [dCP76] Corrado de Concini and Claudio Procesi. “A Characteristic Free Approach to Invariant Theory”. In: *Advances in Mathematics* 21.3 (1976), pp. 330–354. ISSN: 0001-8708. DOI: [10.1016/S0001-8708\(76\)80003-5](https://doi.org/10.1016/S0001-8708(76)80003-5).
- [DG25] Anakin Dey and Zeyu Guo. *Debordering Closure Results in Determinantal and Pfaffian Ideals*. Nov. 21, 2025. Electronic Colloquium on Computational Complexity: [TR25-189](https://eccc.weizmann.ac.il/report/2025/189/). URL: <https://eccc.weizmann.ac.il/report/2025/189/> (visited on 03/24/2026). Pre-published.
- [DL25] Pranjal Dutta and Vladimir Lysikov. *Recent Advances in Debordering Methods*. Oct. 15, 2025. DOI: [10.48550/arXiv.2510.13049](https://doi.org/10.48550/arXiv.2510.13049). arXiv: [2510.13049 \[cs\]](https://arxiv.org/abs/2510.13049). URL: <http://arxiv.org/abs/2510.13049> (visited on 10/20/2025). Pre-published.
- [DRS74] Peter Doubilet, Gian-Carlo Rota, and Joel Stein. “On the Foundations of Combinatorial Theory: IX Combinatorial Methods in Invariant Theory”. In: *Studies in Applied Mathematics* 53.3 (1974), pp. 185–216. DOI: [10.1002/sapm1974533185](https://doi.org/10.1002/sapm1974533185).
- [Ful96] William Fulton. *Young Tableaux: With Applications to Representation Theory and Geometry*. 2nd ed. London Mathematical Society Student Texts 35. Cambridge: Cambridge University Press, 1996.
- [Gro20] Joshua A. Grochow. “Complexity in Ideals of Polynomials: Questions on Algebraic Complexity of Circuits and Proofs”. In: *Bull. EATCS* 131 (2020). URL: <http://eatcs.org/beatcs/index.php/beatcs/article/view/620>.
- [HMM24] Ivan Hu, Dieter van Melkebeek, and Andrew Morgan. “Polynomial Identity Testing via Evaluation of Rational Functions”. In: *Theory of Computing* 20.1 (July 18, 2024), pp. 1–70. DOI: [10.4086/toc.2024.v020a001](https://doi.org/10.4086/toc.2024.v020a001). URL: <https://theoryofcomputing.org/articles/v020a001/> (visited on 07/10/2026).
- [Hod43] W. V. D. Hodge. “Some Enumerative Results in the Theory of Forms”. In: *Mathematical Proceedings of the Cambridge Philosophical Society* 39.1 (Mar. 1943), pp. 22–30. DOI: [10.1017/S0305004100017631](https://doi.org/10.1017/S0305004100017631).
- [HT92] Jürgen Herzog and Ngô Việt Trung. “Gröbner Bases and Multiplicity of Determinantal and Pfaffian Ideals”. In: *Advances in Mathematics* 96.1 (1992), pp. 1–37. DOI: [10.1016/0001-8708\(92\)90050-U](https://doi.org/10.1016/0001-8708(92)90050-U).
- [Kal87] E. Kaltofen. “Single-Factor Hensel Lifting and Its Application to the Straight-Line Complexity of Certain Polynomials”. In: *Proceedings of the Nineteenth Annual ACM Symposium on Theory of Computing*. STOC ’87. New York, NY, USA: Association for Computing Machinery, Jan. 1, 1987, pp. 443–452. ISBN: 978-0-89791-221-1. DOI: [10.1145/28395.28443](https://doi.org/10.1145/28395.28443). URL: <https://dl.acm.org/doi/10.1145/28395.28443> (visited on 07/12/2026).
- [KI04] Valentine Kabanets and Russell Impagliazzo. “Derandomizing Polynomial Identity Tests Means Proving Circuit Lower Bounds”. In: *computational complexity* 13.1–2 (Dec. 2004),

- pp. 1–46. ISSN: 1016-3328, 1420-8954. DOI: [10.1007/s00037-004-0182-6](https://doi.org/10.1007/s00037-004-0182-6). URL: <http://link.springer.com/10.1007/s00037-004-0182-6> (visited on 10/23/2025).
- [KS01] Adam R. Klivans and Daniel Spielman. “Randomness Efficient Identity Testing of Multivariate Polynomials”. In: *Proceedings of the Thirty-Third Annual ACM Symposium on Theory of Computing*. STOC01: 33rd ACM Symposium on Theory of Computing. Herionissos Greece: ACM, July 6, 2001, pp. 216–223. ISBN: 978-1-58113-349-3. DOI: [10.1145/380752.380801](https://doi.org/10.1145/380752.380801). URL: <https://dl.acm.org/doi/10.1145/380752.380801> (visited on 10/14/2025).
- [LB18] V. Lakshmibai and Justin Brown. *Flag Varieties*. Vol. 53. Texts and Readings in Mathematics. Singapore: Springer, 2018. ISBN: 978-981-13-1393-6. DOI: [10.1007/978-981-13-1393-6](https://doi.org/10.1007/978-981-13-1393-6). URL: <http://link.springer.com/10.1007/978-981-13-1393-6> (visited on 07/16/2026).
- [NW94] Noam Nisan and Avi Wigderson. “Hardness vs Randomness”. In: *Journal of Computer and System Sciences* 49.2 (Oct. 1, 1994), pp. 149–167. ISSN: 0022-0000. DOI: [10.1016/S0022-0000\(05\)80043-1](https://doi.org/10.1016/S0022-0000(05)80043-1). URL: <https://www.sciencedirect.com/science/article/pii/S0022000005800431> (visited on 06/17/2026).
- [Ost54] A. Ostrowski. “On Two Problems in Abstract Algebra Connected with Horner’s Rule”. In: *Studies in Mathematics and Mechanics*. Academic Press, 1954, pp. 40–48. ISBN: 978-1-4832-3272-0. URL: <http://dx.doi.org/10.1016/B978-1-4832-3272-0.50010-7>.
- [Pan66] V Ya Pan. “Methods of Computing Values of Polynomials”. In: *Russian Mathematical Surveys* 21.1 (Feb. 1966), pp. 105–136.
- [Rai] Claudiu Raicu. *Algebras with Straightening Law*. URL: [https://www.slmath.org/ckeditor\\_assets/attachments/783/raicu.pdf](https://www.slmath.org/ckeditor_assets/attachments/783/raicu.pdf).
- [Str73] Volker Strassen. “Vermeidung von divisionen.” In: *Journal für die reine und angewandte Mathematik* 264 (1973), pp. 184–202. URL: <http://eudml.org/doc/151394>.
- [Stu08] Bernd Sturmfels. *Algorithms in Invariant Theory*. 2nd ed. Texts & Monographs in Symbolic Computation. Springer Vienna, 2008, p. vii 197. 197 pp. ISBN: 978-3-211-77417-5. DOI: [10.1007/978-3-211-77417-5](https://doi.org/10.1007/978-3-211-77417-5). URL: <http://link.springer.com/book/10.1007%2F978-3-211-77417-5>.
- [Tut47] William T Tutte. “The factorization of linear graphs”. In: *Journal of the London Mathematical Society* 1.2 (1947), pp. 107–111.
- [Val79] L. G. Valiant. “Completeness Classes in Algebra”. In: *Proceedings of the Eleventh Annual ACM Symposium on Theory of Computing*. STOC ’79. New York, NY, USA: Association for Computing Machinery, Apr. 30, 1979, pp. 249–261. ISBN: 978-1-4503-7438-5. DOI: [10.1145/800135.804419](https://doi.org/10.1145/800135.804419). URL: <https://dl.acm.org/doi/10.1145/800135.804419> (visited on 07/11/2026).

- [You28] Alfred Young. “On Quantitative Substitutional Analysis”. In: *Proceedings of the London Mathematical Society* s2-28.1 (Jan. 1928), pp. 255–292. ISSN: 0024-6115. DOI: [10.1112/plms/s2-28.1.255](https://doi.org/10.1112/plms/s2-28.1.255).