

1 Symmetric Functions

First we review the construction of the ring of symmetric functions as an inverse limit in the category of *graded rings*. For a construction in terms of a limit of graded rings, see [Mac95]. We will assume basic notations for partitions and (skew) tableaux following [Ful96].

Definition: For $n \geq 0$, let $\Lambda_n = \mathbb{Z}[x_1, \dots, x_n]^{\mathfrak{S}_n}$ be the ring of symmetric polynomials over $\mathbb{Z}$ in n variables. For $k \geq 0$, let Λ_n^k be the k -th graded component of Λ_n , where each x_i has degree one. The *ring of symmetric functions* Λ is the direct sum $\bigoplus_{k \geq 0} \Lambda^k$. To obtain the ring of symmetric functions Λ_A over rings A other than $\mathbb{Z}$, repeating the above construction gives $\Lambda_A \simeq \Lambda \otimes_{\mathbb{Z}} A$.

1.1 Common Bases for the Ring of Symmetric Functions

Definition:

- Let $\lambda = (\lambda_1 \geq \dots \geq \lambda_n \geq 0)$ be any partition with length $\ell(\lambda) \leq n$. Then the *monomial symmetric function* $m_\lambda(x_1, \dots, x_n)$ is defined by

$$m_\lambda(x_1, \dots, x_n) = \sum_{\alpha} x_1^{\alpha_1} \cdots x_n^{\alpha_n}$$

where the summation is taken over all permutations α of λ . We have that Λ^k is the free $\mathbb{Z}$ -module of rank $p(k)$ with basis $m_\lambda(x_1, \dots, x_n)$ for all $n \geq k$ as $\lambda \vdash k$ ranges over all partitions of k , where $p(k)$ is the number of partitions of k .

With this, we can also define Λ^k as the span of all monomial symmetric functions m_λ such that $|\lambda| = k$.

- For $r \geq 0$, the r -th *elementary symmetric function* e_r is defined by

$$e_r = \sum_{i_1 < \dots < i_r} x_{i_1} \cdots x_{i_r} = m_{(1^r)}.$$

For a partition $\lambda = (\lambda_1, \lambda_2, \dots)$ we define $e_\lambda = e_{\lambda_1} e_{\lambda_2} \cdots$. The *Fundamental Theorem of Symmetric Functions* states that over finitely many variables, say $x_1, \dots, x_n$, the elementary symmetric functions $e_1, \dots, e_n$ are algebraically independant and $\Lambda_n \simeq \mathbb{Z}[e_1, \dots, e_n]$. For a clean proof of this fact due to van der Waerden, see [Stu08, Theorem 1.1.1].

- For $r \geq 0$, the r -th *complete homogeneous symmetric function* h_r is defined by

$$h_r = \sum_{i_1 \leq \dots \leq i_r} x_{i_1} \cdots x_{i_r} = \sum_{\lambda \vdash r} m_\lambda.$$

For a partition $\lambda = (\lambda_1, \lambda_2, \dots)$ we define $h_\lambda = h_{\lambda_1} h_{\lambda_2} \cdots$.

1.2 Schur Functions We follow [Sta23, Chapter 7] for the exposition on Schur functions.

Definition: For a Young tableaux T , its *type* is the vector $\alpha = (\alpha_1, \alpha_2, \dots)$ where α_i is the number of entries of T labeled with i . If T has type α , write $\bar{x}^T = x_1^{\alpha_1} x_2^{\alpha_2} \cdots$. For a partition λ , the *Schur function* $s_\lambda(\bar{x})$ is defined by

$$s_\lambda(\bar{x}) = \sum_{\text{SSYT } T \text{ of shape } \lambda} \bar{x}^T$$

where the summation is taken over all semistandard Young tableau of shape λ .

Note that the Schur functions also form a basis of Λ .

Proposition: Let

$$a_\delta = \det(x_i^{n-j})_{1 \leq i, j \leq n} = \prod_{1 \leq i < j \leq n} (x_i - x_j)$$

be the *Vandermonde determinant*. For a partition $\lambda = (\lambda_1 \geq \dots \geq \lambda_n \geq 0)$, the *alternant* $a_{\lambda+\delta}$ is the determinant

$$a_{\lambda+\delta} = \det(x_i^{\lambda_j+n-j})_{1 \leq i, j \leq n}.$$

Then $s_\lambda(x_1, \dots, x_n) = \frac{a_{\lambda+\delta}}{a_\delta}$.

Proposition: For a partition λ , we have that

$$s_\lambda(x_1, \dots, x_n) = \det(h_{\lambda_i+j-i}(\bar{x}))_{1 \leq i, j \leq n},$$

$$s_{\lambda'}(x_1, \dots, x_n) = \det(e_{\lambda_i+j-i}(\bar{x}))_{1 \leq i, j \leq n},$$

the *Jacobi-Trudi* and *Nägelsbach-Kostka* identities respectively.

In particular we have that $e_r = s_{(1^r)}$ and $h_r = s_{(r)}$.

Theorem (Pieri Rules): For a partition λ and an integer $r \geq 0$, we have that

$$h_r(\bar{x}) \cdot s_\lambda(\bar{x}) = \sum_{\mu} s_\mu$$

where the summation over μ is over all partitions μ formed by added r boxes to the diagram of λ such that no two boxes are added to the same column.

We also have that

$$e_r(\bar{x}) \cdot s_\lambda(\bar{x}) = \sum_{\mu} s_\mu$$

where the summation over μ is over all partitions μ formed by added r boxes to the diagram of λ such that no two boxes are added to the same row.

1.3 An Involution and Inner Product

Definition: We have a map $\omega: \Lambda \rightarrow \Lambda$ given by $\omega(e_\lambda) = h_\lambda$.

Proposition: One can check that $\omega(h_\lambda) = e_\lambda$ so that ω is an involution. Furthermore, we have that $\omega(p_\lambda) = (-1)^{|\lambda|-\ell(\lambda)} p_\lambda$ and $\omega(s_\lambda) = s_{\lambda'}$.

Definition: We define an inner product $\langle \cdot, \cdot \rangle$ on Λ by making the two bases $\{m_\lambda\}_\lambda$ and $\{h_\mu\}_\mu$ dual: $\langle m_\lambda, h_\mu \rangle = \delta_{\lambda, \mu}$ for all partitions λ, μ .

Proposition: Let $z_\lambda = \prod_k k^{n_k(\lambda)} \cdot n_k(\lambda)!$ where $n_k(\lambda)$ is the number of parts of λ of size k . Then $\langle p_\lambda, p_\mu \rangle = z_\lambda \delta_{\lambda, \mu}$. We also have that $\langle s_\lambda, s_\mu \rangle = \delta_{\lambda, \mu}$ so that the Schur functions form an orthonormal basis of Λ .

Proposition: The involution ω is self-adjoint: $\langle f, \omega(g) \rangle = \langle \omega(f), g \rangle$. Thus, we have that $\langle \omega(f), \omega(g) \rangle = \langle f, \omega^2(g) \rangle = \langle f, g \rangle$ for all $f, g \in \Lambda$.

1.4 Skew-Schur Functions and Skewing

Definition: For two partitions λ, μ , the *skew Schur function* $s_{\lambda/\mu}(\bar{x})$ is defined by

$$s_{\lambda/\mu}(\bar{x}) = \sum_{\text{SSYT } T \text{ of shape } \lambda/\mu} \bar{x}^T$$

where the summation is taken over all semistandard Young tableau of shape λ/μ .

Proposition: For partitions λ, μ , we have that

$$s_{\lambda/\mu}(x_1, \dots, x_n) = \det \left(h_{\lambda_i - \mu_j + j - i}(\bar{x}) \right)_{1 \leq i, j \leq n},$$

$$s_{\lambda'/\mu'}(x_1, \dots, x_n) = \det \left(e_{\lambda_i - \mu_j + j - i}(\bar{x}) \right)_{1 \leq i, j \leq n}.$$

Definition: For any symmetric function $f \in \Lambda$, the *skewing operator* $f^\perp : \Lambda \rightarrow \Lambda$ is defined by the property that for all $g, h \in \Lambda$ we have that $\langle f \cdot g, h \rangle = \langle g, f^\perp(h) \rangle$.

The discussion on skewing follows [GR26, Chapter 2].

Example: We have that $s_\mu^\perp(s_\lambda) = s_{\lambda/\mu}$.

Example: We have “dual versions” of the Pieri rules. For a partition λ and an integer $r \geq 0$, we have that

$$h_r^\perp(s_\lambda(\bar{x})) = \sum_{\mu} s_\mu$$

where the summation over μ is over all partitions μ such that λ/μ is a *horizontal strip* meaning no two boxes in the skew diagram are in the same column. We also have that

$$e_r^\perp(s_\lambda(\bar{x})) = \sum_{\mu} s_\mu$$

where the summation over μ is over all partitions μ such that λ/μ is a *vertical strip* meaning no two boxes in the skew diagram are in the same row.

2 The Affine Symmetric Group

We give multiple presentations of the affine symmetric group $\tilde{\mathfrak{S}}_n$. This mostly follows [Lam+14, Chapter 1, §2] but [BB05, Chapter 8, §3] is another phenomenal source.

Definition: The *affine symmetric group* $\tilde{\mathfrak{S}}_n$ is the Coxeter group generated by $\{s_0, \dots, s_{n-1}\}$ such that

$$s_i^2 = e$$

$$s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$$

$$s_i s_j = s_j s_i \text{ for } i - j \not\equiv 0, 1, n-1 \pmod{n},$$

where all of the above indices are taken $\pmod{n}$. Thus, we have a notion of the length ℓ of a word in $\tilde{\mathfrak{S}}_n$.

Equivalently, $aff S_n$ is the group of permutations $w : \mathbb{Z} \rightarrow \mathbb{Z}$ such that for all $k \in \mathbb{Z}$, $w(i + kn) = w(i) + kn$ and $\sum_{i=1}^n w(i) = \sum_{i=1}^n i$. Thus, w is completely determined by $w(1), \dots, w(n)$. Each s_i can be identified with the permutation swapping i and $i + 1$.

Since w is determined by $w(1), \dots, w(n)$, we express $w = [w(1), \dots, w(n)]$ in *window notation*. We have an action of the s_i on a window $[w_1, \dots, w_n]$:

$$[w_1, \dots, w_n] \cdot s_i = [w_1, \dots, w_{i+1}, w_i, \dots, w_n], \quad 1 \leq i \leq n-1,$$

$$[w_1, \dots, w_n] \cdot s_0 = [w_n - n, w_2, \dots, w_{n-1}, w_1 + n].$$

2.1 Affine Grassmannian Elements

Proposition: For $w \in \tilde{\mathfrak{S}}_n$, the length $\ell(w)$ is given by

$$\# \{ (i, j) \in [n] \times [n] \mid i < j, w_i > w_j \} + \sum_{1 \leq i < j \leq n} \left\lfloor \frac{|w_i - w_j|}{n} \right\rfloor.$$

Definition: We have a copy of $\mathfrak{S}_n \subseteq \tilde{\mathfrak{S}}_n$ generated by $\{s_1, \dots, s_{n-1}\}$. The coset representatives of $\tilde{\mathfrak{S}}_n/\mathfrak{S}_n$ are given by *affine grassmannian* elements $w \in \tilde{\mathfrak{S}}_n$. These will be the minimal length representatives of each coset in $\tilde{\mathfrak{S}}_n/\mathfrak{S}_n$. Equivalently, these are the w such that $w(1) < \dots < w(n)$.

2.2 Types of Partitions

Definition: A partition λ is k -bounded if $\lambda_1 \leq k$.

The *hook length* of a cell in a tableau is the number of cells directly below and the number of cells directly to the right, including the cell itself.

A partition κ is an r -core if it has no cells of hook length equal to r . The *size* of κ , $|\kappa|_r$ is the number of cells of hook length less than r .

A k -skew diagram is a skew-partition κ/ρ such that no cell in κ/ρ has hook-length exceeding k and all the cells where ρ would be in the skew-partition have hook length exceeding k .

2.3 Bijections Fix some $n \geq 1$ and throughout let k such that $n = k + 1$. All our examples will focus on $k = 3$, $n = 4$, and $m = 8$.

Proposition ([Lam + 14, Proposition 1.3]): There is a bijection between the set of $(k + 1)$ -cores κ with $|\kappa|_{k+1} = m$ and k -bounded partitions $\lambda \vdash m$.

The map sending a $(k + 1)$ -core to a k -bounded partition is done by simply removing the cells with hook length $\geq k + 1$ and realigning the rows to form a partition. The map sending a k -bounded partition to a $(k + 1)$ -core is computed by the following procedure. Starting from the smallest row in the diagram and moving upwards to the largest row, slide cells in the row (and thus the above rows) to the right until there are no hooks of length at least $k + 1$ in the current row.

Example:



Figure 1: Map sending the $(3 + 1)$ -core $\kappa = (6, 3, 2, 1)$ to the 3-bounded partition $\lambda = (3, 2, 2, 1)$. Cells marked with a red X are the cells eliminated.




Figure 2: Map sending the 3-bounded partition $\lambda = (3, 2, 2, 1)$ to the $(3 + 1)$ -core $\kappa = (6, 3, 2, 1)$. Cells outlined in red are the cells shifted to the right.

In fact, the skew-partition formed by taking the core κ and eliminating all cells with hook length $\geq k + 1$ gives a k -skew diagram. Then “filling in” the k -skew diagram gives us a $(k + 1)$ -core. We record this bijection.

Proposition ([LM04; LM05]): There is a bijection between the $(k + 1)$ -cores κ $\langle\langle$ with $|\kappa|_{k+1} = m \rangle\rangle$ and the k -skew diagrams with $\langle\langle$ something about $m \rangle\rangle$.

We also get a bijection from the k -skew diagrams to the k -bounded partitions. Left-alignment of the rows of the k -skew diagram gives us a k -bounded partition. To go from a k -bounded partition λ to a k -skew diagram, work from the smallest row to the largest row of λ , adding rows to the left-most position that does not form any hooks of length exceeding $k + 1$.

Proposition ([LM04; LM05]): There is a bijection between the k -bounded partitions λ $\langle\langle$ with $|\lambda| = m \rangle\rangle$ and the k -skew diagrams with $\langle\langle$ something about $m \rangle\rangle$.

Example:




Figure 3: An example of the bijection between $(k + 1)$ -cores, k -skew diagrams, and k -bounded partitions.

Proposition ([Lam+14, Proposition 1.9]): There is a bijection between the affine Grassmannian elements of $\tilde{\mathfrak{S}}_n$ with length m and the $(k + 1)$ -cores κ with $|\kappa|_{k+1} = m$.

Definition: For a cell with coordinate (i, j) of a Young diagram, its *content* or *diagonal index* is the value $j - i$. For a fixed n , its *residue* modulo n is the content modulo n .

Example:



Figure 4: Content and residue modulo $3 + 1$ of the 3-bounded partition $\lambda = (3, 2, 2, 1)$.

We define an action of $\tilde{\mathfrak{S}}_n$ on $(k + 1)$ -cores.

Definition: Let κ be a $(k + 1)$ -core and let $0 \leq i \leq n - 1$. Then $s_i \cdot \kappa$ is the $(k + 1)$ -core such that

- If a cell with residue i can be added to κ such that the resulting diagram is still the diagram of a partition, add all such cells,
- If a cell with residue i can be removed from κ such that the resulting diagram is still the diagram of a partition, remove all such cells,
- Else, leave κ as-is.

Note that the first two conditions can never happen simultaneously.

We now give the bijection between $(k + 1)$ -cores κ of size m and affine grassmannian elements $w \in \tilde{\mathfrak{S}}_n$ of length m . To go from an affine grassmannian element w of length m to a $(k + 1)$ -core κ of size m , use the above action of $\tilde{\mathfrak{S}}_n$ starting

