



# Representation Theory Notes and Exercises

With 0 Figures

Anakin Dey

Last Edited on 12/17/23 at 16:49

## TODOs

⟨⟨ Change the style of enumerates from “1.” to “(1)” ⟩⟩

⟨⟨ Proper Exercise Header ⟩⟩

⟨⟨ Proper Chapter Header ⟩⟩

⟨⟨ Remove indent at end of thmtool environments ⟩⟩

# Contents

1	Generalities on Linear Representations	1
---	--	---

# List of Definitions

1.1	Linear Representation, Representation Space . . . . .	1
1.2	Degree . . . . .	1
1.3	Matrix of a Representation . . . . .	1
1.4	Similar/Isomorphic Representations . . . . .	2
1.8	Stable/Invariant Subspaces, Subrepresentation . . . . .	3
1.11	Direct Sum of Representations . . . . .	4

# List of Examples and Counterexamples

1.5	Unit/Trivial Representation . . . . .	2
1.6	Regular Representation . . . . .	2
1.7	Permutation Representation . . . . .	3
1.9	Subrepresentations of the Regular Representation . . . . .	3

# Preface

This is a set of notes on group representation theory mainly based on J.P. Serre's text *Linear Representations of Finite Groups* [Ser77]. Occasionally, other sources may be used, such as the set of notes by Charles Rezk created for Math 427 [Rez20]. The goal of these notes is to eventually work towards algebraic combinatorics such as Fulton's text *Young Tableaux* [Ful96], as much of algebraic combinatorics is motivated by questions stemming from representation theory. At the time of writing this, another goal is the applications of representation theory to computational complexity: see [Pan23] for a recent survey on this connection.

# Chapter 1

## Generalities on Linear Representations

Unless otherwise specified,  $V$  will denote a vector space, usually over the field  $\mathbb{C}$ . We will restrict ourselves to finite dimensional vector spaces. Similarly, we will restrict ourselves to finite groups.

**Definition 1.1 (Linear Representation, Representation Space):** Let  $G$  be a group with identity  $e$ . A *linear representation* of  $G$  in  $V$  is a homomorphism  $\rho : G \rightarrow \text{GL}(V)$ . We will frequently, and often interchangeably, write  $\rho_s := \rho(s)$ . Given  $\rho$ , we will say that  $V$  is a *representation space* or *representation* of  $G$ .

**Definition 1.2 (Degree):** Let  $\rho : G \rightarrow V$  be a representation of  $G$  in a vector space  $V$ . Then the *degree* of  $\rho$  is  $\dim(V)$ .

Let  $\rho : G \rightarrow V$  be a representation of  $G$  in a vector space  $V$  with  $n := \dim(V)$ . Fix a basis  $(e_j)$  of  $V$ . Then since each  $\rho_s$  is an invertible linear transformation of  $V$ , we may define an  $n \times n$  matrix  $R_s \equiv (r_{ij}(s))$  where each  $r_{ij}(s)$  is defined by the identity

$$\rho_s(e_j) = \sum_{i=1}^n r_{ij}(s) e_i.$$

**Definition 1.3 (Matrix of a Representation):** We call  $R_s = (r_{ij}(s))$  above the *matrix of  $\rho_s$*  with respect to the basis  $(e_j)$ .

Note that  $R_s$  satisfies the following:

$$\det(R_s) \neq 0, \quad R_{st} = R_s \cdot R_t \equiv r_{ij}(st) = \sum_{k=1}^n r_{ik}(s) \cdot r_{kj}(s) \quad \forall s, t \in G.$$

Recall that two  $n \times n$  matrices  $A, A'$  are *similar* if there exists an invertible matrix  $T$  such that  $TA = A'T$ . We may extend this notion to representations.

**Definition 1.4 (Similar/Isomorphic Representations):** Let  $\rho$  and  $\rho'$  be two representations of the same group  $G$  in vector spaces  $V$  and  $V'$  respectively. We say  $\rho$  and  $\rho'$  are *similar* or *isomorphic* if there exists an isomorphism  $\tau: V \rightarrow V'$  such that for all  $s \in G$ ,  $\tau$  satisfies  $\tau \circ \rho(s) = \rho'(s) \circ \tau$ . If  $R_s, R'_s$  are the corresponding matrices then this is equivalent to saying there exists an invertible matrix  $T$  such that  $TR_s = R'_s T$  for all  $s \in G$ .

Note that if  $\rho$  and  $\rho'$  are isomorphic, then they must have the same degree.

We now give some examples of these things.

**Example 1.5 (Unit/Trivial Representation):** Let  $G$  be a finite group. Representations of degree 1 must be of the form  $\rho: G \rightarrow \mathbb{C}^\times$ . Since elements  $s$  of  $G$  are of finite order,  $\rho(s)$  must also be of finite order. Thus, for all  $s \in G$ ,  $\rho(s)$  is a root of unity. If we take  $\rho(s) = 1$  for all  $s \in G$ , we obtain the *unit* or *trivial* representation of  $G$ . This also means that  $R_s = 1$  for all  $s$ .

**Example 1.6 (Regular Representation):** Let  $g$  be the order of  $G$ , and let  $V$  be a vector space of dimension  $g$  with a basis  $(e_t)_{t \in G}$ . For each  $s \in G$ , define  $\rho_s$  as the linear map  $\rho_s: V \rightarrow V$  such that  $\rho_s(e_t) = e_{st}$ . This is a linear representation of  $G$  called the *regular* representation of  $G$ . Since for each  $s \in G$ ,  $e_s = \rho_s(e_1)$  and thus the images of  $e_1$  form a basis of  $V$ . Conversely, let  $W$  be a representation of  $G$  with a vector  $w$  satisfying the collection of all  $\rho_s(w)$ ,  $s \in G$ , forms a basis of  $W$ . Then  $W$  is isomorphic to the regular representation of  $G$  by the isomorphism  $\tau(e_s) = \rho_s(w)$ .

For example, let  $G = \mathbb{Z}_3$  and  $V = \mathbb{C}^3$  with  $e_0 = (1, 0, 0)$ ,  $e_1 = (0, 1, 0)$ , and  $e_2 = (0, 0, 1)$ . Then for example,  $\rho_0, \rho_1, \rho_2: \mathbb{C}^3 \rightarrow \mathbb{C}^3$  are the linear maps such that

$$\begin{array}{lll} \rho_0(e_0) = e_{0+0} = e_0 & \rho_0(e_1) = e_{0+1} = e_1 & \rho_0(e_2) = e_{0+2} = e_2 \\ \rho_1(e_0) = e_{1+0} = e_1 & \rho_1(e_1) = e_{1+1} = e_2 & \rho_1(e_2) = e_{1+2} = e_0 \\ \rho_2(e_0) = e_{2+0} = e_2 & \rho_2(e_1) = e_{2+1} = e_0 & \rho_2(e_2) = e_{2+2} = e_1 \end{array}$$

With this, the matrix representations of  $\rho_0, \rho_1$  and  $\rho_2$  is similarly straightforward:

$$R_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad R_1 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad R_2 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$



**Example 1.7 (Permutation Representation):** We may generalize the regular representation to any group action  $G \curvearrowright X$ ,  $X$  a finite set. Recall that for such an action, the map  $x \mapsto sx$  for each  $s \in G$  is a permutation  $X \leftrightarrow X$ . Let  $V$  be a vector space with dimension the size of  $X$ , and so a basis  $(e_x)_{x \in X}$ . Define a representation  $\rho$  of  $G$  by defining  $\rho_s$  as the linear map sending  $e_x \mapsto e_{sx}$ . This representation is known as the *Permutation* representation of  $G$  associated with  $X$ . If we consider  $X = [n]$  and  $G = S_n$ , then take  $V = \mathbb{C}^n$  as our vector space and  $e_i$  as the standard basis vector. Then  $\rho_\sigma(e_j) = e_{\sigma_j}$ . Thus for each  $\sigma \in S_n$ , we have that  $R_\sigma = (r_{ij}(\sigma))$  where entry  $r_{ij}(\sigma) = 1$  if  $i = \sigma_j$  and 0 otherwise.

**Definition 1.8 (Stable/Invariant Subspaces, Subrepresentation):** Let  $\rho : G \rightarrow \text{GL}(V)$  be a linear representation and  $W \subseteq V$  a subspace of  $V$ . We say that  $W$  is *stable* under the action of  $G$  if  $x \in W$  implies that  $\rho_s(x) \in W$  for all  $s \in G$ . Thus, the restriction  $\rho_s^W := \rho_s|_W$  is an isomorphism of  $W$  onto itself. Restrictions satisfy the property that  $\rho_s^W \circ \rho_t^W = \rho_{st}^W$ . Thus,  $\rho^W : G \rightarrow \text{GL}(W)$  is a linear representation of  $G$  in  $W$  and we say that  $W$  is a *subrepresentation* of  $V$ .

**Example 1.9 (Subrepresentations of the Regular Representation):** Let  $G$  be a group. Recall the regular representation  $V$  given in Example 1.6. Let  $W$  be the 1 dimensional subspace of  $V$  generated by the element  $x = \sum_{s \in G} e_s$ . Then note that  $\rho_s(x) = x$  for all  $s \in G$  and thus  $W$  is a subrepresentation of  $V$ . Furthermore, this is isomorphic to the unit representation Example 1.5 with  $\tau : \mathbb{C}^\times \rightarrow W$  such that  $\tau(1) = x$ . For example, let  $G = Z_3$  and  $\rho : Z_3 \rightarrow \mathbb{C}^3$  the representation given in Example 1.6. Then  $x = (1, 1, 1)$  and for example we have that

$$\rho_1(x) = \rho(1)(e_0) + \rho_1(e_1) + \rho_1(e_2) = e_1 + e_2 + e_0 = x.$$

**Theorem 1.10:** Let  $\rho : G \rightarrow \text{GL}(V)$  be a linear representation of  $G$  in  $V$  and let  $W$  be a subspace of  $V$  stable under  $G$ . Then there exists a complement  $W^0$  of  $W$  in  $V$  which is stable under  $G$ .

**Proof:** Let  $W'$  be an arbitrary complement of  $W$  in  $V$ , and let  $p : V \rightarrow W$  be the projection. Then we form the average  $p^0$  of conjugates of  $p$  by elements in  $G$ :

$$p^0 := \frac{1}{|G|} \sum_{t \in G} \rho_t \circ p \circ \rho_t^{-1}.$$

Since  $p : V \rightarrow W$  and  $\rho_t$  preserves  $W$ , we have that  $p^0$  maps  $V$  onto  $W$ . Furthermore, note that  $\rho_t^{-1}$  also preserves  $W$ .

Thus we have that

$$(p \circ \rho_t^{-1})(x) = \rho_t^{-1}(x), \quad (\rho_t \circ p \circ \rho_t^{-1})(x) = x, \quad p^0(x) = x.$$

Thus,  $p^0$  is a projection of  $V$  onto  $W$ , corresponding to some complement  $W^0$  of  $W$ . Moreover, we have that  $\rho_s \circ p^0 = p^0 \circ \rho_s$  for all  $s \in G$  because

$$\rho_s \circ p^0 \circ \rho_s^{-1} = \frac{1}{|G|} \sum_{t \in G} \rho_s \circ \rho_t \circ p \circ \rho_t^{-1} \circ \rho_s^{-1} = \frac{1}{|G|} \sum_{t \in G} \rho_{st} \circ p \rho_{st}^{-1} = p^0.$$

Now suppose that  $x \in W^0$  and  $s \in G$ , we have that  $p^0(x) = 0$  and hence  $(p^0 \circ \rho_s)(x) = (\rho_s \circ p^0)(x) = 0$ , meaning that  $\rho_s(x) \in W^0$ . This,  $W^0$  is stable under  $G$ .  $\square$

Suppose that  $V$  had an innerproduct  $\langle x, y \rangle$ , and furthermore suppose this inner product was invariant under  $G$  meaning that for all  $s \in G$ ,  $\langle \rho_s(x), \rho_s(y) \rangle = \langle x, y \rangle$ . We may also reduce to this case by replacing  $\langle x, y \rangle$  with  $\sum_{t \in G} \langle \rho_t(x), \rho_t(y) \rangle$ . With this, the orthogonal complement  $W^\perp$  of  $W$  in  $V$  is a complement of  $W$  stable under  $G$ . Note that the invariance of  $\langle x, y \rangle$  means that if  $(e_i)$  is an orthonormal basis of  $V$ , then  $R_s$  is a unitary matrix  $\langle \langle \text{proof?} \rangle \rangle$ .

Using the notation of Theorem 1.10, let  $x \in V$  and  $w, w^0$  be the projections of  $x$  on  $W$  and  $W^0$  respectively. Thus for all  $s \in G$ ,  $\rho_s(x) = \rho_s(w) + \rho_s(w^0)$ . Since  $W$  and  $W^0$  are stable under  $G$ , we have that  $\rho_s(w) \in W$  and  $\rho_s(w^0) \in W^0$ . This means that  $\rho_s(w)$  and  $\rho_s(w^0)$  are the projections of  $\rho_s(x)$  and in turn the representations of  $W$  and  $W^0$  determine the representations of  $V$ .

**Definition 1.11 (Direct Sum of Representations):** Given the above, we write  $V = W \oplus W^0$  as the *direct sum* of  $W$  and  $W^0$ . We identify elements  $v \in V$  as pairs  $(w, w^0)$  given by their projections.

If the representations  $W$  and  $W^0$  are given in matrices  $R_s$  and  $R_s^0$ , then the matrix form of the representation  $V$  is given by

$$\begin{pmatrix} R_s & 0 \\ 0 & R_s^0 \end{pmatrix}.$$

Similar results hold for arbitrarily many, but finite, direct sums of representations.

# Bibliography

- [Ful96] William Fulton. *Young Tableaux: With Applications to Representation Theory and Geometry*. London Mathematical Society Student Texts. Cambridge University Press, 1996. DOI: [10.1017/CB09780511626241](https://doi.org/10.1017/CB09780511626241).
- [Pan23] Greta Panova. *Computational Complexity in Algebraic Combinatorics*. 2023. arXiv: [2306.17511](https://arxiv.org/abs/2306.17511) [math.CO].
- [Rez20] Charles Rezk. *A short course on finite group representations*. 2020. URL: <https://rezk.web.illinois.edu/Finite%5C%20Group%5C%20Reps/short-course-finite-group-representations.html>.
- [Ser77] Jean-Pierre Serre. *Linear Representations of Finite Groups*. Springer New York, 1977. ISBN: 9781468494587. DOI: [10.1007/978-1-4684-9458-7](https://doi.org/10.1007/978-1-4684-9458-7).