



Representation Theory Notes and Exercises

With 0 Figures

Anakin Dey

Last Edited on 12/19/23 at 00:09

TODOs

⟨⟨ Change the style of enumerates from “1.” to “(1)” ⟩⟩

⟨⟨ Proper Exercise Header ⟩⟩

⟨⟨ Proper Chapter Header ⟩⟩

⟨⟨ Remove indent at end of thmtool environments ⟩⟩

Contents

1	Generalities on Linear Representations	1
2	Character Theory	7

List of Definitions

1.1	Linear Representation, Representation Space	1
1.2	Degree	1
1.3	Matrix of a Representation	1
1.4	Similar/Isomorphic Representations	2
1.8	Stable/Invariant Subspaces, Subrepresentation	3
1.11	Direct Sum of Representations	4
1.12	Irreducible/Simple Representations	4
1.15	Tensor/Kronecker Product of Representations	5
1.16	Symmetric Square, Alternating Square	6
2.1	Character	7

List of Examples and Counterexamples

1.5	Unit/Trivial Representation	2
1.6	Regular Representation	2
1.7	Permutation Representation	3
1.9	Subrepresentations of the Regular Representation	3
1.14	Decomposition of Representation of \mathbb{Z}_3 into Irreducibles	5

Preface

This is a set of notes on group representation theory mainly based on J.P. Serre's text *Linear Representations of Finite Groups* [Ser77]. Occasionally, other sources may be used, such as the set of notes by Charles Rezk created for Math 427 [Rez20]. The goal of these notes is to eventually work towards algebraic combinatorics such as Fulton's text *Young Tableaux* [Ful96], as much of algebraic combinatorics is motivated by questions stemming from representation theory. At the time of writing this, another goal is the applications of representation theory to computational complexity: see [Pan23] for a recent survey on this connection.

Chapter 1

Generalities on Linear Representations

Unless otherwise specified, V will denote a vector space, usually over the field \mathbb{C} . We will restrict ourselves to finite dimensional vector spaces. Similarly, we will restrict ourselves to finite groups.

Definition 1.1 (Linear Representation, Representation Space): Let G be a group with identity e . A *linear representation* of G in V is a homomorphism $\rho : G \rightarrow \text{GL}(V)$. We will frequently, and often interchangeably, write $\rho_s := \rho(s)$. Given ρ , we will say that V is a *representation space* or *representation* of G .

Definition 1.2 (Degree): Let $\rho : G \rightarrow V$ be a representation of G in a vector space V . Then the *degree* of ρ is $\dim(V)$.

Let $\rho : G \rightarrow V$ be a representation of G in a vector space V with $n := \dim(V)$. Fix a basis (e_j) of V . Then since each ρ_s is an invertible linear transformation of V , we may define an $n \times n$ matrix $R_s \equiv (r_{ij}(s))$ where each $r_{ij}(s)$ is defined by the identity

$$\rho_s(e_j) = \sum_{i=1}^n r_{ij}(s) e_i.$$

Definition 1.3 (Matrix of a Representation): We call $R_s = (r_{ij}(s))$ above the *matrix of ρ_s* with respect to the basis (e_j) .

Note that R_s satisfies the following:

$$\det(R_s) \neq 0, \quad R_{st} = R_s \cdot R_t \equiv r_{ij}(st) = \sum_{k=1}^n r_{ik}(s) \cdot r_{kj}(s) \quad \forall s, t \in G.$$

Recall that two $n \times n$ matrices A, A' are *similar* if there exists an invertible matrix T such that $TA = A'T$. We may extend this notion to representations.

Definition 1.4 (Similar/Isomorphic Representations): Let ρ and ρ' be two representations of the same group G in vector spaces V and V' respectively. We say ρ and ρ' are *similar* or *isomorphic* if there exists an isomorphism $\tau: V \rightarrow V'$ such that for all $s \in G$, τ satisfies $\tau \circ \rho(s) = \rho'(s) \circ \tau$. If R_s, R'_s are the corresponding matrices then this is equivalent to saying there exists an invertible matrix T such that $TR_s = R'_s T$ for all $s \in G$.

Note that if ρ and ρ' are isomorphic, then they must have the same degree.

We now give some examples of these things.

Example 1.5 (Unit/Trivial Representation): Let G be a finite group. Representations of degree 1 must be of the form $\rho: G \rightarrow \mathbb{C}^\times$. Since elements s of G are of finite order, $\rho(s)$ must also be of finite order. Thus, for all $s \in G$, $\rho(s)$ is a root of unity. If we take $\rho(s) = 1$ for all $s \in G$, we obtain the *unit* or *trivial* representation of G . This also means that $R_s = 1$ for all s .

Example 1.6 (Regular Representation): Let g be the order of G , and let V be a vector space of dimension g with a basis $(e_t)_{t \in G}$. For each $s \in G$, define ρ_s as the linear map $\rho_s: V \rightarrow V$ such that $\rho_s(e_t) = e_{st}$. This is a linear representation of G called the *regular* representation of G . Since for each $s \in G$, $e_s = \rho_s(e_1)$ and thus the images of e_1 form a basis of V . Conversely, let W be a representation of G with a vector w satisfying the collection of all $\rho_s(w)$, $s \in G$, forms a basis of W . Then W is isomorphic to the regular representation of G by the isomorphism $\tau(e_s) = \rho_s(w)$.

For example, let $G = \mathbb{Z}_3$ and $V = \mathbb{C}^3$ with $e_0 = (1, 0, 0)$, $e_1 = (0, 1, 0)$, and $e_2 = (0, 0, 1)$. Then for example, $\rho_0, \rho_1, \rho_2: \mathbb{C}^3 \rightarrow \mathbb{C}^3$ are the linear maps such that

$$\begin{array}{lll} \rho_0(e_0) = e_{0+0} = e_0 & \rho_0(e_1) = e_{0+1} = e_1 & \rho_0(e_2) = e_{0+2} = e_2 \\ \rho_1(e_0) = e_{1+0} = e_1 & \rho_1(e_1) = e_{1+1} = e_2 & \rho_1(e_2) = e_{1+2} = e_0 \\ \rho_2(e_0) = e_{2+0} = e_2 & \rho_2(e_1) = e_{2+1} = e_0 & \rho_2(e_2) = e_{2+2} = e_1 \end{array}$$

With this, the matrix representations of ρ_0, ρ_1 and ρ_2 is similarly straightforward:

$$R_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad R_1 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad R_2 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$

Example 1.7 (Permutation Representation): We may generalize the regular representation to any group action $G \curvearrowright X$, X a finite set. Recall that for such an action, the map $x \mapsto sx$ for each $s \in G$ is a permutation $X \leftrightarrow X$. Let V be a vector space with dimension the size of X , and so a basis $(e_x)_{x \in X}$. Define a representation ρ of G by defining ρ_s as the linear map sending $e_x \mapsto e_{sx}$. This representation is known as the *Permutation* representation of G associated with X . If we consider $X = [n]$ and $G = S_n$, then take $V = \mathbb{C}^n$ as our vector space and e_i as the standard basis vector. Then $\rho_\sigma(e_j) = e_{\sigma_j}$. Thus for each $\sigma \in S_n$, we have that $R_\sigma = (r_{ij}(\sigma))$ where entry $r_{ij}(\sigma) = 1$ if $i = \sigma(j)$ and 0 otherwise.

Definition 1.8 (Stable/Invariant Subspaces, Subrepresentation): Let $\rho: G \rightarrow \text{GL}(V)$ be a linear representation and $W \subseteq V$ a subspace of V . We say that W is *stable* under the action of G if $x \in W$ implies that $\rho_s(x) \in W$ for all $s \in G$. Thus, the restriction $\rho_s^W := \rho_s|_W$ is an isomorphism of W onto itself. Restrictions satisfy the property that $\rho_s^W \circ \rho_t^W = \rho_{st}^W$. Thus, $\rho^W: G \rightarrow \text{GL}(W)$ is a linear representation of G in W and we say that W is a *subrepresentation* of V .

Example 1.9 (Subrepresentations of the Regular Representation): Let G be a group. Recall the regular representation V given in Example 1.6. Let W be the 1 dimensional subspace of V generated by the element $x = \sum_{s \in G} e_s$. Then note that $\rho_s(x) = x$ for all $s \in G$ and thus W is a subrepresentation of V . Furthermore, this is isomorphic to the unit representation Example 1.5 with $\tau: C^\times \rightarrow W$ such that $\tau(1) = x$. For example, let $G = \mathbb{Z}_3$ and $\rho: \mathbb{Z}_3 \rightarrow \text{GL}(\mathbb{C}^3)$ the representation given in Example 1.6. Then $x = (1, 1, 1)$ and for example we have that

$$\rho_1(x) = \rho_1(e_0) + \rho_1(e_1) + \rho_1(e_2) = e_1 + e_2 + e_0 = x.$$

Theorem 1.10: Let $\rho: G \rightarrow \text{GL}(V)$ be a linear representation of G in V and let W be a subspace of V stable under G . Then there exists a complement W^0 of W in V which is stable under G .

Proof: Let W' be an arbitrary complement of W in V , and let $p: V \rightarrow W$ be the projection. Then we form the average p^0 of conjugates of p by elements in G :

$$p^0 := \frac{1}{|G|} \sum_{t \in G} \rho_t \circ p \circ \rho_t^{-1}.$$

Since $p: V \rightarrow W$ and ρ_t preserves W , we have that p^0 maps V onto W . Furthermore, note that ρ_t^{-1} also preserves W .

Thus we have that

$$(p \circ \rho_t^{-1})(x) = \rho_t^{-1}(x), \quad (\rho_t \circ p \circ \rho_t^{-1})(x) = x, \quad p^0(x) = x.$$

Thus, p^0 is a projection of V onto W , corresponding to some complement W^0 of W . Moreover, we have that $\rho_s \circ p^0 = p^0 \circ \rho_s$ for all $s \in G$ because

$$\rho_s \circ p^0 \circ \rho_s^{-1} = \frac{1}{|G|} \sum_{t \in G} \rho_s \circ \rho_t \circ p \circ \rho_t^{-1} \circ \rho_s^{-1} = \frac{1}{|G|} \sum_{t \in G} \rho_{st} \circ p \rho_{st}^{-1} = p^0.$$

Now suppose that $x \in W^0$ and $s \in G$, we have that $p^0(x) = 0$ and hence $(p^0 \circ \rho_s)(x) = (\rho_s \circ p^0)(x) = 0$, meaning that $\rho_s(x) \in W^0$. This, W^0 is stable under G . \square

Suppose that V had an innerproduct $\langle x, y \rangle$, and furthermore suppose this inner product was invariant under G meaning that for all $s \in G$, $\langle \rho_s(x), \rho_s(y) \rangle = \langle x, y \rangle$. We may also reduce to this case by replacing $\langle x, y \rangle$ with $\sum_{t \in G} \langle \rho_t(x), \rho_t(y) \rangle$. With this, the orthogonal complement W^\perp of W in V is a complement of W stable under G . Note that the invariance of $\langle x, y \rangle$ means that if (e_i) is an orthonormal basis of V , then R_s is a unitary matrix.

Using the notation of Theorem 1.10, let $x \in V$ and w, w^0 be the projections of x on W and W^0 respectively. Thus for all $s \in G$, $\rho_s(x) = \rho_s(w) + \rho_s(w^0)$. Since W and W^0 are stable under G , we have that $\rho_s(w) \in W$ and $\rho_s(w^0) \in W^0$. This means that $\rho_s(w)$ and $\rho_s(w^0)$ are the projections of $\rho_s(x)$ and in turn the representations of W and W^0 determine the representations of V .

Definition 1.11 (Direct Sum of Representations): Given the above, we write $V = W \oplus W^0$ as the *direct sum* of W and W^0 . We identify elements $v \in V$ as pairs (w, w^0) given by their projections.

If the representations W and W^0 are given in matrices R_s and R_s^0 , then the matrix form of the representation V is given by

$$\begin{pmatrix} R_s & 0 \\ 0 & R_s^0 \end{pmatrix}.$$

Similar results hold for arbitrarily many, but finite, direct sums of representations.

The above is a discussion on how to compose two or more representations into one larger representation. The natural question is then about the opposite.

Definition 1.12 (Irreducible/Simple Representations): Let $\rho : G \rightarrow \text{GL}(V)$ be a linear representation of G . Then this representation is *irreducible* or *simple* if V has no subspaces stable under G besides 0 and V itself.

By Theorem 1.10, this is equivalent to saying that V is not the direct sum of two representations besides $V = 0 \oplus V$. A representation of degree 1 is reducible. We may use irreducible representations to construct other ones via the direct sum.

Theorem 1.13: Every representation is a direct sum of irreducible representations.

Proof: Let V be a linear representation of G . We induct on $\dim(V)$. If $\dim(V) = 0$, then $V = 0$ which is the direct sum of an empty family of irreducible representations. So suppose that

$\dim(V) \geq 1$. If V is irreducible, then we are done. Otherwise, there exists a subspace $W \subsetneq V$ stable under G and by Theorem 1.10 a stable complement W^0 such that $V = W \oplus W^0$. By assumption, $W \neq 0 \neq W^0$ and so $\dim(W) < \dim(V)$ and $\dim(W^0) < \dim(V)$. By induction, we have obtained a decomposition of V into irreducibles. \square

Example 1.14 (Decomposition of Representation of \mathbb{Z}_3 into Irreducibles): Recall from Example 1.6 the regular representation $\rho : \mathbb{Z}_3 \rightarrow \text{GL}(\mathbb{C}^3)$ with $e_0 = (1, 0, 0)$, $e_1 = (0, 1, 0)$, and $e_2 = (0, 0, 1)$ and

$$\begin{array}{lll} \rho_0(e_0) = e_0 & \rho_0(e_1) = e_1 & \rho_0(e_2) = e_2 \\ \rho_1(e_0) = e_1 & \rho_1(e_1) = e_2 & \rho_1(e_2) = e_0 \\ \rho_2(e_0) = e_2 & \rho_2(e_1) = e_0 & \rho_2(e_2) = e_1 \end{array}$$

Our goal will be to decompose ρ into $\rho^1 \oplus \rho^2 \oplus \rho^3$. We aim to find the elements fixed by \mathbb{Z}_3 . Note that if an element is fixed by 1, the generator of \mathbb{Z}_3 , then it is fixed by all of \mathbb{Z}_3 . We want to find 1-dimensional \mathbb{Z}_3 -invariant subspaces of \mathbb{C}^3 . This is equivalent to finding the eigenvalues of the matrix

$$R_1 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

The eigenvalues and their eigenvectors of R_1 are as follows, note the inclusion of the trivial subrepresentation from Example 1.9:

$$\lambda_1 = 1, v_1 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad \lambda_2 = \frac{-1 - i\sqrt{3}}{2}, v_2 = \begin{pmatrix} \frac{-1+i\sqrt{3}}{2} \\ \frac{-1-i\sqrt{3}}{2} \\ 1 \end{pmatrix} \quad \lambda_3 = \frac{-1 + i\sqrt{3}}{2}, v_3 = \begin{pmatrix} \frac{-1-i\sqrt{3}}{2} \\ \frac{-1+i\sqrt{3}}{2} \\ 1 \end{pmatrix}$$

Thus $\mathbb{C}^3 = V_1 \oplus V_2 \oplus V_3$ where $V_i := \text{span}(v_i)$. Note that there are only 3 morphisms $\mathbb{Z}_3 \rightarrow \mathbb{C}^\times$ mapping 1 to 1, ω , or ω^2 where ω is a cube root of unity. Thus ρ^1, ρ^2 , and ρ^3 must correspond to these morphisms **<< but which ones >>**.

A natural question is if such a decomposition $V = W_1 \oplus \cdots \oplus W_k$ is unique. However, suppose that ρ is the trivial representation (Example 1.5). Then each component of the decomposition is a line and we can decompose a vector space into the direct sum of lines in a number of ways. However, we will see that the number of W_i that are isomorphic to a given irreducible representation does not depend on the choice of decomposition.

Definition 1.15 (Tensor/Kronecker Product of Representations): Let $\rho^1: G \rightarrow \text{GL}(V_1)$ and $\rho^2: G \rightarrow \text{GL}(V_2)$ be two representations of a group G . We construct a representation $\rho: G \rightarrow \text{GL}(V_1 \otimes V_2)$ such that

$$\rho_s(x_1 \otimes x_2) = \rho_s^1(x_1) \circ \rho_s^2(x_2) \quad \text{for } x_1 \in V_1, x_2 \in V_2.$$

The existence and uniqueness of ρ follow immediately from the existence and uniqueness of the tensor product. We write $\rho_s \equiv \rho_s^1 \otimes \rho_s^2$ as the *tensor product* of the given representations.

Recall that if (e_{i_1}) and (e_{i_2}) be bases of V_1 and V_2 respectively, then $(e_{i_1} \otimes e_{i_2})$ is a basis of $V_1 \otimes V_2$. If $(r_{i_1 j_1}(s))$ and $(r_{i_2 j_2}(s))$ are the matrices of ρ_s^1 and ρ_s^2 respectively satisfying

$$\rho_s^1(e_{j_1}) = \sum_{i_1} r_{i_1 j_1}(s) e_{i_1} \qquad \rho_s^2(e_{j_2}) = \sum_{i_2} r_{i_2 j_2}(s) e_{i_2}$$

then the matrix of ρ_s is $(r_{i_1 j_1}(s) \cdot r_{i_2 j_2}(s))$ satisfying

$$\rho_s(e_{j_1} \otimes e_{j_2}) = \sum_{i_1, i_2} r_{i_1 j_1}(s) \cdot r_{i_2 j_2}(s) \cdot e_{i_1} \otimes e_{i_2}.$$

⟨ TODO: example of tensor product ⟩ Note that the tensor product of two irreducible representations is not in general irreducible **⟨ TODO: example? ⟩**.

We now consider the special case of $V \otimes V$. Let (e_i) be a basis of V and define an automorphism θ of $V \otimes V$ such that $\theta(e_i \otimes e_j) = e_j \otimes e_i$. Then note that $\theta^2 \equiv \text{id}_{V \otimes V}$. We may decompose $V \otimes V$ into the direct sum

$$V \otimes V = \text{Sym}^2(V) \oplus \text{Alt}^2(V).$$

Here, $\text{Sym}^2(V)$ is the set of $z \in V \otimes V$ such that $\theta(z) = z$ and $\text{Alt}^2(V)$ is the set of $z \in V \otimes V$ where $\theta(z) = -z$. These have bases $(e_i \otimes e_j + e_j \otimes e_i)_{i \leq j}$ and $(e_i \otimes e_j - e_j \otimes e_i)_{i < j}$ respectively. As such, $\dim(\text{Sym}^2(V)) = \frac{n(n+1)}{2}$ and $\dim(\text{Alt}^2(V)) = \frac{n(n-1)}{2}$ where $n := \dim(V)$.

Definition 1.16 (Symmetric Square, Alternating Square): These subspaces $\text{Sym}^2(V)$ and $\text{Alt}^2(V)$ of $V \otimes V$ are respectively called the *symmetric square* and *alternating square* of the given representation.

Chapter 2

Character Theory

Definition 2.1 (Character): Let $\rho : G \rightarrow \text{GL}(V)$ be a linear representation of a finite group G in V . Then the character χ_ρ of ρ is the function

$$\chi_\rho(s) := \text{Tr}(R_s) \equiv \text{Tr}(\rho_s).$$

for each $s \in G$.

Proposition 2.2: If χ is the character of a representation ρ of degree n then

1. $\chi(e) = 1$;
2. $\chi(s^{-1}) = \chi(s)^*$, the complex conjugate of $\chi(s)$,
3. $\chi(tst^{-1}) = \chi(s)$.

Proof: The first is immediate since ρ_1 is the identity matrix I and $\text{Tr}(I) = n$. Then recall that we may choose our basis to be orthonormal, and as such ρ_s is a unitary matrix. Thus, each eigenvalue $\lambda_1, \dots, \lambda_n$ has an absolute value equal to 1. Thus

$$\chi(s)^* = \text{Tr}(\rho_s)^* = \sum \lambda_i^* = \sum \lambda_i^{-1} = \text{Tr}(\rho_s)^{-1} = \text{Tr}(\rho_{s^{-1}}) = \chi(s^{-1}).$$

This uses the fact that the conjugate of the sum is the sum of the conjugates as well as the fact that the eigenvalues of R_s^{-1} are the inverses of the eigenvalues of R_s . Finally, letting $u = ts$ and $v = t^{-1}$ allows us to write $\chi(tst^{-1}) = \chi(s)$ as $\chi(uv) = \chi(vu)$ which is immediate since for any complex matrices A, B we have that $\text{Tr}(AB) = \text{Tr}(BA)$. □

Proposition 2.3: Let $\rho^1: G \rightarrow \text{GL}(V_1)$ and $\rho^2: G \rightarrow \text{GL}(V_2)$ be two linear representations with characters χ_1 and χ_2 respectively. Then

1. The character χ of the direct sum representation $V_1 \oplus V_2$ is $\chi_1 + \chi_2$.
2. The character ψ of the tensor product representation $V_1 \otimes V_2$ is $\chi_1 \cdot \chi_2$.

Proof: Let R_s^1, R_s^2 be the matrix forms of ρ_s^1 and ρ_s^2 respectively. Then the matrix form R_s of the representation of $V_1 \oplus V_2$ is given by

$$R_s = \begin{pmatrix} R_s^1 & 0 \\ 0 & R_s^2 \end{pmatrix}$$

and thus $\text{Tr}(R_s) = \text{Tr}(R_s^1) + \text{Tr}(R_s^2)$. Let (e_{i_1}) and (e_{i_2}) be bases for V_1 and V_2 . Then we have that

$$\psi(s) = \sum_{i_1 i_2} r_{i_1 i_1}(s) \cdot r_{i_2 i_2}(s) = \left(\sum_{i_1} r_{i_1 i_1}(s) \right) \cdot \left(\sum_{i_2} r_{i_2 i_2}(s) \right) = \chi_1(s) \cdot \chi_2(s).$$

□

Proposition 2.4: Let $\rho: G \rightarrow \text{GL}(V)$ be a linear representation of G with character χ . Let χ_σ^2 be the character of $\text{Sym}^2(V)$ and χ_α^2 be the character of $\text{Alt}^2(V)$ from Definition 1.16. Then

$$\begin{aligned} \chi_\sigma^2(s) &= \frac{1}{2}(\chi(s)^2 + \chi(s^2)) \\ \chi_\alpha^2(s) &= \frac{1}{2}(\chi(s)^2 - \chi(s^2)) \end{aligned}$$

which directly implies that $\chi_\sigma^2 + \chi_\alpha^2 = \chi$.

Proof: Let $s \in G$ and (e_i) a basis of V consisting solely of eigenvectors for ρ_s . Then $\rho_s(e_i) = \lambda_i e_i$ for some $\lambda_i \in \mathbb{C}$. Thus

$$\chi(s) = \sum \lambda_i \qquad \chi(s^2) = \sum \lambda_i^2.$$

We also have that

$$\begin{aligned} (\rho_s \otimes \rho_s)(e_i \otimes e_j + e_j \otimes e_i) &= \lambda_i \lambda_j (e_i \otimes e_j + e_j \otimes e_i) \\ (\rho_s \otimes \rho_s)(e_i \otimes e_j - e_j \otimes e_i) &= \lambda_i \lambda_j (e_i \otimes e_j - e_j \otimes e_i) \end{aligned}$$

which yields that

$$\chi_\sigma^2(s) = \sum_{i \leq j} \lambda_i \lambda_j = \sum \lambda_i^2 + \sum_{i < j} \lambda_i \lambda_j = \frac{1}{2} \left(\sum \lambda_i \right)^2 + \frac{1}{2} \sum \lambda_i^2 \chi_\alpha^2(s) = \sum_{i < j} \lambda_i \lambda_j = \frac{1}{2} \left(\sum \lambda_i \right)^2 - \frac{1}{2} \sum \lambda_i^2.$$

The proposition then directly follows. Note that the equality $\chi_\sigma^2 + \chi_\alpha^2 = \chi^2$ directly reflects the fact that $V \otimes V = \text{Sym}^2(V) \oplus \text{Alt}^2(V)$. \square

Proposition 2.5 (Schur's Lemma): Let $\rho^1: G \rightarrow \text{GL}(V_1)$ and $\rho^2: G \rightarrow \text{GL}(V_2)$ be two irreducible representations of G . Let $f: V_1 \rightarrow V_2$ be a linear map such that $f \circ \rho_s^1 = \rho_s^2 \circ f$ for all $s \in G$. Then

1. If ρ^1 and ρ^2 are not isomorphic, then $f = 0$
2. If $V_1 = V_2$ and $\rho^1 = \rho^2$ then f is a *homothety*, a scalar multiple of the identity.

Proof: The case of $f = 0$ is trivial, so suppose that $f \neq 0$. Let $W_1 = \ker(f)$ and $W_2 = \text{im}(f)$. Then for $x \in W_1$ we have that $f(\rho_s^1(x)) = \rho_s^2(f(x)) = 0$ which means that $\rho_s^1(x) \in W_1$. Thus W_1 is stable under G and irreducibility of V_1 combined with the assumption that $f \neq 0$ implies that $W_1 = 0$. Similarly, we have that for $f(x) \in W_2$, we have that $\rho_s^2(f(x)) = f(\rho_s^1(x)) \in W_2$, so $\rho_s^2(f(x)) \in W_2$. Thus W_2 is also stable under G meaning that by a similar argument, $W_2 = V_2$. Since $\ker(f) = 0$ and $\text{im}(f) = V_2$, we must have that f is an isomorphism $V_1 \rightarrow V_2$. This proves the first claim.

Now suppose that $V_1 = V_2$, $\rho^1 = \rho^2$, and that λ is some eigenvalue of f . Let $f' = f - \lambda$. Since λ is an eigenvalue, then $\ker(f') \neq 0$. However, we also have that $f' \circ \rho_s^1 = \rho_s^2 \circ f'$. The first part of this proof shows that this implies that $f' = 0$. Thus, $f = \lambda$ and f is a homothety. \square

Corollary 2.6: Let $\rho^1: G \rightarrow \text{GL}(V_1)$ and $\rho^2: G \rightarrow \text{GL}(V_2)$ be two irreducible representations of G . Let $h: V_1 \rightarrow V_2$ and define h^0 such that

$$h^0 = \frac{1}{|G|} \sum_{t \in G} (\rho_t^2)^{-1} \circ h \circ \rho_t^1.$$

Then

1. If ρ^1 and ρ^2 are not isomorphic, then $h^0 = 0$
2. If $V_1 = V_2$ and $\rho^1 = \rho^2$, then h^0 is a homothety of ratio $\frac{1}{n} \text{Tr}(h)$, with $n = \dim(V_1)$.

Proof: First for $s \in G$ we have that

$$(\rho_s^2)^{-1} \circ h^0 \circ \rho_s^1 = \frac{1}{|G|} \sum_{t \in G} (\rho_s^2)^{-1} (\rho_t^2)^{-1} \circ h \circ \rho_t^1 \rho_s^1. \quad = \frac{1}{|G|} \sum_{t \in G} (\rho_{ts}^2)^{-1} \circ h \circ \rho_{ts}^1 = h^0.$$

Thus, Proposition 2.5 applies to h^0 and in the first case $h^0 = 0$ and in the second h^0 is a homothety of scalar λ . Moreover we have that

$$n \cdot \lambda = \text{Tr}(\lambda) = \frac{1}{|G|} \sum_{t \in G} \text{Tr}((\rho_t^1)^{-1} \circ h \circ \rho_t^1) = \text{Tr}(h).$$

Thus, $\lambda = \frac{1}{n} \text{Tr}(h)$. \square

Consider Corollary 2.6 in matrix form where $\rho_s^1 = (r_{i_1 j_1}(s))$ and $\rho_s^2 = (r_{i_2 j_2}(s))$. Then our linear map h is given by the matrix $(x_{i_2 i_1})$ and similarly h^0 is given by the matrix $(x_{i_2 i_1}^0)$. Then by definition of h^0 we have that

$$x_{i_2 i_1}^0 = \frac{1}{|G|} \sum_{t \in G, j_1, j_2} r_{i_2 j_2}(t^{-1}) \cdot x_{j_2 j_1} \cdot r_{j_1 i_1}(t).$$

In case (1) of Corollary 2.6, the right hand side must vanish and so all coefficients of 0:

Corollary 2.7: In case (1) of Corollary 2.6 we have that

$$\frac{1}{|G|} \sum_{t \in G} r_{i_2 j_2}(t^{-1}) \cdot r_{j_1 i_1}(t) = 0$$

for all i_1, j_1, i_2, j_2 .

We now consider case (2) of Corollary 2.6 in the matrix form. We have that $h^0 = \lambda$, with $\lambda = \frac{1}{n} \text{Tr}(h)$, meaning that $x_{i_2 i_1}^0 = \lambda \delta_{i_2 i_1}$. That is, $\lambda = \frac{1}{n} \sum \delta_{i_2 i_1} \cdot x_{i_2 i_1}$. This we have that

$$\frac{1}{|G|} \sum_{t \in G, j_1, j_2} r_{i_2 j_2}(t^{-1}) \cdot x_{j_2 j_1} \cdot r_{j_1 i_1}(t) = \frac{1}{n} \sum_{j_1, j_2} \delta_{i_2 i_1} \cdot \delta_{j_2 j_1} \cdot x_{j_2 j_1}.$$

Equating coefficients of the $x_{j_2 j_1}$ yields the following corollary:

Corollary 2.8: In case (2) of Corollary 2.6 we have that

$$\frac{1}{|G|} \sum_{t \in G} r_{i_2 j_2}(t^{-1}) \cdot r_{j_1 i_1}(t) = \frac{1}{n} \delta_{i_2 i_1} \cdot \delta_{j_2 j_1}.$$

We may generalize these notions to further functions. Let ϕ, ψ be **<< complex valued? >>** functions on G . Define

$$\langle \phi | \psi \rangle := \frac{1}{|G|} \sum_{t \in G} \phi(t^{-1}) \cdot \psi(t) = \frac{1}{|G|} \sum_{t \in G} \phi(t) \cdot \psi(t^{-1}).$$

Then $\langle \phi | \psi \rangle = \langle \psi | \phi \rangle$ and $\langle \phi | \psi \rangle$ is linear in ϕ and in ψ . Thus, Corollary 2.7 and Corollary 2.8 respectively become

$$\langle r_{i_2 j_2}, r_{j_1 i_1} \rangle = 0 \qquad \langle r_{i_2 j_2}, r_{j_1 i_1} \rangle = \frac{1}{n} \delta_{i_2 i_1} \cdot \delta_{j_2 j_1}.$$

If the matrices $(r_{ij}(t))$ are unitary, realized by a suitable choice of basis, then $r_{ij}(t^{-1}) = r_{ji}(t)^*$ and Corollary 2.7 and Corollary 2.8 are just *orthogonality relations*.

Exercises

Exercise 2.1 (Ser77 2.1): Let χ, χ' be the characters of two representations. Prove the following formulas:

$$(\chi + \chi')_{\sigma}^2 = \chi_{\sigma}^2 + \chi'_{\sigma}^2 + \chi \chi'$$

$$(\chi + \chi')_{\alpha}^2 = \chi_{\alpha}^2 + \chi'_{\alpha}^2 + \chi \chi'$$

Proof: Let $s \in G$. Then by Proposition 2.4 we have that

$$\begin{aligned} (\chi + \chi')_{\sigma}^2(s) &= \frac{1}{2}((\chi + \chi')^2(s) + (\chi + \chi')(s^2)) \\ &= \frac{1}{2}(\chi(s)^2 + \chi'(s)^2 + 2\chi(s)\chi'(s) + \chi(s^2) + \chi'(s^2)) \\ &= \frac{1}{2}(\chi(s)^2 + \chi(s^2)) + \frac{1}{2}(\chi'(s) + \chi'(s^2)) + \chi(s)\chi'(s) = \chi_{\sigma}^2(s) + \chi'_{\sigma}^2(s) + \chi(s)\chi'(s). \end{aligned}$$

Since this holds for all $s \in G$, the formula holds in general. The proof of the other formula is similar. \square

Exercise 2.2 (Ser77 2.2): Let X be a finite set on which G acts, and $\rho: G \rightarrow \text{GL}(V)$ the corresponding permutation representation (Example 1.7), and χ_X the character of ρ . Then show that for $s \in G$, $\chi_X(s)$ is equal to the number of elements fixed by s .

Proof: Suppose $X = [n]$ and so $s \in S_n$, meaning $G \leq S_n$. We may assume this without loss of generality. Note that $R_s = (r_{ij}(s))$ where $r_{ij}(s) = 1$ if $s(j) = i$ and 0 otherwise. We want to count the number of elements in $[n]$ fixed by s , i.e. the number of i such that $\sigma(i) = i$. These correspond exactly to the entries in R_s where $r_{ii}(s) = 1$. Thus, the claim follows. \square

Bibliography

- [Ful96] William Fulton. *Young Tableaux: With Applications to Representation Theory and Geometry*. London Mathematical Society Student Texts. Cambridge University Press, 1996. DOI: [10.1017/CB09780511626241](https://doi.org/10.1017/CB09780511626241).
- [Pan23] Greta Panova. *Computational Complexity in Algebraic Combinatorics*. 2023. arXiv: [2306.17511](https://arxiv.org/abs/2306.17511) [math.CO].
- [Rez20] Charles Rezk. *A short course on finite group representations*. 2020. URL: <https://rezk.web.illinois.edu/Finite%5C%20Group%5C%20Reps/short-course-finite-group-representations.html>.
- [Ser77] Jean-Pierre Serre. *Linear Representations of Finite Groups*. Springer New York, 1977. ISBN: 9781468494587. DOI: [10.1007/978-1-4684-9458-7](https://doi.org/10.1007/978-1-4684-9458-7).