

# Algebraic Curves

With 0 Figures

Anakin Dey

Last Edited on 9/2/24 at 14:55

# Contents

<b>1</b>	<b>Affine Algebraic Sets</b>	<b>1</b>
1.1	Algebraic Preliminaries . . . . .	1
1.2	Affine Space and Algebraic Sets . . . . .	4
1.3	The Ideal of a Set of Points . . . . .	4

# Preface

At the time of writing this, I am starting my PhD at The Ohio State University. Currently a large part of my interests in algebra are about algorithms as they relate to polynomials and algebraic geometry. As such, I've been primarily looking at *Ideals, Varieties, and Algorithms* [CLO15], I've been doing a bunch of problems from *Ideals, Varieties, and Algorithms* [CLO15], *Using Algebraic Geometry* [CLO05] and stealing glances at Sturmfel's *Algorithms in Invariant Theory* [Str08]. However, this book is such a classic and I'm of the opinion that you can never know too much about curves and varieties. To put my money where my mouth is, I've started to  $\text{\TeX}$  up some exercises from Fulton's *Algebraic Curves* [Ful08].

# Chapter 1

## Affine Algebraic Sets

### 1.1 Algebraic Preliminaries

**Solution:** [Ful08] Ex. 1.1: As normal, we denote a monomial  $x_1^{i_1} \cdots x_n^{i_n}$  as  $\bar{x}^I$  where  $|I| = i_1 + \cdots + i_n = d \in \mathbb{N}$  and  $I \in \mathbb{N}^n$ . Thus, if  $f(x_1, \dots, x_n), g(x_1, \dots, x_n) \in R[x_1, \dots, x_n]$  are forms of degree  $r$  and  $s$  respectively, we have that

$$\begin{aligned} f(x_1, \dots, x_n) &= \sum_{I \in \mathbb{N}^n, |I|=r} f_I \bar{x}^I, \\ g(x_1, \dots, x_n) &= \sum_{J \in \mathbb{N}^n, |J|=s} g_J \bar{x}^J. \end{aligned}$$

Then, we have by distribution that

$$f(x_1, \dots, x_n) \cdot g(x_1, \dots, x_n) = \sum_{\substack{I \in \mathbb{N}^n, |I|=r \\ J \in \mathbb{N}^n, |J|=s}} f_I \bar{x}^I g_J \bar{x}^J.$$

For each  $I, J$  we have that if  $f_I, g_J \neq 0$  then  $f_I g_J \bar{x}^I \bar{x}^J = (f g)_{I,J} \bar{x}^{I+J}$  where  $(f g)_{I,J} = (f g)_{I,J} = f_I g_J$ . Notice that  $|I + J| = r + s$  by coordinatewise addition and so each term is either 0 or a degree  $r + s$  homogeneous form. Thus after combining like terms, we have that  $f(x_1, \dots, x_n) \cdot g(x_1, \dots, x_n)$  is either degree 0 or a degree  $r + s$  homogeneous form. As 0 has any degree we choose, we have that  $f \cdot g$  is a degree  $r + s$  form.

If we wanted to show that the product of two nonzero polynomials is still nonzero, it can be shown that  $R$  a domain implies  $R[x_1, \dots, x_n]$  is a domain. This can be seen by looking at leading terms and inducting on the number of variables. □

**Solution: [Ful08] Ex. 1.2:** The expression of  $z \in K = \text{Frac}(R)$  as  $z = \frac{a}{b}$  for some  $a \in R, b \in R$  such that  $a, b$  have no common factors, is obvious as we only consider finite sums so we can always take common denominators. Suppose that  $z = \frac{a}{b} = \frac{a'}{b'}$  two different representatives. Since  $R$  is a UFD, say that  $a = u_a a_1 \cdots a_{n_a}$  and  $a' = u_{a'} a'_1 \cdots a'_{n_{a'}}$  are factorizations of  $a$  and  $a'$ . We have similar factorizations for  $b$  and  $b'$ . Then we have that  $ab' = a'b \in R$ . However, as  $R$  is a unique factorization domain, we must have

$$u_a a_1 \cdots a_{n_a} u_{b'} b'_1 \cdots b'_{n_{b'}} = u_{a'} a'_1 \cdots a'_{n_{a'}} u_b b_1 \cdots b_{n_b}$$

are two equivalent factorizations up to unit. This implies that  $\frac{a}{b}$  and  $\frac{a'}{b'}$  differ only up to unit.  $\square$

**Solution: [Ful08] Ex. 1.3:**

- (a) We know that  $P = (p)$  for some nonzero, nonunit  $p \in R$ . Suppose that  $p$  was reducible, so  $p = ab$  for some nonzero nonunit  $a, b \in R$ . Then as  $p \nmid a$  and  $p \nmid b$ , since such divisibility would imply that  $a$  or  $b$  were equal to  $p$  up to unit, we have that  $a, b \notin (p)$ . But then we have elements  $a, b \in R$  such that  $a \notin (p)$ ,  $b \notin (p)$  but  $ab = p \in (p)$  which contradicts the primality of  $P = (p)$ . Then  $p$  is irreducible.
- (b) We show that  $P = (p)$  is maximal. There exists a maximal ideal  $M = (m)$  such that  $(p) \subseteq (m)$ . Thus,  $p \mid m$ . Notice that by (a) that both  $p$  and  $m$  are irreducible elements of  $R$ . As such, if  $p \mid m$  we must have that  $m \mid p$  as if not,  $m$  would be reducible. Thus,  $p = um$  for some unit  $u \in R^\times$ . Thus,  $(p) = (m)$  and  $P = (p)$  is maximal.

$\square$

**Solution: [Ful08] Ex. 1.4:** We prove by induction on the number of variables  $n$ . If  $n = 1$  then any nonzero polynomial of degree  $d$  has at most  $d$  distinct zeros. Thus if  $F(x) \in k[x]$  is a nonzero polynomial, there exists a  $z \in k$  such that  $F(z) \neq 0$ . Now let  $n \geq 1$ . Let  $F(x_1, \dots, x_n, y) \in k[x_1, \dots, x_n, y]$  be a nonzero polynomial. Letting  $d = \deg_y(F)$ , we can find  $F_0[x_1, \dots, x_n], \dots, F_d[x_1, \dots, x_n]$  such that

$$F(x_1, \dots, x_n, y) = \sum_{i=0}^d F_i(x_1, \dots, x_n) \cdot y^i.$$

Regard  $F(x_1, \dots, x_n, y)$  as a nonzero polynomial  $g(y)$  in  $F(x_1, \dots, x_n)[y]$ . As  $k$  is infinite and  $g(y)$  is nonzero, then we must have there exists  $z_y \in k$  such that  $g(z_y) \neq 0$ . Thus, we have that  $F(x_1, \dots, x_n, z_y) \in k[x_1, \dots, x_n]$  is nonzero. By induction, there exists  $(z_1, \dots, z_n) \in k^n$  such that  $F(z_1, \dots, z_n, z_y) \neq 0$ .  $\square$

**Solution: [Ful08] Ex. 1.5:** This is Euclid's proof of there being infinitely many primes. Omitted.  $\square$

**Solution: [Ful08] Ex. 1.6:** Let  $k$  be an algebraically closed field. Then the irreducible monic polynomials of  $k[x]$  are of the form  $(x - a)$  for each  $a \in k$ . Thus, they are in bijection with the elements of  $k$ . By [Ful08] Ex. 1.5, there are infinitely many monic irreducible polynomials over any field. Thus,  $k$  is an infinite field.  $\square$

**Solution: [Ful08] Ex. 1.7:**

(a) Let  $F(x_1, \dots, x_n) \in k[x_1, \dots, x_n]$ . Then we may express  $F(x_1, \dots, x_n)$  as

$$F(x_1, \dots, x_n) = \sum_{I=(i_1, \dots, i_n) \in \mathbb{N}^n} \lambda_I (x_1)^{i_1} \cdots (x_n)^{i_n} = \sum_{I=(i_1, \dots, i_n) \in \mathbb{N}^n} \lambda_I ((x_1 - a_1) + a_1)^{i_1} \cdots ((x_n - a_n) + a_n)^{i_n}.$$

Expanding and combining like terms yields the desired results.

(b) Suppose  $F(a_1, \dots, a_n) = 0$ . By (a), we may express  $F(x_1, \dots, x_n)$  as

$$F(x_1, \dots, x_n) = \sum_{I=(i_1, \dots, i_n) \in \mathbb{N}^n} \lambda_I (x_1 - a_1)^{i_1} \cdots (x_n - a_n)^{i_n}$$

The claim says that this expression has no constant term, i.e.  $\lambda_{\vec{0}} = 0$ . This is immediate as the constant term corresponds to  $F(a_1, \dots, a_n) = 0$ .  $\square$

## 1.2 Affine Space and Algebraic Sets

**Solution:** [Ful08] Ex. 1.8: This corresponds to the fact that any nonzero univariate polynomial has finitely many zeroes and the zero polynomial vanishes everywhere by definition.  $\square$

**Solution:** [Ful08] Ex. 1.14:  $\square$

**Solution:** [Ful08] Ex. 1.15: Let  $V = \mathbf{V}(I)$  for some ideal  $I \subseteq k[\bar{x}]$  and  $W = \mathbf{V}(J)$  for some ideal  $J \subseteq k[\bar{y}]$ . We may consider  $I, J \subseteq k[\bar{x}, \bar{y}]$  as well by identifying  $I$  with  $\langle I \rangle$  where we generated the ideal over  $k[\bar{x}, \bar{y}]$  and do similarly for  $J$ . **⟨ It is clear ⟩** that  $V \times W = \langle I \cup J \rangle \subseteq k[\bar{x}, \bar{y}]$ .  $\square$

## 1.3 The Ideal of a Set of Points

**Solution:** [Ful08] Ex. 1.16: This follows immediately from the fact that  $V \subseteq W \implies \mathbf{I}(V) \supseteq \mathbf{I}(W)$ .  $\square$

# Bibliography

- [CLO05] D.A. Cox, J. Little, and D. O'Shea. *Using Algebraic Geometry*. Springer-Verlag, 2005. ISBN: 0387207066. DOI: [10.1007/b138611](https://doi.org/10.1007/b138611). URL: <http://dx.doi.org/10.1007/b138611>.
- [CLO15] D.A. Cox, J. Little, and D. O'Shea. *Ideals, Varieties, and Algorithms: An Introduction to Computational Algebraic Geometry and Commutative Algebra*. Undergraduate Texts in Mathematics. Springer International Publishing, 2015. ISBN: 9783319167213. URL: <https://books.google.com/books?id=yL7yCAAQBAJ>.
- [Ful08] W. Fulton. *Algebraic Curves: An Introduction to Algebraic Geometry*. 3rd ed. 2008.
- [Str08] Bernd Strumfels. *Algorithms in Invariant Theory*. Springer Vienna, 2008. ISBN: 9783211774175. DOI: [10.1007/978-3-211-77417-5](https://doi.org/10.1007/978-3-211-77417-5). URL: <http://dx.doi.org/10.1007/978-3-211-77417-5>.