



Representation Theory Notes and Exercises

With 0 Figures

Anakin Dey

Last Edited on 12/16/23 at 22:47

TODOs

⟨⟨ Change the style of enumerates from “1.” to “(1)” ⟩⟩

⟨⟨ Proper Exercise Header ⟩⟩

⟨⟨ Proper Chapter Header ⟩⟩

Contents

1	Generalities on Linear Representations	1
----------	---	----------

List of Definitions

1.1	Linear Representation, Representation Space	1
1.2	Degree	1
1.3	Matrix of a Representation	1
1.4	Similar/Isomorphic Representations	2

List of Examples and Counterexamples

1.5	Unit/Trivial Representation	2
1.6	Regular Representation	2
1.7	Permutation Representation	2

Preface

This is a set of notes on group representation theory mainly based on J.P. Serre's text *Linear Representations of Finite Groups* [Ser77]. Occasionally, other sources may be used. The goal of these notes is to eventually work towards algebraic combinatorics such as Fulton's text *Young Tableaux* [Ful96], as much of algebraic combinatorics is motivated by questions stemming from representation theory. At the time of writing this, another goal is the applications of representation theory to computational complexity: see [Pan23] for a recent survey on this connection.

Chapter 1

Generalities on Linear Representations

Unless otherwise specified, V will denote a vector space, usually over the field \mathbb{C} . We will restrict ourselves to finite dimensional vector spaces. Similarly, we will restrict ourselves to finite groups.

Definition 1.1 (Linear Representation, Representation Space): Let G be a group with identity e . A *linear representation* of G in V is a homomorphism $\rho : G \rightarrow \text{GL}(V)$. We will frequently, and often interchangeably, write $\rho_s := \rho(s)$. Given ρ , we will say that V is a *representation space* or *representation* of G .

Definition 1.2 (Degree): Let $\rho : G \rightarrow V$ be a representation of G in a vector space V . Then the *degree* of ρ is $\dim(V)$.

Let $\rho : G \rightarrow V$ be a representation of G in a vector space V with $n := \dim(V)$. Fix a basis (e_j) of V . Then since each ρ_s is an invertible linear transformation of V , we may define an $n \times n$ matrix $R_s \equiv (r_{ij}(s))$ where each $r_{ij}(s)$ is defined by the identity

$$\rho_s(e_j) = \sum_{i=1}^n r_{ij}(s) e_i.$$

Definition 1.3 (Matrix of a Representation): We call $R_s = (r_{ij}(s))$ above the *matrix of ρ_s* with respect to the basis (e_j) .

Note that R_s satisfies the following:

$$\det(R_s) \neq 0, \quad R_{st} = R_s \cdot R_t \equiv r_{ij}(st) = \sum_{k=1}^n r_{ik}(s) \cdot r_{kj}(t) \quad \forall s, t \in G.$$

Recall that two $n \times n$ matrices A, A' are *similar* if there exists an invertible matrix T such that $TA = A'T$. We may extend this notion to representations.

Definition 1.4 (Similar/Isomorphic Representations): Let ρ and ρ' be two representations of the same group G in vector spaces V and V' respectively. We say ρ and ρ' are *similar* or *isomorphic* if there exists an isomorphism $\tau: V \rightarrow V'$ such that for all $s \in G$, τ satisfies $\tau \circ \rho(s) = \rho'(s) \circ \tau$. If R_s, R'_s are the corresponding matrices then this is equivalent to saying there exists an invertible matrix T such that $TR_s = R'_sT$ for all $s \in G$.

Note that if ρ and ρ' are isomorphic, then they must have the same degree.

We now give some examples of these things.

Example 1.5 (Unit/Trivial Representation): Let G be a finite group. Representations of degree 1 must be of the form $\rho: G \rightarrow \mathbb{C}^\times$. Since elements s of G are of finite order, $\rho(s)$ must also be of finite order. Thus, for all $s \in G$, $\rho(s)$ is a root of unity. If we take $\rho(s) = 1$ for all $s \in G$, we obtain the *unit* or *trivial* representation of G .

Example 1.6 (Regular Representation): Let g be the order of G , and let V be a vector space of dimension g with a basis $(e_t)_{t \in G}$. For each $s \in G$, define ρ_s as the linear map $\rho_s: V \rightarrow V$ sending $e_t \mapsto e_{st}$. This is a linear representation of G called the *regular* representation of G . Since for each $s \in G$, $e_s = \rho_s(e_1)$ and thus the images of e_1 form a basis of V . Conversely, let W be a representation of G with a vector w satisfying the collection of all $\rho_s(w)$, $s \in G$, forms a basis of W . Then W is isomorphic to the regular representation of G by the isomorphism $\tau(e_s) = \rho_s(w)$.

Example 1.7 (Permutation Representation): We may generalize the regular representation to any group action $G \curvearrowright X$, X a finite set. Recall that for such an action, the map $x \mapsto sx$ for each $s \in G$ is a permutation $X \leftrightarrow X$. Let V be a vector space with dimension the size of X , and so a basis $(e_x)_{x \in X}$. Define a representation ρ of G by defining ρ_s as the linear map sending $e_x \mapsto e_{sx}$. This representation is known as the *Permutation* representation of G associated with X .

Bibliography

- [Ful96] William Fulton. *Young Tableaux: With Applications to Representation Theory and Geometry*. London Mathematical Society Student Texts. Cambridge University Press, 1996. DOI: [10.1017/CB09780511626241](https://doi.org/10.1017/CB09780511626241).
- [Pan23] Greta Panova. *Computational Complexity in Algebraic Combinatorics*. 2023. arXiv: [2306.17511](https://arxiv.org/abs/2306.17511) [math.CO].
- [Ser77] Jean-Pierre Serre. *Linear Representations of Finite Groups*. Springer New York, 1977. ISBN: 9781468494587. DOI: [10.1007/978-1-4684-9458-7](https://doi.org/10.1007/978-1-4684-9458-7).