



# Representation Theory Notes and Exercises

With 0 Figures

Anakin Dey

Last Edited on 12/21/23 at 00:19

## TODOs

⟨⟨ Change the style of enumerates from “1.” to “(1)” ⟩⟩

⟨⟨ Proper Exercise Header ⟩⟩

⟨⟨ Proper Chapter Header ⟩⟩

⟨⟨ Remove indent at end of thmtool environments ⟩⟩

# Contents

1	Generalities on Linear Representations	1
2	Character Theory	7

# List of Definitions

1.1	Linear Representation, Representation Space . . . . .	1
1.2	Degree . . . . .	1
1.3	Matrix of a Representation . . . . .	1
1.4	Similar/Isomorphic Representations . . . . .	2
1.8	Stable/Invariant Subspaces, Subrepresentation . . . . .	3
1.11	Direct Sum of Representations . . . . .	4
1.12	Irreducible/Simple Representations . . . . .	4
1.15	Tensor/Kronecker Product of Representations . . . . .	5
1.16	Symmetric Square, Alternating Square . . . . .	6
2.1	Character . . . . .	7
2.9	Scalar Product . . . . .	11
2.18	Class Function . . . . .	13
2.24	Canonical Decomposition of a Representation . . . . .	15

# List of Examples and Counterexamples

1.5	Unit/Trivial Representation . . . . .	2
1.6	Regular Representation . . . . .	2
1.7	Permutation Representation . . . . .	3
1.9	Subrepresentations of the Regular Representation . . . . .	3
1.14	Decomposition of Representation of $\mathbb{Z}_3$ into Irreducibles . . . . .	5
2.23	Character Table of $S_3$ . . . . .	15
2.26	Decomposition of $C_2$ . . . . .	16

# Preface

This is a set of notes on group representation theory mainly based on J.P. Serre's text *Linear Representations of Finite Groups* [Ser77]. Occasionally, other sources may be used, such as the set of notes by Charles Rezk created for Math 427 [Rez20]. The goal of these notes is to eventually work towards algebraic combinatorics such as Fulton's text *Young Tableaux* [Ful96], as much of algebraic combinatorics is motivated by questions stemming from representation theory. At the time of writing this, another goal is the applications of representation theory to computational complexity: see [Pan23] for a recent survey on this connection.

# Chapter 1

## Generalities on Linear Representations

Unless otherwise specified,  $V$  will denote a vector space, usually over the field  $\mathbb{C}$ . We will restrict ourselves to finite dimensional vector spaces. Similarly, we will restrict ourselves to finite groups.

**Definition 1.1 (Linear Representation, Representation Space):** Let  $G$  be a group with identity  $e$ . A *linear representation* of  $G$  in  $V$  is a homomorphism  $\rho : G \rightarrow \text{GL}(V)$ . We will frequently, and often interchangeably, write  $\rho_s := \rho(s)$ . Given  $\rho$ , we will say that  $V$  is a *representation space* or *representation* of  $G$ .

**Definition 1.2 (Degree):** Let  $\rho : G \rightarrow V$  be a representation of  $G$  in a vector space  $V$ . Then the *degree* of  $\rho$  is  $\dim(V)$ .

Let  $\rho : G \rightarrow V$  be a representation of  $G$  in a vector space  $V$  with  $n := \dim(V)$ . Fix a basis  $(e_j)$  of  $V$ . Then since each  $\rho_s$  is an invertible linear transformation of  $V$ , we may define an  $n \times n$  matrix  $R_s \equiv (r_{ij}(s))$  where each  $r_{ij}(s)$  is defined by the identity

$$\rho_s(e_j) = \sum_{i=1}^n r_{ij}(s) e_i.$$

**Definition 1.3 (Matrix of a Representation):** We call  $R_s = (r_{ij}(s))$  above the *matrix of  $\rho_s$*  with respect to the basis  $(e_j)$ .

Note that  $R_s$  satisfies the following:

$$\det(R_s) \neq 0, \quad R_{st} = R_s \cdot R_t \equiv r_{ij}(st) = \sum_{k=1}^n r_{ik}(s) \cdot r_{kj}(t) \quad \forall s, t \in G.$$

Recall that two  $n \times n$  matrices  $A, A'$  are *similar* if there exists an invertible matrix  $T$  such that  $TA = A'T$ . We may extend this notion to representations.

**Definition 1.4 (Similar/Isomorphic Representations):** Let  $\rho$  and  $\rho'$  be two representations of the same group  $G$  in vector spaces  $V$  and  $V'$  respectively. We say  $\rho$  and  $\rho'$  are *similar* or *isomorphic* if there exists an isomorphism  $\tau: V \rightarrow V'$  such that for all  $s \in G$ ,  $\tau$  satisfies  $\tau \circ \rho(s) = \rho'(s) \circ \tau$ . If  $R_s, R'_s$  are the corresponding matrices then this is equivalent to saying there exists an invertible matrix  $T$  such that  $TR_s = R'_s T$  for all  $s \in G$ .

Note that if  $\rho$  and  $\rho'$  are isomorphic, then they must have the same degree.

We now give some examples of these things.

**Example 1.5 (Unit/Trivial Representation):** Let  $G$  be a finite group. Representations of degree 1 must be of the form  $\rho: G \rightarrow \mathbb{C}^\times$ . Since elements  $s$  of  $G$  are of finite order,  $\rho(s)$  must also be of finite order. Thus, for all  $s \in G$ ,  $\rho(s)$  is a root of unity. If we take  $\rho(s) = 1$  for all  $s \in G$ , we obtain the *unit* or *trivial* representation of  $G$ . This also means that  $R_s = 1$  for all  $s$ .

**Example 1.6 (Regular Representation):** Let  $g$  be the order of  $G$ , and let  $V$  be a vector space of dimension  $g$  with a basis  $(e_t)_{t \in G}$ . For each  $s \in G$ , define  $\rho_s$  as the linear map  $\rho_s: V \rightarrow V$  such that  $\rho_s(e_t) = e_{st}$ . This is a linear representation of  $G$  called the *regular* representation of  $G$ . Since for each  $s \in G$ ,  $e_s = \rho_s(e_1)$  and thus the images of  $e_1$  form a basis of  $V$ . Conversely, let  $W$  be a representation of  $G$  with a vector  $w$  satisfying the collection of all  $\rho_s(w)$ ,  $s \in G$ , forms a basis of  $W$ . Then  $W$  is isomorphic to the regular representation of  $G$  by the isomorphism  $\tau(e_s) = \rho_s(w)$ .

For example, let  $G = \mathbb{Z}_3$  and  $V = \mathbb{C}^3$  with  $e_0 = (1, 0, 0)$ ,  $e_1 = (0, 1, 0)$ , and  $e_2 = (0, 0, 1)$ . Then for example,  $\rho_0, \rho_1, \rho_2: \mathbb{C}^3 \rightarrow \mathbb{C}^3$  are the linear maps such that

$$\begin{array}{lll} \rho_0(e_0) = e_{0+0} = e_0 & \rho_0(e_1) = e_{0+1} = e_1 & \rho_0(e_2) = e_{0+2} = e_2 \\ \rho_1(e_0) = e_{1+0} = e_1 & \rho_1(e_1) = e_{1+1} = e_2 & \rho_1(e_2) = e_{1+2} = e_0 \\ \rho_2(e_0) = e_{2+0} = e_2 & \rho_2(e_1) = e_{2+1} = e_0 & \rho_2(e_2) = e_{2+2} = e_1 \end{array}$$

With this, the matrix representations of  $\rho_0, \rho_1$  and  $\rho_2$  is similarly straightforward:

$$R_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad R_1 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad R_2 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$



**Example 1.7 (Permutation Representation):** We may generalize the regular representation to any group action  $G \curvearrowright X$ ,  $X$  a finite set. Recall that for such an action, the map  $x \mapsto sx$  for each  $s \in G$  is a permutation  $X \leftrightarrow X$ . Let  $V$  be a vector space with dimension the size of  $X$ , and so a basis  $(e_x)_{x \in X}$ . Define a representation  $\rho$  of  $G$  by defining  $\rho_s$  as the linear map sending  $e_x \mapsto e_{sx}$ . This representation is known as the *Permutation* representation of  $G$  associated with  $X$ . If we consider  $X = [n]$  and  $G = S_n$ , then take  $V = \mathbb{C}^n$  as our vector space and  $e_i$  as the standard basis vector. Then  $\rho_\sigma(e_j) = e_{\sigma(j)}$ . Thus for each  $\sigma \in S_n$ , we have that  $R_\sigma = (r_{ij}(\sigma))$  where entry  $r_{ij}(\sigma) = 1$  if  $i = \sigma(j)$  and 0 otherwise.

**Definition 1.8 (Stable/Invariant Subspaces, Subrepresentation):** Let  $\rho: G \rightarrow \text{GL}(V)$  be a linear representation and  $W \subseteq V$  a subspace of  $V$ . We say that  $W$  is *stable* under the action of  $G$  if  $x \in W$  implies that  $\rho_s(x) \in W$  for all  $s \in G$ . Thus, the restriction  $\rho_s^W := \rho_s|_W$  is an isomorphism of  $W$  onto itself. Restrictions satisfy the property that  $\rho_s^W \circ \rho_t^W = \rho_{st}^W$ . Thus,  $\rho^W: G \rightarrow \text{GL}(W)$  is a linear representation of  $G$  in  $W$  and we say that  $W$  is a *subrepresentation* of  $V$ .

**Example 1.9 (Subrepresentations of the Regular Representation):** Let  $G$  be a group. Recall the regular representation  $V$  given in Example 1.6. Let  $W$  be the 1 dimensional subspace of  $V$  generated by the element  $x = \sum_{s \in G} e_s$ . Then note that  $\rho_s(x) = x$  for all  $s \in G$  and thus  $W$  is a subrepresentation of  $V$ . Furthermore, this is isomorphic to the unit representation Example 1.5 with  $\tau: C^\times \rightarrow W$  such that  $\tau(1) = x$ . For example, let  $G = \mathbb{Z}_3$  and  $\rho: \mathbb{Z}_3 \rightarrow \text{GL}(\mathbb{C}^3)$  the representation given in Example 1.6. Then  $x = (1, 1, 1)$  and for example we have that

$$\rho_1(x) = \rho_1(e_0) + \rho_1(e_1) + \rho_1(e_2) = e_1 + e_2 + e_0 = x.$$

**Theorem 1.10:** Let  $\rho: G \rightarrow \text{GL}(V)$  be a linear representation of  $G$  in  $V$  and let  $W$  be a subspace of  $V$  stable under  $G$ . Then there exists a complement  $W^0$  of  $W$  in  $V$  which is stable under  $G$ .

**Proof:** Let  $W'$  be an arbitrary complement of  $W$  in  $V$ , and let  $p: V \rightarrow W$  be the projection. Then we form the average  $p^0$  of conjugates of  $p$  by elements in  $G$ :

$$p^0 := \frac{1}{|G|} \sum_{t \in G} \rho_t \circ p \circ \rho_t^{-1}.$$

Since  $p: V \rightarrow W$  and  $\rho_t$  preserves  $W$ , we have that  $p^0$  maps  $V$  onto  $W$ . Furthermore, note that  $\rho_t^{-1}$  also preserves  $W$ .

Thus we have that

$$(p \circ \rho_t^{-1})(x) = \rho_t^{-1}(x), \quad (\rho_t \circ p \circ \rho_t^{-1})(x) = x, \quad p^0(x) = x.$$

Thus,  $p^0$  is a projection of  $V$  onto  $W$ , corresponding to some complement  $W^0$  of  $W$ . Moreover, we have that  $\rho_s \circ p^0 = p^0 \circ \rho_s$  for all  $s \in G$  because

$$\rho_s \circ p^0 \circ \rho_s^{-1} = \frac{1}{|G|} \sum_{t \in G} \rho_s \circ \rho_t \circ p \circ \rho_t^{-1} \circ \rho_s^{-1} = \frac{1}{|G|} \sum_{t \in G} \rho_{st} \circ p \rho_{st}^{-1} = p^0.$$

Now suppose that  $x \in W^0$  and  $s \in G$ , we have that  $p^0(x) = 0$  and hence  $(p^0 \circ \rho_s)(x) = (\rho_s \circ p^0)(x) = 0$ , meaning that  $\rho_s(x) \in W^0$ . This,  $W^0$  is stable under  $G$ .  $\square$

Suppose that  $V$  had an inner product  $\langle x, y \rangle$ , and furthermore suppose this inner product was invariant under  $G$  meaning that for all  $s \in G$ ,  $\langle \rho_s(x), \rho_s(y) \rangle = \langle x, y \rangle$ . We may also reduce to this case by replacing  $\langle x, y \rangle$  with  $\sum_{t \in G} \langle \rho_t(x), \rho_t(y) \rangle$ . With this, the orthogonal complement  $W^\perp$  of  $W$  in  $V$  is a complement of  $W$  stable under  $G$ . Note that the invariance of  $\langle x, y \rangle$  means that if  $(e_i)$  is an orthonormal basis of  $V$ , then  $R_s$  is a unitary matrix.

Using the notation of Theorem 1.10, let  $x \in V$  and  $w, w^0$  be the projections of  $x$  on  $W$  and  $W^0$  respectively. Thus for all  $s \in G$ ,  $\rho_s(x) = \rho_s(w) + \rho_s(w^0)$ . Since  $W$  and  $W^0$  are stable under  $G$ , we have that  $\rho_s(w) \in W$  and  $\rho_s(w^0) \in W^0$ . This means that  $\rho_s(w)$  and  $\rho_s(w^0)$  are the projections of  $\rho_s(x)$  and in turn the representations of  $W$  and  $W^0$  determine the representations of  $V$ .

**Definition 1.11 (Direct Sum of Representations):** Given the above, we write  $V = W \oplus W^0$  as the *direct sum* of  $W$  and  $W^0$ . We identify elements  $v \in V$  as pairs  $(w, w^0)$  given by their projections.

If the representations  $W$  and  $W^0$  are given in matrices  $R_s$  and  $R_s^0$ , then the matrix form of the representation  $V$  is given by

$$\begin{pmatrix} R_s & 0 \\ 0 & R_s^0 \end{pmatrix}.$$

Similar results hold for arbitrarily many, but finite, direct sums of representations.

The above is a discussion on how to compose two or more representations into one larger representation. The natural question is then about the opposite.

**Definition 1.12 (Irreducible/Simple Representations):** Let  $\rho : G \rightarrow \text{GL}(V)$  be a linear representation of  $G$ . Then this representation is *irreducible* or *simple* if  $V$  has no subspaces stable under  $G$  besides 0 and  $V$  itself.

By Theorem 1.10, this is equivalent to saying that  $V$  is not the direct sum of two representations besides  $V = 0 \oplus V$ . A representation of degree 1 is reducible. We may use irreducible representations to construct other ones via the direct sum.

**Theorem 1.13:** Every representation is a direct sum of irreducible representations.

**Proof:** Let  $V$  be a linear representation of  $G$ . We induct on  $\dim(V)$ . If  $\dim(V) = 0$ , then  $V = 0$  which is the direct sum of an empty family of irreducible representations. So suppose that

$\dim(V) \geq 1$ . If  $V$  is irreducible, then we are done. Otherwise, there exists a subspace  $W \subsetneq V$  stable under  $G$  and by Theorem 1.10 a stable complement  $W^0$  such that  $V = W \oplus W^0$ . By assumption,  $W \neq 0 \neq W^0$  and so  $\dim(W) < \dim(V)$  and  $\dim(W^0) < \dim(V)$ . By induction, we have obtained a decomposition of  $V$  into irreducibles.  $\square$

**Example 1.14 (Decomposition of Representation of  $\mathbb{Z}_3$  into Irreducibles):** Recall from Example 1.6 the regular representation  $\rho : \mathbb{Z}_3 \rightarrow \text{GL}(\mathbb{C}^3)$  with  $e_0 = (1, 0, 0)$ ,  $e_1 = (0, 1, 0)$ , and  $e_2 = (0, 0, 1)$  and

$$\begin{array}{lll} \rho_0(e_0) = e_0 & \rho_0(e_1) = e_1 & \rho_0(e_2) = e_2 \\ \rho_1(e_0) = e_1 & \rho_1(e_1) = e_2 & \rho_1(e_2) = e_0 \\ \rho_2(e_0) = e_2 & \rho_2(e_1) = e_0 & \rho_2(e_2) = e_1 \end{array}$$

Our goal will be to decompose  $\rho$  into  $\rho^1 \oplus \rho^2 \oplus \rho^3$ . We aim to find the elements fixed by  $\mathbb{Z}_3$ . Note that if an element is fixed by 1, the generator of  $\mathbb{Z}_3$ , then it is fixed by all of  $\mathbb{Z}_3$ . We want to find 1-dimensional  $\mathbb{Z}_3$ -invariant subspaces of  $\mathbb{C}^3$ . This is equivalent to finding the eigenvalues of the matrix

$$R_1 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

The eigenvalues and their eigenvectors of  $R_1$  are as follows, note the inclusion of the trivial subrepresentation from Example 1.9:

$$\lambda_1 = 1, v_1 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad \lambda_2 = \frac{-1 - i\sqrt{3}}{2}, v_2 = \begin{pmatrix} \frac{-1+i\sqrt{3}}{2} \\ \frac{-1-i\sqrt{3}}{2} \\ 1 \end{pmatrix} \quad \lambda_3 = \frac{-1 + i\sqrt{3}}{2}, v_3 = \begin{pmatrix} \frac{-1-i\sqrt{3}}{2} \\ \frac{-1+i\sqrt{3}}{2} \\ 1 \end{pmatrix}$$

Thus  $\mathbb{C}^3 = V_1 \oplus V_2 \oplus V_3$  where  $V_i := \text{span}(v_i)$ . Note that there are only 3 morphisms  $\mathbb{Z}_3 \rightarrow \mathbb{C}^\times$  mapping 1 to 1,  $\omega$ , or  $\omega^2$  where  $\omega$  is a cube root of unity. Thus  $\rho^1, \rho^2$ , and  $\rho^3$  must correspond to these morphisms **<< but which ones >>**.

A natural question is if such a decomposition  $V = W_1 \oplus \cdots \oplus W_k$  is unique. However, suppose that  $\rho$  is the trivial representation (Example 1.5). Then each component of the decomposition is a line and we can decompose a vector space into the direct sum of lines in a number of ways. However, we will see that the number of  $W_i$  that are isomorphic to a given irreducible representation does not depend on the choice of decomposition.

**Definition 1.15 (Tensor/Kronecker Product of Representations):** Let  $\rho^1: G \rightarrow \text{GL}(V_1)$  and  $\rho^2: G \rightarrow \text{GL}(V_2)$  be two representations of a group  $G$ . We construct a representation  $\rho: G \rightarrow \text{GL}(V_1 \otimes V_2)$  such that

$$\rho_s(x_1 \otimes x_2) = \rho_s^1(x_1) \circ \rho_s^2(x_2) \quad \text{for } x_1 \in V_1, x_2 \in V_2.$$

The existence and uniqueness of  $\rho$  follow immediately from the existence and uniqueness of the tensor product. We write  $\rho_s \equiv \rho_s^1 \otimes \rho_s^2$  as the *tensor product* of the given representations.

Recall that if  $(e_{i_1})$  and  $(e_{i_2})$  be bases of  $V_1$  and  $V_2$  respectively, then  $(e_{i_1} \otimes e_{i_2})$  is a basis of  $V_1 \otimes V_2$ . If  $(r_{i_1 j_1}(s))$  and  $(r_{i_2 j_2}(s))$  are the matrices of  $\rho_s^1$  and  $\rho_s^2$  respectively satisfying

$$\rho_s^1(e_{j_1}) = \sum_{i_1} r_{i_1 j_1}(s) e_{i_1} \qquad \rho_s^2(e_{j_2}) = \sum_{i_2} r_{i_2 j_2}(s) e_{i_2}$$

then the matrix of  $\rho_s$  is  $(r_{i_1 j_1}(s) \cdot r_{i_2 j_2}(s))$  satisfying

$$\rho_s(e_{j_1} \otimes e_{j_2}) = \sum_{i_1, i_2} r_{i_1 j_1}(s) \cdot r_{i_2 j_2}(s) \cdot e_{i_1} \otimes e_{i_2}.$$

**⟨ TODO: example of tensor product ⟩** Note that the tensor product of two irreducible representations is not in general irreducible **⟨ TODO: example? ⟩**.

We now consider the special case of  $V \otimes V$ . Let  $(e_i)$  be a basis of  $V$  and define an automorphism  $\theta$  of  $V \otimes V$  such that  $\theta(e_i \otimes e_j) = e_j \otimes e_i$ . Then note that  $\theta^2 \equiv \text{id}_{V \otimes V}$ . We may decompose  $V \otimes V$  into the direct sum

$$V \otimes V = \text{Sym}^2(V) \oplus \text{Alt}^2(V).$$

Here,  $\text{Sym}^2(V)$  is the set of  $z \in V \otimes V$  such that  $\theta(z) = z$  and  $\text{Alt}^2(V)$  is the set of  $z \in V \otimes V$  where  $\theta(z) = -z$ . These have bases  $(e_i \otimes e_j + e_j \otimes e_i)_{i \leq j}$  and  $(e_i \otimes e_j - e_j \otimes e_i)_{i < j}$  respectively. As such,  $\dim(\text{Sym}^2(V)) = \frac{n(n+1)}{2}$  and  $\dim(\text{Alt}^2(V)) = \frac{n(n-1)}{2}$  where  $n := \dim(V)$ .

**Definition 1.16 (Symmetric Square, Alternating Square):** These subspaces  $\text{Sym}^2(V)$  and  $\text{Alt}^2(V)$  of  $V \otimes V$  are respectively called the *symmetric square* and *alternating square* of the given representation.

## Chapter 2

# Character Theory

**Definition 2.1 (Character):** Let  $\rho : G \rightarrow \text{GL}(V)$  be a linear representation of a finite group  $G$  in  $V$ . Then the character  $\chi_\rho$  of  $\rho$  is the function

$$\chi_\rho(s) := \text{Tr}(R_s) \equiv \text{Tr}(\rho_s).$$

for each  $s \in G$ .

**Proposition 2.2:** If  $\chi$  is the character of a representation  $\rho$  of degree  $n$  then

1.  $\chi(e) = 1$ ;
2.  $\chi(s^{-1}) = \chi(s)^*$ , the complex conjugate of  $\chi(s)$ ,
3.  $\chi(tst^{-1}) = \chi(s)$ .

**Proof:** The first is immediate since  $\rho_1$  is the identity matrix  $I$  and  $\text{Tr}(I) = n$ . Then recall that we may choose our basis to be orthonormal, and as such  $\rho_s$  is a unitary matrix. Thus, each eigenvalue  $\lambda_1, \dots, \lambda_n$  has an absolute value equal to 1. Thus

$$\chi(s)^* = \text{Tr}(\rho_s)^* = \sum \lambda_i^* = \sum \lambda_i^{-1} = \text{Tr}(\rho_s)^{-1} = \text{Tr}(\rho_{s^{-1}}) = \chi(s^{-1}).$$

This uses the fact that the conjugate of the sum is the sum of the conjugates as well as the fact that the eigenvalues of  $R_s^{-1}$  are the inverses of the eigenvalues of  $R_s$ . Finally, letting  $u = ts$  and  $v = t^{-1}$  allows us to write  $\chi(tst^{-1}) = \chi(s)$  as  $\chi(uv) = \chi(vu)$  which is immediate since for any complex matrices  $A, B$  we have that  $\text{Tr}(AB) = \text{Tr}(BA)$ . □

**Proposition 2.3:** Let  $\rho^1: G \rightarrow \text{GL}(V_1)$  and  $\rho^2: G \rightarrow \text{GL}(V_2)$  be two linear representations with characters  $\chi_1$  and  $\chi_2$  respectively. Then

1. The character  $\chi$  of the direct sum representation  $V_1 \oplus V_2$  is  $\chi_1 + \chi_2$ .
2. The character  $\psi$  of the tensor product representation  $V_1 \otimes V_2$  is  $\chi_1 \cdot \chi_2$ .

**Proof:** Let  $R_s^1, R_s^2$  be the matrix forms of  $\rho_s^1$  and  $\rho_s^2$  respectively. Then the matrix form  $R_s$  of the representation of  $V_1 \oplus V_2$  is given by

$$R_s = \begin{pmatrix} R_s^1 & 0 \\ 0 & R_s^2 \end{pmatrix}$$

and thus  $\text{Tr}(R_s) = \text{Tr}(R_s^1) + \text{Tr}(R_s^2)$ . Let  $(e_{i_1})$  and  $(e_{i_2})$  be bases for  $V_1$  and  $V_2$ . Then we have that

$$\psi(s) = \sum_{i_1 i_2} r_{i_1 i_1}(s) \cdot r_{i_2 i_2}(s) = \left( \sum_{i_1} r_{i_1 i_1}(s) \right) \cdot \left( \sum_{i_2} r_{i_2 i_2}(s) \right) = \chi_1(s) \cdot \chi_2(s).$$

□

**Proposition 2.4:** Let  $\rho: G \rightarrow \text{GL}(V)$  be a linear representation of  $G$  with character  $\chi$ . Let  $\chi_\sigma^2$  be the character of  $\text{Sym}^2(V)$  and  $\chi_\alpha^2$  be the character of  $\text{Alt}^2(V)$  from Definition 1.16. Then

$$\begin{aligned} \chi_\sigma^2(s) &= \frac{1}{2}(\chi(s)^2 + \chi(s^2)) \\ \chi_\alpha^2(s) &= \frac{1}{2}(\chi(s)^2 - \chi(s^2)) \end{aligned}$$

which directly implies that  $\chi_\sigma^2 + \chi_\alpha^2 = \chi$ .

**Proof:** Let  $s \in G$  and  $(e_i)$  a basis of  $V$  consisting solely of eigenvectors for  $\rho_s$ . Then  $\rho_s(e_i) = \lambda_i e_i$  for some  $\lambda_i \in \mathbb{C}$ . Thus

$$\chi(s) = \sum \lambda_i \quad \chi(s^2) = \sum \lambda_i^2.$$

We also have that

$$\begin{aligned} (\rho_s \otimes \rho_s)(e_i \otimes e_j + e_j \otimes e_i) &= \lambda_i \lambda_j (e_i \otimes e_j + e_j \otimes e_i) \\ (\rho_s \otimes \rho_s)(e_i \otimes e_j - e_j \otimes e_i) &= \lambda_i \lambda_j (e_i \otimes e_j - e_j \otimes e_i) \end{aligned}$$

which yields that

$$\chi_\sigma^2(s) = \sum_{i \leq j} \lambda_i \lambda_j = \sum \lambda_i^2 + \sum_{i < j} \lambda_i \lambda_j = \frac{1}{2} \left( \sum \lambda_i \right)^2 + \frac{1}{2} \sum \lambda_i^2 \chi_\alpha^2(s) = \sum_{i < j} \lambda_i \lambda_j = \frac{1}{2} \left( \sum \lambda_i \right)^2 - \frac{1}{2} \sum \lambda_i^2.$$

The proposition then directly follows. Note that the equality  $\chi_\sigma^2 + \chi_\alpha^2 = \chi^2$  directly reflects the fact that  $V \otimes V = \text{Sym}^2(V) \oplus \text{Alt}^2(V)$ .  $\square$

**Proposition 2.5 (Schur's Lemma):** Let  $\rho^1: G \rightarrow \text{GL}(V_1)$  and  $\rho^2: G \rightarrow \text{GL}(V_2)$  be two irreducible representations of  $G$ . Let  $f: V_1 \rightarrow V_2$  be a linear map such that  $f \circ \rho_s^1 = \rho_s^2 \circ f$  for all  $s \in G$ . Then

1. If  $\rho^1$  and  $\rho^2$  are not isomorphic, then  $f = 0$
2. If  $V_1 = V_2$  and  $\rho^1 = \rho^2$  then  $f$  is a *homothety*, a scalar multiple of the identity.

**Proof:** The case of  $f = 0$  is trivial, so suppose that  $f \neq 0$ . Let  $W_1 = \ker(f)$  and  $W_2 = \text{im}(f)$ . Then for  $x \in W_1$  we have that  $f(\rho_s^1(x)) = \rho_s^2(f(x)) = 0$  which means that  $\rho_s^1(x) \in W_1$ . Thus  $W_1$  is stable under  $G$  and irreducibility of  $V_1$  combined with the assumption that  $f \neq 0$  implies that  $W_1 = 0$ . Similarly, we have that for  $f(x) \in W_2$ , we have that  $\rho_s^2(f(x)) = f(\rho_s^1(x)) \in W_2$ , so  $\rho_s^2(f(x)) \in W_2$ . Thus  $W_2$  is also stable under  $G$  meaning that by a similar argument,  $W_2 = V_2$ . Since  $\ker(f) = 0$  and  $\text{im}(f) = V_2$ , we must have that  $f$  is an isomorphism  $V_1 \rightarrow V_2$ . This proves the first claim.

Now suppose that  $V_1 = V_2$ ,  $\rho^1 = \rho^2$ , and that  $\lambda$  is some eigenvalue of  $f$ . Let  $f' = f - \lambda$ . Since  $\lambda$  is an eigenvalue, then  $\ker(f') \neq 0$ . However, we also have that  $f' \circ \rho_s^1 = \rho_s^2 \circ f'$ . The first part of this proof shows that this implies that  $f' = 0$ . Thus,  $f = \lambda$  and  $f$  is a homothety.  $\square$

**Corollary 2.6:** Let  $\rho^1: G \rightarrow \text{GL}(V_1)$  and  $\rho^2: G \rightarrow \text{GL}(V_2)$  be two irreducible representations of  $G$ . Let  $h: V_1 \rightarrow V_2$  and define  $h^0$  such that

$$h^0 = \frac{1}{|G|} \sum_{t \in G} (\rho_t^2)^{-1} \circ h \circ \rho_t^1.$$

Then

1. If  $\rho^1$  and  $\rho^2$  are not isomorphic, then  $h^0 = 0$
2. If  $V_1 = V_2$  and  $\rho^1 = \rho^2$ , then  $h^0$  is a homothety of ratio  $\frac{1}{n} \text{Tr}(h)$ , with  $n = \dim(V_1)$ .

**Proof:** First for  $s \in G$  we have that

$$(\rho_s^2)^{-1} \circ h^0 \circ \rho_s^1 = \frac{1}{|G|} \sum_{t \in G} (\rho_s^2)^{-1} (\rho_t^2)^{-1} \circ h \circ \rho_t^1 \rho_s^1 = \frac{1}{|G|} \sum_{t \in G} (\rho_{ts}^2)^{-1} \circ h \circ \rho_{ts}^1 = h^0.$$

Thus, Proposition 2.5 applies to  $h^0$  and in the first case  $h^0 = 0$  and in the second  $h^0$  is a homothety of scalar  $\lambda$ . Moreover we have that

$$n \cdot \lambda = \text{Tr}(\lambda) = \frac{1}{|G|} \sum_{t \in G} \text{Tr}((\rho_t^1)^{-1} \circ h \circ \rho_t^1) = \text{Tr}(h).$$

Thus,  $\lambda = \frac{1}{n} \text{Tr}(h)$ .  $\square$

Consider Corollary 2.6 in matrix form where  $\rho_s^1 = (r_{i_1 j_1}(s))$  and  $\rho_s^2 = (r_{i_2 j_2}(s))$ . Then our linear map  $h$  is given by the matrix  $(x_{i_2 i_1})$  and similarly  $h^0$  is given by the matrix  $(x_{i_2 i_1}^0)$ . Then by definition of  $h^0$  we have that

$$x_{i_2 i_1}^0 = \frac{1}{|G|} \sum_{t \in G, j_1, j_2} r_{i_2 j_2}(t^{-1}) \cdot x_{j_2 j_1} \cdot r_{j_1 i_1}(t).$$

In case (1) of Corollary 2.6, the right hand side must vanish and so all coefficients of 0:

**Corollary 2.7:** In case (1) of Corollary 2.6 we have that

$$\frac{1}{|G|} \sum_{t \in G} r_{i_2 j_2}(t^{-1}) \cdot r_{j_1 i_1}(t) = 0$$

for all  $i_1, j_1, i_2, j_2$ .

We now consider case (2) of Corollary 2.6 in the matrix form. We have that  $h^0 = \lambda$ , with  $\lambda = \frac{1}{n} \text{Tr}(h)$ , meaning that  $x_{i_2 i_1}^0 = \lambda \delta_{i_2 i_1}$ . That is,  $\lambda = \frac{1}{n} \sum \delta_{i_2 i_1} \cdot x_{i_2 i_1}$ . This we have that

$$\frac{1}{|G|} \sum_{t \in G, j_1, j_2} r_{i_2 j_2}(t^{-1}) \cdot x_{j_2 j_1} \cdot r_{j_1 i_1}(t) = \frac{1}{n} \sum_{j_1, j_2} \delta_{i_2 i_1} \cdot \delta_{j_2 j_1} \cdot x_{j_2 j_1}.$$

Equating coefficients of the  $x_{j_2 j_1}$  yields the following corollary:

**Corollary 2.8:** In case (2) of Corollary 2.6 we have that

$$\frac{1}{|G|} \sum_{t \in G} r_{i_2 j_2}(t^{-1}) \cdot r_{j_1 i_1}(t) = \frac{1}{n} \delta_{i_2 i_1} \cdot \delta_{j_2 j_1}.$$

We may generalize these notions to further functions. Let  $\phi, \psi$  be **<< complex valued? >>** functions on  $G$ . Define

$$\langle \phi, \psi \rangle := \frac{1}{|G|} \sum_{t \in G} \phi(t^{-1}) \cdot \psi(t) = \frac{1}{|G|} \sum_{t \in G} \phi(t) \cdot \psi(t^{-1}).$$

Then  $\langle \phi, \psi \rangle = \langle \psi, \phi \rangle$  and  $\langle \phi, \psi \rangle$  is linear in  $\phi$  and in  $\psi$ . Thus, Corollary 2.7 and Corollary 2.8 respectively become

$$\langle r_{i_2 j_2}, r_{j_1 i_1} \rangle = 0 \qquad \langle r_{i_2 j_2}, r_{j_1 i_1} \rangle = \frac{1}{n} \delta_{i_2 i_1} \cdot \delta_{j_2 j_1}.$$

If the matrices  $(r_{ij}(t))$  are unitary, realized by a suitable choice of basis, then  $r_{ij}(t^{-1}) = r_{ji}(t)^*$  and Corollary 2.7 and Corollary 2.8 are just *orthogonality relations*.



**Definition 2.9 (Scalar Product):** If  $\phi, \psi$  are two complex valued functions on  $G$ , then let

$$(\phi | \psi) := \frac{1}{|G|} \sum_{t \in G} \phi(t) \psi(t)^*.$$

This is a *scalar product*. It is linear in  $\phi$ , semilinear in  $\psi$ , and  $(\phi | \phi) > 0$  for all  $\phi \neq 0$ .

Define  $\check{\psi}(t) := \psi(t^{-1})^*$ . Then  $(\phi | \psi) = \langle \phi, \check{\psi} \rangle$ . In particular, suppose  $\chi$  is a character so that by Proposition 2.2 we have that  $\chi = \check{\chi}$  then for all complex valued functions  $\phi$  on  $G$  we have that  $(\phi | \chi) = \langle \phi, \chi \rangle$ . Thus, we may use the two interchangeably in the context of characters.

**Theorem 2.10:**

1. If  $\chi$  is the character of an irreducible representation, we have that  $(\chi | \chi) = 1$ , i.e.  $\chi$  has “norm 1.”
2. If  $\chi$  and  $\chi'$  are characters of two non-isomorphic irreducible representations, then  $(\chi | \chi') = 0$ , i.e.  $\chi$  and  $\chi'$  are “orthogonal.”

**Proof:** Suppose  $\rho$  is an irreducible representation with matrix form  $\rho_t = (r_{ij}(t))$  and  $\chi$  its character. Then  $\chi(t) = \sum r_{ii}(t)$  and so

$$(\chi | \chi) = \langle \chi, \chi \rangle = \sum_{i,j} \langle r_{ii}, r_{jj} \rangle = \frac{\delta_{ij}}{n}$$

where the last equality is by Corollary 2.8 and  $n$  is the degree of  $\rho$ . Thus

$$(\chi | \chi) = \sum_{i,j} \frac{\delta_{ij}}{n} = \frac{n}{n} = 1.$$

This proves the first claim. Applying Corollary 2.7 yields the second claim □

**Theorem 2.11:** Let  $V$  be a linear representation of  $G$  with character  $\phi$  such that  $V$  decomposes into a direct sum of irreducible representations  $V = W_1 \oplus \cdots \oplus W_k$ . Then if  $W$  is an irreducible representation with character  $\chi$ , the number of  $W_i$  isomorphic to  $W$  is equal to the scalar product  $(\phi | \chi) = \langle \phi, \chi \rangle$ .

**Proof:** Let  $\chi_i$  be the character of  $W_i$ . Then by Proposition 2.3 we have that  $\phi = \chi_1 + \cdots + \chi_k$ . By linearity of  $(\cdot | \cdot)$  in the first argument we have that  $(\phi | \chi) = (\chi_1 | \chi) + \cdots + (\chi_k | \chi)$ . The result follows by Theorem 2.10. □

**Corollary 2.12:** Let  $V$  be a linear representation of  $G$  with character  $\phi$  such that  $V$  decomposes into a direct sum of irreducible representations  $V = W_1 \oplus \cdots \oplus W_k$ . Then if  $W$  is an irreducible representation with character  $\chi$ , the number of  $W_i$  isomorphic to  $W$  does not depend on the chosen decomposition.

**Proof:** Note that  $(\phi | \chi)$  does not depend on choice of decomposition. □

**Corollary 2.13:** Two representations are isomorphic if and only if they have the same character.

**Proof:** The forward direction is obvious, and the reverse is true by the prior corollary.  $\square$

Thus, our study of representations is reduced to that of the study of characters. If  $\chi_1, \dots, \chi_k$  are the distinct irreducible characters of  $G$  (**How do we know there are finitely many?**) and if  $W_1, \dots, W_k$  their corresponding representation, then each representation  $V$  of  $G$  is isomorphic to a direct sum

$$V = m_1 W_1 \oplus \dots \oplus m_h W_h \quad m_i \neq 0.$$

The character  $\phi$  of  $V$  is equal to  $m_1 \chi_1 + \dots + m_h \chi_h$  and we have that  $m_i = (\phi | \chi_i)$ . This is especially useful when considering the tensor product  $W_i \otimes W_j$  of two irreducible representations. It shows that the product  $\chi_i \cdot \chi_j$  decomposes into a sum  $\chi_i \chi_j = \sum m_{ij}^k \chi_k$ , each integer  $m_{ij}^k \geq 0$ . The orthogonality relations among the  $\chi_i$  imply that

$$(\phi | \phi) = \sum_{i=1}^h m_i^2.$$

We now obtain a useful irreducibility criterion:

**Theorem 2.14:** If  $\phi$  is the character of a representation  $V$ ,  $(\phi | \phi)$  is a positive integer and  $(\phi | \phi) = 1$  if and only if  $V$  is irreducible.

**Proof:** We have that  $\sum m_i^2 = 1$  if and only if one of the  $m_i = 1$  and all the others are equal to 0. This means that  $V$  is isomorphic to one of the  $W_i$ .  $\square$

We now explore the decomposition of the regular representation  $\rho: G \rightarrow \text{GL}(R)$  of a group  $G$  (Example 1.6). Suppose  $\chi_1, \dots, \chi_h$  are the irreducible characters of  $G$  with degrees  $n_1, \dots, n_k$ . Note that by Proposition 2.2,  $n_i = \chi_i(e)$ . Recall that  $R$  has basis  $(e_t)_{t \in G}$  where  $\rho_s(e_t) = e_{st}$ . This means that for  $s \neq e$ , the diagonal terms of the matrix for  $\rho_s$  are all 0, so  $\text{Tr}(\rho_s) = 0$ . On the otherhand, we have that

$$\text{Tr}(\rho_e) = \dim(R) = |G|.$$

**Proposition 2.15:** The character  $r_G$  of the regular representation is given by

$$r_G(e) = |G| \quad r_G(s) = 0 \text{ if } s \neq e.$$

**Corollary 2.16:** Every irreducible representation  $W_i$  is contained in the regular representation with multiplicity equal to its degree  $n_i$ .

**Proof:** By Theorem 2.11, the number of times  $W_i$  is contained in the regular representation is  $\langle r_G, \chi_i \rangle$ . We have that

$$\langle r_G, \chi_i \rangle = \frac{1}{|G|} \sum_{s \in G} r_G(s^{-1}) \chi_i(s) = \frac{1}{|G|} \cdot |G| \chi_i(1) = \chi_i(1) = n_i.$$

□

**Corollary 2.17:** 1. The degrees satisfy  $\sum_{i=1}^h n_i^2 = |G|$ .

2. if  $e \neq s \in G$ , we have that  $\sum_{i=1}^h n_i \chi_i(s) = 0$ .

**Proof:** By Corollary 2.16, we have that  $r_G(s) = \sum n_i \chi_i(s)$  for all  $s \in G$  **<< how? >>**. The claim is thus immediate. □

The above result lets us determine the irreducible representations of a group  $G$ . Suppose we have constructed some mutually non-isomorphic irreducible representations of degrees  $n_1, \dots, n_h$ . In order to check if we have found all such representations, it is necessary and sufficient to verify that  $n_1^2 + \dots + n_h^2 = |G|$ . Also, we shall later see that each of the  $n_i$  divide the order of  $G$ .

**Definition 2.18 (Class Function):** A function  $f$  on a group  $G$  is a *class function* if for all  $s, t \in G$ ,  $f(tst^{-1}) = f(s)$ .

**Proposition 2.19:** Let  $f$  be a class function on a group  $G$  and  $\rho : G \rightarrow \text{GL}(V)$  a linear representation of  $G$  with character  $\chi$ . Define  $\rho_f : V \rightarrow V$  by  $\rho_f = \sum_{t \in G} f(t) \rho_t$ . If  $V$  is irreducible of degree  $n$ , the  $\rho_f$  is a homothety of ratio  $\lambda$  where

$$\lambda = \frac{1}{n} \sum_{t \in G} f(t) \chi(t) = \frac{|G|}{n} (f \mid \chi^*).$$

**Proof:** We have that

$$\rho_s^{-1} \rho_f \rho_s = \sum_{t \in G} f(t) \rho_s^{-1} \rho_t \rho_s = \sum_{t \in G} f(t) \rho_{s^{-1}ts} = \sum_{t \in G} f(s^{-1}ts) \rho_{s^{-1}ts} = \rho_f.$$

Thus, by Proposition 2.5 we have that  $\rho_f$  is a homothety  $\lambda$ . The trace of  $\lambda$  is  $n\lambda$ . Thus, the trace of  $\rho_f$  is  $\sum_{t \in G} f(t) \text{Tr}(\rho_t) = \sum_{t \in G} f(t) \chi(t)$ . Thus,  $\lambda = \frac{1}{n} \sum_{t \in G} f(t) \chi(t) = \frac{|G|}{n} (f \mid \chi^*)$ . □

Let  $H$  be the space of class functions on  $G$ .

**Theorem 2.20:** The characters  $\chi_1, \dots, \chi_h$  of  $G$  form an orthonormal basis of  $H$ .

**Proof:** Note that Theorem 2.10 says that the  $\chi_i$  are all orthonormal to each other. To show that they generate  $H$ , it is enough to show that the only element of  $H$  orthogonal to  $\chi_i^*$  is 0. Let  $f$  be such an element. For each

representation  $\rho$  of  $G$ , let  $\rho_f$  be as in Proposition 2.19. Since  $f$  is orthogonal to the  $\chi_i^*$ , Proposition 2.19 says that  $\rho_f$  is 0 as long as  $\rho$  is irreducible. From the **<< direct sum decomposition >>**, we conclude that  $\rho_f$  is always 0. Now consider the regular representation of  $G$  and compute the image of the basis vector  $e_e$  under  $\rho_f$ :

$$0 = \rho_f(e_e) = \sum_{t \in G} f(t) \rho_t(e_e) = \sum_{t \in G} f(t) \rho_t.$$

Thus,  $f(t) = 0$  for each  $t \in G$  and  $f = 0$ . □

**Theorem 2.21:** The number of irreducible representations of  $G$ , up to isomorphism, is the number of conjugacy classes of  $G$ .

**Proof:** Let  $C_1, \dots, C_k$  be the distinct conjugacy classes of  $G$ . Then all class functions are constant on each class, their value determined by some  $\lambda_i$  for each  $C_i$ . These  $\lambda_i$  may be chosen arbitrarily. Thus, the dimension of the space  $H$  of class functions is equal to  $k$ . But we already know by Theorem 2.20 that the dimension of  $H$  is  $h$ , the number of irreducible representations of  $G$ . □

**Proposition 2.22:** Let  $s \in G$  and  $c(s)$  the number of elements in the conjugacy class of  $s$ .

1. We have  $\sum_{i=1}^h \chi_i(s)^* \chi_i(s) = \frac{|G|}{c(s)}$ .
2. For  $t$  not conjugate to  $s$ , we have  $\sum_{i=1}^h \chi_i(s)^* \chi_i(t) = 0$ .

**Proof:** Let  $f_s$  be the class function equal to 1 on the class of  $s$  and 0 otherwise. By Theorem 2.21, we have that

$$f_s = \sum_{i=1}^h \lambda_i \chi_i \qquad \lambda_i = (f_s | \chi_i) = \frac{c(s)}{|G|} \chi_i(s)^*.$$

We have then, for each  $t \in G$ , that

$$f_s(t) = \frac{c(s)}{|G|} \sum_{i=1}^h \chi_i(s)^* \chi_i(t).$$

If  $t = s$ , we get claim 1 and for  $t$  not conjugate to  $s$  we get claim 2. □

**Example 2.23 (Character Table of  $S_3$ ):** Consider the group  $S_3$ . There are three conjugacy classes: the identity (), the 3 transpositions, and the 2 cyclic permutations. Let  $t$  be one of the transpositions and  $c$  one of the cyclic permutations. Then  $t^2 = 1 = c^3$  and  $tc = c^2t$ . There are just two characters of degree 1: the unit character  $\chi_1$  and the character  $\chi_2$  giving the sign of the permutation **<< why? >>**. By Theorem 2.21, there exists one more irreducible character  $\theta$ . If  $n$  is the degree of  $\theta$ , then we must have that  $1 + 1 + n^2 = 6$ , so  $n = 2$ . By Proposition 2.15, we have that  $\chi_1 + \chi_2 + 2\theta$  is the character of the regular representation. Thus, we get the following *character table*:

**<< TODO: Center >>**

	1	$t$	$c$
$\chi_1$	1	1	1
$\chi_2$	1	-1	1
$\theta$	2	0	-1

We obtain an irreducible representation of  $G$  with character  $\theta$  by having  $G$  permute the coordinates of elements of  $\mathbb{C}^3$  satisfying  $x + y + z = 0$ .

Let  $\rho: G \rightarrow \text{GL}(V)$  be a linear representation of  $G$ . Recall that the direct sum decomposition of  $V$  into irreducible representation is not necessarily unique. Thus, we shall now define a “coarser” decomposition which has the advantage of being unique.

**Definition 2.24 (Canonical Decomposition of a Representation):** Let  $\chi_1, \dots, \chi_h$  be the distinct characters of the irreducible representations of  $W_1, \dots, W_h$  of  $G$  with degrees  $n_1, \dots, n_h$ . Let  $V = U_1 \oplus \dots \oplus U_m$  be a decomposition of  $V$  into a direct sum of irreducible representations. For  $i = 1, \dots, h$ , let  $V_i$  be the direct sum of the  $U_i$  which are isomorphic to  $W_i$ . Then  $V = V_1 \oplus \dots \oplus V_h$ . We have decomposed  $V$  into a direct sum of irreducible representations and combined the ones which are isomorphic to each other.

This decomposition satisfies some nice properties:

**Theorem 2.25:** 1. The decomposition  $V = V_1 \oplus V_h$  does not depend on the initially chosen decomposition of  $V$  into irreducibles.

2. The projection  $p_i: V \rightarrow V_i$  is given by

$$p_i = \frac{n_i}{|G|} \sum_{t \in G} \chi_i^*(t) \rho_t.$$

**Proof:** We shall prove claim 2 since claim 1 follows as the  $p_i$  determine the  $V_i$ . Let  $q_i = \frac{n_i}{|G|} \sum_{t \in G} \chi_i(t)^* \rho_t$ . By Proposition 2.19, we have that the restriction of  $q_i$  to an irreducible representation  $W$  with character  $\chi$  and degree  $n$  is a homothety of ratio  $\frac{n_i}{n}(\chi_i | \chi)$ . Thus,  $q_i$  is 0 if  $\chi_i \neq \chi$  and 1 if  $\chi = \chi_i$ . This yields that  $q_i$  is the identity on an irreducible representation isomorphic to  $W_i$ , and 0 on the others. Thus,  $q_i$  is the identity on  $V_i$  and 0 on  $V_j$  for  $j \neq i$ . Decomposing  $x \in V$  into  $x_i \in V_i$  such that  $x = x_1 + \cdots + x_h$  yields that

$$q_i(x) = q_i(x_1) + \cdots + q_i(x_h) = x_i.$$

Thus  $q_i = p_i$ . □

This allows us to decompose representations  $V$  in two stages. First, we determine  $V_1 \oplus \cdots \oplus V_h$ . This is done easily using the given formula for  $p_i$  in Theorem 2.25. Finally, for each  $V_i$  we may choose a decomposition of  $V_i$  into a direct sum of irreducible representations, each isomorphic to  $W_i$ . This last decomposition may be done in any number of ways.

**Example 2.26 (Decomposition of  $C_2$ ):** Let  $G = C_2 = \{e, s\}$  be the cyclic group of two elements generated by  $s$ . Let  $\rho: G \rightarrow \text{GL}(V)$  be any representation of  $C_2$ . Note that  $C_2$  has two irreducible representations of degree 1,  $W^+$  and  $W^-$  with respective characters  $\rho^+ = 1$  and  $\rho_s = -1$ . The canonical decomposition of  $V$  is  $V = V^+ \oplus V^-$ , where  $V^+$  consists of elements  $x \in V$  which are symmetric and  $V^-$  consists of elements which are antisymmetric. In other words,  $V^+$  consists of elements  $x \in V$  where  $\rho_s(x) = x$  and  $V^-$  consists of elements  $x \in V$  where  $\rho_s(x) = -x$ . This, the projections are

$$p^+(x) = \frac{1}{2}(x + \rho_s(x)) \qquad p^-(x) = \frac{1}{2}(x - \rho_s(x)).$$

To decompose  $V^+$  and  $V^-$  into irreducible components means to decompose these subspaces into a direct sum of lines, which can be in arbitrarily many ways.

## Exercises

**Exercise 2.1 (Ser77 2.1):** Let  $\chi, \chi'$  be the characters of two representations. Prove the following formulas:

$$\begin{aligned}(\chi + \chi')_{\sigma}^2 &= \chi_{\sigma}^2 + \chi'_{\sigma}^2 + \chi \chi' \\ (\chi + \chi')_{\alpha}^2 &= \chi_{\alpha}^2 + \chi'_{\alpha}^2 + \chi \chi'\end{aligned}$$

**Proof:** Let  $s \in G$ . Then by Proposition 2.4 we have that

$$\begin{aligned}(\chi + \chi')_{\sigma}^2(s) &= \frac{1}{2}((\chi + \chi')^2(s) + (\chi + \chi')(s^2)) \\ &= \frac{1}{2}(\chi(s)^2 + \chi'(s)^2 + 2\chi(s)\chi'(s) + \chi(s^2) + \chi'(s^2)) \\ &= \frac{1}{2}(\chi(s)^2 + \chi(s^2)) + \frac{1}{2}(\chi'(s) + \chi'(s^2)) + \chi(s)\chi'(s) = \chi_{\sigma}^2(s) + \chi'_{\sigma}^2(s) + \chi(s)\chi'(s).\end{aligned}$$

Since this holds for all  $s \in G$ , the formula holds in general. The proof of the other formula is similar.  $\square$

**Exercise 2.2 (Ser77 2.2):** Let  $X$  be a finite set on which  $G$  acts, and  $\rho: G \rightarrow \text{GL}(V)$  the corresponding permutation representation (Example 1.7), and  $\chi_X$  the character of  $\rho$ . Then show that for  $s \in G$ ,  $\chi_X(s)$  is equal to the number of elements fixed by  $s$ .

**Proof:** Suppose  $X = [n]$  and so  $s \in S_n$ , meaning  $G \leq S_n$ . We may assume this without loss of generality. Note that  $R_s = (r_{ij}(s))$  where  $r_{ij}(s) = 1$  if  $s(j) = i$  and 0 otherwise. We want to count the number of elements in  $[n]$  fixed by  $s$ , i.e. the number of  $i$  such that  $\sigma(i) = i$ . These correspond exactly to the entries in  $R_s$  where  $r_{ii}(s) = 1$ . Thus, the claim follows.  $\square$

**Exercise 2.3 (Ser77 2.6):** Let  $G$  act on a finite set  $X$ ,  $\rho$  the corresponding permutation representation, and  $\chi$  its character.

1. Let  $c$  be the number of distinct orbits. Show that  $c$  is equal to the number of times  $\rho$  contains the unit representation 1. Deduce that  $(\chi | 1) = c$ . In particular if  $G$  is transitive and thus  $c = 1$ , then  $\rho = 1 \oplus \theta$  where  $\theta$  does not contain the unit representation. If  $\psi$  is the character of  $\theta$ , then  $\chi = 1 + \psi$  and  $(\psi | 1) = 0$ .
2. Let  $G$  act on the product  $X \times X$  in the natural way. Show that the character of the corresponding permutation representation is equal to  $\chi^2$ .
3. Suppose that  $G$  is transitive on  $X$  and  $|X| \geq 2$ . We say  $G$  is *doubly transitive* if for all  $x, y, x', y' \in X$  with  $x \neq y$  and  $x' \neq y'$  there exists  $s \in G$  such that  $s(x, y) = (sx, sy) = (x', y')$ . Prove that the following are equivalent:

- (a)  $G$  is doubly transitive.
- (b) The action of  $G$  on  $X \times X$  has two orbits, the diagonal and the complement.
- (c)  $(\chi^2 | 1) = 2$
- (d) The representation  $\theta$  defined in the first part of this exercise is irreducible.

**Proof:** We know that the number of times the unit representation is contained in  $\chi$  is equal to  $(\chi | 1)$  by Theorem 2.11. By [Exercise 2.5](#) we have that  $(\chi | 1) = \frac{1}{|G|} \sum_{s \in G} \chi(s)$ . We prove that  $\frac{1}{|G|} \sum \chi(s) = c$  by double counting. Consider the set  $\{(s, x) \in G \times X \mid s \cdot x = x\}$ . Then we have that

$$\sum_{x \in X} |G_x| = \sum_{x \in X} |\{s \in G \mid s \cdot x = x\}| = |\{(s, x) \in G \times X \mid s \cdot x = x\}| = \sum_{s \in G} |\{x \in X \mid s \cdot x = x\}| = \sum_{s \in G} \chi(s).$$

Let  $O_1, \dots, O_c$  be the distinct orbits. By the Orbit-Stabilizer theorem, each  $O_i$  is in bijection with  $G/G_x$  for all  $x \in O_i$ . Note that the orbits  $O_i$  partition  $X$ . Thus we have that

$$\sum_{s \in G} \chi(s) = \sum_{i=1}^c \sum_{x \in O_i} |G_x| = \sum_{i=1}^c \sum_{x \in O_i} \frac{|G|}{|O_i|} = c \cdot |G|$$

and  $\frac{1}{|G|} \sum_{s \in G} \chi(s) = c$ . Following this, the rest of the claim is immediate.

Now suppose that  $\phi$  is the character of the permutation representation of  $G \curvearrowright X \times X$ . Then by Exercise 2.2,  $\phi(s)$  is equal to the number of elements fixed by  $s$ . An element  $(x, y) \in X \times X$  is fixed by  $s \in G$  if and only if both  $x$  and  $y$  are fixed. Thus if there are  $\chi(s)$  elements of  $X$  fixed by  $s$ , then  $\chi^2(s)$  elements of  $X \times X$  are fixed by  $s$  and  $\phi = \chi^2$ .

To prove 3, we have that  $(a) \iff (b)$  is immediate and  $(b) \iff (c)$  follows from 1 and 2. Now suppose  $(c)$  holds and let  $\psi$  be the character of  $\theta$ . Then  $1 + \psi = \theta$ . Since  $(\chi | 1) = (1 | 1) = 1$  we must have that  $(\psi | 1) = 0$ . Since  $\chi^2 = 1 + 2\psi + \psi^2$ , we have that  $(c)$  is equivalent to saying  $(\psi^2 | 1) = 1$ . Thus

$$\frac{1}{|G|} \sum_{s \in G} \psi(s)^2 = 1.$$

However, note that  $\psi(s)$  is real valued, not just complex valued. This is because  $\chi$  is real valued, it counts fixed points, and clearly 1 is real valued. Thus  $\psi^* = \psi(s)^*$  and so the above equality implies that  $(\psi | \psi = 1)$ . By Theorem 2.14, we have that this is true if and only if  $\theta$  is irreducible, i.e.  $(c) \iff (d)$  holds.  $\square$



# Bibliography

- [Ful96] William Fulton. *Young Tableaux: With Applications to Representation Theory and Geometry*. London Mathematical Society Student Texts. Cambridge University Press, 1996. DOI: [10.1017/CB09780511626241](https://doi.org/10.1017/CB09780511626241).
- [Pan23] Greta Panova. *Computational Complexity in Algebraic Combinatorics*. 2023. arXiv: [2306.17511](https://arxiv.org/abs/2306.17511) [math.CO].
- [Rez20] Charles Rezk. *A short course on finite group representations*. 2020. URL: <https://rezk.web.illinois.edu/Finite%20Group%20Reps/short-course-finite-group-representations.html>.
- [Ser77] Jean-Pierre Serre. *Linear Representations of Finite Groups*. Springer New York, 1977. ISBN: 9781468494587. DOI: [10.1007/978-1-4684-9458-7](https://doi.org/10.1007/978-1-4684-9458-7).