

Algorithms in Invariant Theory Exercises

With 0 Figures

Anakin Dey

Last Edited on 4/3/24 at 00:24

Contents

1	Introduction
---	--------------

1

Preface

A core interest of mine (at the time of writing this) is algorithms in the context of commutative algebra and algebraic geometry. As such, Bernd Sturmfel's text *Algorithms in Invariant Theory* [\[Str08\]](#) is a good resource for first learning this stuff. This solely just contains exercises for now. Perhaps in the future I'll include notes and some source code

Chapter 1

Introduction

Symmetric Polynomials

Exercise 1.1 (Str08 1.1.5): Prove the following explicit formula for elementary symmetric polynomials in terms of the power sums [Mac98, Page 29].

$$\sigma_k = \frac{1}{k!} \det \begin{pmatrix} p_1 & 1 & 0 & \cdots & 0 \\ p_2 & p_1 & 2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ p_{k-1} & p_{k-2} & \cdots & p_1 & k-1 \\ p_k & p_{k-1} & \cdots & \cdots & p_1 \end{pmatrix}$$

Proof: From [Page 23, 2.11'] [Mac98], we have the following identities

$$1 \cdot \sigma_1 = p_1,$$

$$2 \cdot \sigma_2 = p_1 \sigma_1 - p_2,$$

$$3 \cdot \sigma_3 = p_1 \sigma_2 - p_2 \sigma_1 + p_3,$$

$$\vdots$$

$$k \cdot \sigma_k = \sum_{r=1}^k (-1)^{r-1} p_r \sigma_{k-r}.$$

Treating the σ_i as indeterminants, we can re-express the above system of equations:

$$\begin{aligned}
p_1 &= 1 \cdot \sigma_1, \\
p_2 &= p_1 \sigma_1 - 2 \cdot \sigma_2, \\
p_3 &= p_2 \sigma_1 - p_1 \sigma_2 + 3 \cdot \sigma_3, \\
&\vdots \\
p_k &= \left(\sum_{r=1}^{k-1} (-1)^{r-1} p_{k-r} \sigma_r \right) + (-1)^{k-1} \cdot k \sigma_k.
\end{aligned}$$

Consider $\sigma_1, -\sigma_2, \sigma_3, \dots, (-1)^n \sigma_{n-1}, \sigma_n$ as indeterminants. Thus, we obtain the following matrix equation:

$$\begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ p_1 & 2 & 0 & \cdots & 0 \\ p_2 & p_1 & 3 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ p_{k-1} & p_{k-2} & \cdots & \cdots & (-1)^k \cdot k \end{pmatrix} \begin{pmatrix} \sigma_1 \\ -\sigma_2 \\ \vdots \\ (-1)^k \sigma_{k-1} \\ \sigma_k \end{pmatrix} = \begin{pmatrix} p_1 \\ p_2 \\ \vdots \\ p_{k-1} \\ p_k \end{pmatrix}.$$

Then, Cramer's rule yields that

$$\begin{aligned}
\sigma_k &= \det \begin{pmatrix} 1 & 0 & 0 & \cdots & p_1 \\ p_1 & 2 & 0 & \cdots & p_2 \\ p_2 & p_1 & 3 & \cdots & p_3 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ p_{k-1} & p_{k-2} & \cdots & \cdots & p_k \end{pmatrix} \det \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ p_1 & 2 & 0 & \cdots & 0 \\ p_2 & p_1 & 3 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ p_{k-1} & p_{k-2} & \cdots & \cdots & (-1)^k \cdot k \end{pmatrix}^{-1} \\
&= \frac{(-1)^k}{k!} \det \begin{pmatrix} 1 & 0 & 0 & \cdots & p_1 \\ p_1 & 2 & 0 & \cdots & p_2 \\ p_2 & p_1 & 3 & \cdots & p_3 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ p_{k-1} & p_{k-2} & \cdots & \cdots & p_k \end{pmatrix} \\
&= \frac{(-1)^k}{k!} \cdot (-1)^k \det \begin{pmatrix} p_1 & 1 & 0 & \cdots & 0 \\ p_2 & p_1 & 2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ p_{k-1} & p_{k-2} & \cdots & p_1 & k-1 \\ p_k & p_{k-1} & \cdots & \cdots & p_1 \end{pmatrix} = \frac{1}{k!} \det \begin{pmatrix} p_1 & 1 & 0 & \cdots & 0 \\ p_2 & p_1 & 2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ p_{k-1} & p_{k-2} & \cdots & p_1 & k-1 \\ p_k & p_{k-1} & \cdots & \cdots & p_1 \end{pmatrix}.
\end{aligned}$$

□

Gröbner Bases

Exercise 1.2 (Str08 1.2.1): Let $<$ be an monomial order and let I be any ideal in $\mathbb{C}[x_1, \dots, x_n]$. A monomial m is called *minimally nonstandard* if M is nonstandard and all proper divisors of m are standard. Show that the set of minimally nonstandard monomials is finite.

Proof: ⟨ Have ugly inductive proof ⟩

□

Exercise 1.3 (Str08 1.2.2): Prove that the reduced Gröbner basis \mathcal{G}_{red} of I with respect to $<$ is unique (up to multiplicative constants from \mathbb{C}). Given an algorithm which transforms an arbitrary Gröbner basis into \mathcal{G}_{red} .

Proof: This is [CLO15, Chapter 2, §7, Theorem 5].

□

Exercise 1.4 (Str08 1.2.3): Let $I \subseteq \mathbb{C}[x_1, \dots, x_n]$ be an ideal, given by a finite set of generators. Using Gröbner bases, describe an algorithm for computing the *elimination ideals* $I \cap \mathbb{C}[x_1, \dots, x_i]$ for $i = 1, \dots, n-1$, and prove its correctness.

Proof: This is [CLO15, Chapter 3, §1, Theorem 2].

□

Exercise 1.5 (Str08 1.2.4): Find a characterization of all monomial orders on the polynomial ring $\mathbb{C}[x_1, x_2]$. Hint: each variable receives a certain “weight” which behaves additively under multiplication of variables. Generalize your result to n variables.

Bibliography

- [CLO15] D.A. Cox, J. Little, and D. O'Shea. *Ideals, Varieties, and Algorithms: An Introduction to Computational Algebraic Geometry and Commutative Algebra*. Undergraduate Texts in Mathematics. Springer International Publishing, 2015. ISBN: 9783319167213. URL: <https://books.google.com/books?id=yL7yCAAQBAJ>.
- [Mac98] Ian Grant Macdonald. *Symmetric functions and Hall polynomials*. Oxford university press, 1998.
- [Str08] Bernd Strumfels. *Algorithms in Invariant Theory*. Springer Vienna, 2008. ISBN: 9783211774175. DOI: [10.1007/978-3-211-77417-5](http://dx.doi.org/10.1007/978-3-211-77417-5). URL: <http://dx.doi.org/10.1007/978-3-211-77417-5>.