

Banner appropriate to article type will appear here in typeset article

Strain-coupled one-dimensional turbulence: a single-line closure for the rapid pressure–strain and its measured limits

Sparsh Sharma¹ and Alan R. Kerstein²

¹German Aerospace Center (DLR), 38108 Braunschweig, Germany

²Consultant, Danville, CA 94526, USA

Corresponding author: Sparsh Sharma, sparsh.sharma@dlr.de

(Received xx; revised xx; accepted xx)

The rapid distortion of turbulence by a mean strain is central to flows from wind-tunnel contractions to the stagnation regions of lifting surfaces, yet its economical prediction founders on a closure problem: the rapid pressure–strain term is a nonlocal three-dimensional spectral integral, while affordable models evolve at most one-dimensional or single-point information. We present a strain-coupled formulation of one-dimensional turbulence (ODT) and, as its analytical core, a systematic treatment of what a single line of turbulence data determines about the rapid pressure–strain term: the obstruction to a single-line closure is stated exactly, sharp bounds on the rapid term are derived from line data alone, and the closure assumption of axisymmetry about the line is not assumed but measured—it is exact for axisymmetric distortion and its plane-strain error is quantified as a function of accumulated strain. A strained-box direct numerical simulation with an exact rapid-distortion companion provides the independent spectral benchmark: it shows that the correct linear reference for line-projected spectra is broadband rather than a rigid spectral translation, and a three-way comparison localises the model’s remaining deficiency in its wavenumber-uniform rapid kinematics, quantified by a distance-from-linear-theory envelope as a function of strain rapidity. Hierarchical application of the model’s energy-redistributing kernel to eddy sub-scales is shown to supply the measured missing ingredient—scale-local relaxation—where it reaches the scales of the imbalance.

Key words:

1. Introduction

[M4: introduction rewrite — see markers above.]

2. Strain-coupled one-dimensional turbulence

The linear rapid-distortion equations, recalled below, contain two distinct ingredients: an evolution of the Fourier *amplitudes*, comprising production and the solenoidal pressure projection, and an evolution of the *wavevectors* that carries the kinematic straining of scales. The pressure part of the amplitude equation requires the full three-dimensional wavevector and is therefore not directly representable on a one-dimensional line; that incompatibility organises this section. The distortion physics is split according to its structure and each part assigned to the mechanism that can carry it faithfully: the production, which is local in physical space, is imposed as a continuous forcing of the line velocity (§ 2.3); the rapid pressure–strain, which is not, is supplied as a continuous, energy-conserving redistribution operator constructed to reproduce the homogeneous rapid-distortion evolution of the component energies under an explicit spectral closure (§ 2.4); and the wavevector kinematics are recovered by a consistent straining of the domain (§ 2.5). The slow, turbulence-driven pressure–strain and the inertial cascade remain the discrete eddy events of the standard model, reviewed first.

Throughout, $A_{ij} = \partial U_i / \partial x_j$ denotes the imposed mean velocity gradient, taken uniform across the line at each instant; the medium is incompressible, $A_{ii} = 0$. The line carries the fluctuation field $u_i(y, t)$ about this mean; $R_{ij} \equiv \overline{u_i u_j}$ is the Reynolds-stress tensor, the overline denoting the average over the statistically homogeneous directions (equivalently, the ensemble of independent realisations), and $k_t \equiv \frac{1}{2} R_{ii}$ the turbulent kinetic energy. In the homogeneous configurations used to establish and test the formulation the spatial line average and the ensemble average coincide.

2.1. The standard model

This subsection reviews the one-dimensional turbulence model of Kerstein (1999) in the vector-valued formulation of Kerstein *et al.* (2001), in the notation used throughout; nothing in it is original. The model evolves a three-component velocity field $u_i(y, t)$, $i = 1, 2, 3$, on a one-dimensional domain whose coordinate y is identified with the direction x_2 . The fields represent individual realisations; ensemble statistics are accumulated over many independent simulations. Two interleaved mechanisms advance the state: between events the fields obey ordinary one-dimensional diffusion, $(\partial_t - \nu \partial_y^2) u_i = 0$, with kinematic viscosity ν ; and a stochastic sequence of instantaneous *eddy events* represents the effect of individual turbulent overturns.

Because a continuum rotational motion cannot be represented on a line, an eddy event is an instantaneous, measure-preserving rearrangement of the profiles together with an additive velocity change,

$$u_i(y) \longrightarrow u_i(f(y)) + c_i K(y), \quad (2.1)$$

where f is the *triplet map* (Kerstein 1991, 1999), defined on an interval $[y_0, y_0 + l]$ of size l by

$$f(y) = y_0 + \begin{cases} 3(y - y_0), & y_0 \leq y \leq y_0 + \frac{1}{3}l, \\ 2l - 3(y - y_0), & y_0 + \frac{1}{3}l \leq y \leq y_0 + \frac{2}{3}l, \\ 3(y - y_0) - 2l, & y_0 + \frac{2}{3}l \leq y \leq y_0 + l, \\ y - y_0, & \text{otherwise,} \end{cases} \quad (2.2)$$

written through its inverse, so that fluid at $f(y)$ is placed at y : the map compresses the interval to one third of its length, inserts three copies, reverses the central copy, and leaves the exterior unchanged. It is *measure preserving* (the non-local analogue of a divergence-

free rearrangement), *continuous*, and *scale local* (it changes gradients by at most an order-unity factor) (Kerstein *et al.* 2001)—the three invariants whose preservation under an imposed mean strain is established in § 2.3. The additive term is built from the *kernel*

$$K(y) \equiv y - f(y), \quad \int K(y) dy = 0, \quad \int K^2(y) dy = \frac{4}{27} l^3, \quad (2.3)$$

non-zero only within the eddy interval; the first identity guarantees that the additive change conserves the momentum of each component, and the second fixes the energetics of the redistribution.

The kernel term is the model surrogate for the pressure-mediated redistribution of energy among velocity components (Wunsch & Kerstein 2001; Kerstein *et al.* 2001). Requiring that an event redistribute kinetic energy among components while conserving the total, with a rule invariant under permutation of the component indices, gives

$$\Delta E_i = \alpha \sum_j T_{ij} Q_j, \quad \mathbf{T} = \frac{1}{2} \begin{pmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{pmatrix}, \quad Q_i \equiv \frac{27}{8} \rho_0 l u_{i,K}^2, \quad (2.4)$$

where $u_{i,K} \equiv l^{-2} \int u_i(f(y)) K(y) dy$, Q_i is the available kinetic energy of component i within the eddy, ρ_0 the constant density, and $\alpha \in [0, 1]$ the parameter setting the fraction of extractable energy transferred towards isotropy. Solving the per-component energy balance for the amplitudes yields

$$c_i = \frac{27}{4l} \left[-u_{i,K} + \text{sgn}(u_{i,K}) \sqrt{u_{i,K}^2 + \alpha \sum_j T_{ij} u_{j,K}^2} \right], \quad (2.5)$$

real for $0 \leq \alpha \leq 1$. The transfer matrix $\mathbf{T}$ equalises the component energies: the construction is a model of the return-to-isotropy (*slow*) part of the pressure–strain interaction, a correspondence made precise in § 2.2. The standard model contains no counterpart of the rapid, mean-strain-driven part.

The event sequence is a Poisson process whose rate reflects the instantaneous flow state: each candidate eddy (y_0, l) is assigned a time scale τ through $(l/\tau)^2 \sim u_{2,K}^2 + \alpha \sum_j T_{2j} u_{j,K}^2 - Z v^2/l^2$, the last term a viscous penalty with order-unity parameter Z , and the event-rate density is $\lambda(y_0, l; t) = C/(l^2 \tau)$ with rate constant C ; candidates with negative discriminant are forbidden. The inertial-range cascade and the $k^{-5/3}$ spectrum are *outcomes* of repeated sampling from this rate, not inputs (Kerstein 1999); C , Z and a large-eddy suppression parameter are the only tunable constants (Stephens & Lignell 2021). Finally, one diagnostic caveat of Kerstein *et al.* (2001) is respected throughout: because the line velocity does not itself advect fluid, off-diagonal products of line velocity components are not momentum fluxes; diagonal component spectra and variances are read from the fields, whereas shear stresses are read from eddy-induced transport.

2.2. Rapid and slow pressure–strain, and linear rapid distortion

The Reynolds-stress transport equation for a fluctuating field about a mean with constant gradient A_{ij} reads, in homogeneous turbulence,

$$\frac{dR_{ij}}{dt} = \mathcal{P}_{ij} + \Pi_{ij} - \varepsilon_{ij}, \quad \mathcal{P}_{ij} = -A_{ik} R_{kj} - A_{jk} R_{ki}, \quad \Pi_{ij} = \left\langle \frac{p'}{\rho} (\partial_j u'_i + \partial_i u'_j) \right\rangle, \quad (2.6)$$

with ε_{ij} the dissipation tensor (Pope 2000). Taking the divergence of the fluctuating momentum equation gives the pressure as the solution of a Poisson equation whose source

splits into a part linear in the mean gradient and a part quadratic in the fluctuations,

$$\frac{1}{\rho} \nabla^2 p' = \underbrace{-2 A_{lm} \partial_l u'_m}_{\text{rapid}} - \underbrace{\partial_l \partial_m (u'_l u'_m - \langle u'_l u'_m \rangle)}_{\text{slow}}, \quad (2.7)$$

81 and the pressure–strain correlation inherits the decomposition $\Pi_{ij} = \Pi_{ij}^{(r)} + \Pi_{ij}^{(s)}$ into a
 82 *rapid* part driven by the mean strain and a *slow* part driven by the turbulence alone. The
 83 slow part relaxes the Reynolds stress towards isotropy—the mechanism that the kernel
 84 (2.4)–(2.5) reproduces in model form. The rapid part is linear in A_{lm} and is determined by
 85 the velocity spectrum tensor $\Phi_{ij}(\kappa)$; the standard model represents neither $\mathcal{P}_{ij}$ nor $\Pi_{ij}^{(r)}$.

When the total strain accumulates in a time short compared with the eddy turnover time, the response is computed exactly by linear theory (Batchelor & Proudman 1954; Hunt 1973; Hunt & Carruthers 1990). Writing the fluctuation as Fourier modes that follow the mean flow, each mode of time-dependent wavevector $\kappa(t)$ and amplitude $\hat{u}_i$ evolves, in the inviscid rapid limit, according to

$$\frac{d\kappa_i}{dt} = -A_{ji} \kappa_j, \quad \frac{d\hat{u}_i}{dt} = \left(\frac{2 \kappa_i \kappa_m}{\kappa^2} - \delta_{im} \right) A_{mn} \hat{u}_n. \quad (2.8)$$

86 The bracketed operator combines the production $-A_{im} \hat{u}_m$ with the rapid pressure response,
 87 the solenoidal projection $P_{ij} = \delta_{ij} - \kappa_i \kappa_j / \kappa^2$ applied to the strained field; the factor of
 88 two is fixed by continuity, since only that coefficient makes $d(\kappa_i \hat{u}_i)/dt$ vanish under (2.8).
 89 Two features are central to what follows. First, the pressure response is *non-local* and
 90 intrinsically *three-dimensional*: it requires all of $\kappa_1, \kappa_2, \kappa_3$, whereas a single line resolves
 91 only the wavenumber along its own coordinate, so (2.8) cannot be evaluated on the line
 92 as written. Second, the rapid/slow decomposition furnishes the organising principle of the
 93 formulation: the slow part is already carried by the kernel, and it remains to supply the
 94 production and a rapid redistribution consistent with (2.8), expressed in quantities available
 95 on the line.

2.3. The strain-coupled line equation and its structural admissibility

The production term of the fluctuation momentum equation, $-A_{ij} u_j$, is a pointwise linear combination of the local velocity components and is therefore exactly representable on the line. Between eddy events the molecular evolution is replaced by

$$\partial_t u_i = \nu \partial_y^2 u_i - A_{ij} u_j + B_{ij} u_j, \quad (2.9)$$

97 in which $-A_{ij} u_j$ is the production and $B_{ij} u_j$ is the rapid pressure–strain operator
 98 constructed in § 2.4; the eddy events (2.1)–(2.5) are retained without modification, and the
 99 advancement is split between the continuous phase (2.9) and the discrete event sequence.

The forcing must not violate the structural properties on which the model rests, and it does not; we record the argument as a consistency check. The measure preservation, scale locality and kernel identities (2.3) are geometric properties of the map f and kernel K on a fixed domain, and the operator splitting applies the discrete event exactly as in the standard model, so they hold verbatim. Continuity of the fields is preserved because, for a mean gradient uniform along the line, the operator $\mathcal{A}_{ij} \equiv -A_{ij} + B_{ij}$ acts pointwise and the solution of the non-diffusive part of (2.9) over a sub-step is

$$u_i(y, t) = \left[\exp \int_{t_0}^t \mathcal{A} dt' \right]_{ij} u_j(y, t_0), \quad (2.10)$$

a y -independent matrix, which maps continuous profiles to continuous profiles while the molecular term is smoothing. The forcing displaces no fluid along y , so the Lebesgue measure of the domain is untouched; the deliberate straining of the domain is a separate operation (§ 2.5).

The energetic role of the forcing is exact. At fixed domain length the production term evolves the Reynolds stress as

$$\left. \frac{dR_{ij}}{dt} \right|_{\text{prod}} = -A_{ik}R_{kj} - A_{jk}R_{ki} \equiv \mathcal{P}_{ij}, \quad (2.11)$$

exactly the Reynolds-stress production tensor of (2.6): the term carries no model assumption. The kinetic-energy budget of the formulation thus separates into an exact production by the mean strain, energy-conserving redistribution (the kernels), and dissipation by molecular diffusion—the structure of the exact transport equation. The forcing adds the one term the standard model lacks without disturbing the terms it already represents.

2.4. The rapid redistribution operator

Were the continuous evolution to consist of production and diffusion only ($B_{ij} = 0$), the line would reproduce $dR_{ij}/dt = \mathcal{P}_{ij}$, whereas the exact inviscid rapid-distortion balance is $dR_{ij}/dt = \mathcal{P}_{ij} + \Pi_{ij}^{(r)}$. The omission is not small: for turbulence isotropic at the onset of straining the classical result derived below gives $\Pi_{ij}^{(r)} = -\frac{3}{5}\mathcal{P}_{ij}$, so production acting alone overstates the initial distortion rate by a factor of $\frac{5}{2}$; under sustained strain it also lets a component energy grow without the bound that incompressibility imposes. Representing $\Pi_{ij}^{(r)}$ is essential, not optional.

2.4.1. The exact rapid pressure–strain and the closure problem

The exact rapid term follows from the amplitude equation. Writing (2.8) as $d\hat{u}_i/dt = M_{in}\hat{u}_n$ with $M_{in} = -A_{in} + 2(\kappa_i\kappa_m/\kappa^2)A_{mn}$, and noting that the incompressible wavevector dynamics preserve the measure $d\kappa$ (their phase-space divergence is $-A_{ii} = 0$), the spectrum tensor $\Phi_{ij} = \langle \hat{u}_i^* \hat{u}_j \rangle$ yields

$$\frac{dR_{ij}}{dt} = \int (M_{in}\Phi_{nj} + M_{jn}\Phi_{in}) d\kappa = \mathcal{P}_{ij} + \Pi_{ij}^{(r)}, \quad (2.12)$$

in which the production separates from the pressure contribution,

$$\Pi_{ij}^{(r)} = 2A_{mn}(\mathcal{R}_{ijmn} + \mathcal{R}_{jimn}), \quad \mathcal{R}_{ijmn} \equiv \int \frac{\kappa_i\kappa_m}{\kappa^2} \Phi_{nj}(\kappa) d\kappa. \quad (2.13)$$

The tensor is traceless—by solenoidality $\kappa_n\Phi_{nj} = 0$, the trace integrand contains $\kappa_i\Phi_{ni} = 0$ —so the rapid term redistributes energy among components without changing the total, in keeping with the energy-conserving character of the kernel mechanism. Equation (2.13) also exposes the central difficulty: the rapid term is a functional of the spectrum tensor through the orientation-weighted integral $\mathcal{R}_{ijmn}$, not of the single-point Reynolds stress—the classical closure problem of the rapid pressure–strain, common to every single-point model (Crow 1968; Launder *et al.* 1975; Pope 2000) and intrinsic to the physics. A line carries R_{ij} and its own one-dimensional spectra, but not the distribution of energy over wavevector *orientation* that $\mathcal{R}_{ijmn}$ requires; a spectral-tensor closure is therefore unavoidable, and we make it explicit. (What a line’s data *do* determine about this integral, and how sharply, is the analytical question taken up in § 5.)

129 2.4.2. *Baseline isotropic closure*

The minimal closure takes the spectrum tensor instantaneously isotropic, $\Phi_{nj} = \frac{E(\kappa)}{4\pi\kappa^2} (\delta_{nj} - \kappa_n \kappa_j / \kappa^2)$ with $\int E d\kappa = k_t$. With the angular averages $\overline{e_i e_m} = \frac{1}{3} \delta_{im}$ and $\overline{e_i e_m e_n e_j} = \frac{1}{15} (\delta_{im} \delta_{nj} + \delta_{in} \delta_{mj} + \delta_{ij} \delta_{mn})$ for $e_i = \kappa_i / \kappa$,

$$\mathcal{R}_{ijmn}^{\text{iso}} = k_t \left[\frac{4}{15} \delta_{im} \delta_{nj} - \frac{1}{15} (\delta_{in} \delta_{mj} + \delta_{ij} \delta_{mn}) \right], \quad (2.14)$$

and substitution into (2.13) gives, with $S_{ij} = \frac{1}{2} (A_{ij} + A_{ji})$ and $A_{nn} = 0$,

$$\Pi_{ij}^{(r), \text{iso}} = \frac{4}{5} k_t S_{ij} = -\frac{3}{5} \mathcal{P}_{ij} \quad (\text{isotropic } R_{ij}), \quad (2.15)$$

130 the exact response of initially isotropic turbulence to a weak rapid strain (Crow 1968) and
 131 the content of the isotropisation-of-production model with $C_2 = \frac{3}{5}$ (Pope 2000)—obtained
 132 here, with no adjustable constant, as the isotropic limit of (2.13). For anisotropic turbulence
 133 $\mathcal{R}_{ijmn}$ must be modelled as a tensor function of R_{ij} ; the general representation linear in
 134 the anisotropy is that of Launder *et al.* (1975), to which the construction reduces in form.
 135 We adopt (2.14) as the baseline and assess the anisotropic correction in § 3.

 136 2.4.3. *Realisation on the line*

Given the closed rapid term—symmetric and traceless—we require a pointwise operator B_{ij} whose action on the line reproduces it. Demanding that the operator evolve the Reynolds stress by exactly $\Pi_{ij}^{(r)}$,

$$(\text{BR} + \text{RB})_{ij} = \Pi_{ij}^{(r)}, \quad (2.16)$$

a continuous Lyapunov equation with, for positive-definite $\mathbf{R}$, a unique symmetric solution $\mathbf{B}$. The solution conserves kinetic energy automatically—contracting (2.16) gives $2B_{ij}R_{ij} = \Pi_{ii}^{(r)} = 0$, so the rate of working of the operator on the line vanishes. In the isotropic case $\mathbf{R} = \frac{2}{3} k_t \mathbf{I}$ it reduces to the closed form

$$B_{ij} = \frac{3}{5} S_{ij}, \quad \mathcal{A}_{ij} = -A_{ij} + \frac{3}{5} S_{ij}. \quad (2.17)$$

137 B_{ij} is evaluated from the running Reynolds stress and applied continuously, paced by the
 138 mean strain, exactly as the production is. This is deliberate: the rapid pressure–strain is by
 139 construction the part of the redistribution driven by the mean gradient and therefore acts
 140 at the strain rate, in contrast to the slow redistribution that the eddy kernel supplies at the
 141 eddy rate. Strain-driven effects (production and rapid pressure–strain) are continuous and
 142 enter through $\mathcal{A}_{ij}$; turbulence-driven effects (the inertial cascade and the slow return to
 143 isotropy) remain the discrete events.

144 The sense in which the formulation is rapid-distortion consistent can now be stated
 145 precisely, again as a consistency check rather than a theorem. By (2.11) the production
 146 term contributes $\mathcal{P}_{ij}$ to dR_{ij}/dt and by construction (2.16) the operator contributes $\Pi_{ij}^{(r)}$;
 147 their sum is the exact balance (2.12) for the prescribed closure, with no adjustable constant,
 148 recovering (2.15) at the onset of distortion of isotropic turbulence. The scope of the
 149 statement must be drawn as carefully as the result: the consistency holds at the level of
 150 the *component energies* and is *conditional on the spectral closure*; the isotropic closure is
 151 exact only at onset, and its accuracy for anisotropic states is that of the isotropisation-of-
 152 production approximation it reproduces. The *spectral* distribution of the distorted energy is
 153 not enforced by (2.9): the operator acts uniformly on every resolved wavenumber, and the
 154 distribution of energy across scales emerges from its interaction with diffusion, the cascade

155 and the domain straining. It is therefore a genuine prediction, tested in §§ 3–4—where the
 156 uniform, wavenumber-blind action of B_{ij} is also identified as the model’s measured spectral
 157 deficiency (§ 4.2.1).

158 2.5. Wavevector kinematics by domain straining

The second ingredient of (2.8), the wavevector evolution, carries the kinematic compression and stretching of scales and is independent of the amplitude evolution: in the method-of-characteristics reading of linear theory, the wavevectors are the characteristics along which the amplitudes evolve. On the line the relevant wavenumber is κ_2 . We restrict attention to a line aligned with a principal axis of the mean strain, so that $A_{12} = A_{32} = 0$ and a material element of the line remains a coordinate line—as holds for the irrotational mean flow of a stagnation region. Then $d\kappa_2/dt = -A_{22} \kappa_2$, and since a material element of length ℓ stretches as $\dot{\ell}/\ell = A_{22}$ while wavenumbers scale as $1/\ell$, a uniform dilatation of the domain reproduces the wavevector kinematics exactly:

$$\frac{1}{L} \frac{dL}{dt} = A_{22}, \quad \implies \quad \kappa_2(t) = \kappa_2(t_0) \exp\left(-\int_{t_0}^t A_{22} dt'\right), \quad (2.18)$$

159 with L the domain length, implemented within the Lagrangian adaptive-mesh advancement
 160 of the model (Lignell *et al.* 2013) by imposing the mean-flow stretching on the mesh.

161 This step is not redundant with the amplitude operator. The Reynolds stress is an intensive
 162 second moment and is invariant under the uniform dilatation: the dilatation relabels the
 163 scales that carry the energy without changing the component energies, while $\mathcal{A}_{ij}$ changes
 164 the component energies without relabelling scales. The two operations are the physical-
 165 space counterparts of the characteristic motion and the amplitude evolution in (2.8), and
 166 together they account for the mean straining without double counting. Two limitations
 167 are inherent to representing a three-dimensional kinematics on one axis: the transverse
 168 wavenumbers also strain but are not represented, their effect being subsumed in the spectral
 169 closure, which already supplies the orientation information the line lacks; and a rotational
 170 mean flow, or a line not aligned with a principal axis, tilts the material line out of the
 171 coordinate direction and lies outside the formulation.

172 2.6. Summary of the formulation

173 The strain-coupled model advances the line field over a sub-step by composing (a) a
 174 *continuous strain–diffusion phase on a dilating domain*—integrate $\partial_i u_i = \nu \partial_y^2 u_i + \mathcal{A}_{ij} u_j$
 175 with $\mathcal{A}_{ij} = -A_{ij} + B_{ij}$, the production exact and B_{ij} given by (2.16) from the running
 176 Reynolds stress and the spectral closure (baseline (2.17)), while dilating the domain by
 177 (2.18)—and (b) *discrete eddy events* sampled from the standard rate distribution: the
 178 triplet map and the slow kernel, unchanged. Step (a) carries the mean-strain physics; step
 179 (b) carries the turbulence-driven physics; standard ODT is recovered exactly when $A_{ij} = 0$.

The relative importance of the two steps is governed by the ratio of the mean strain rate $S \equiv (2S_{ij}S_{ij})^{1/2}$ to the turbulence relaxation rate,

$$\chi \equiv \frac{S k_t}{\varepsilon}. \quad (2.19)$$

180 For $\chi \gg 1$ the distortion accumulates faster than the turbulence can respond, the eddy
 181 events are negligible over the distortion time, and the formulation reduces to the rapid-
 182 distortion limit of step (a), exact for the component energies under the closure. For $\chi \ll 1$
 183 the turbulence relaxes between increments of strain and the slow kernel dominates. The
 184 stagnation region of a leading edge is the intermediate regime $\chi = O(1)$, in which neither

limit is uniformly valid and both mechanisms contribute; the fidelity of the formulation across this range, and of the emergent distorted spectrum, are assessed against rapid-distortion solutions and scale-resolving data in §§ 3–4.

3. Verification, and moment-level validation

We separate two questions. *Verification* asks whether the formulation and its implementation reproduce a known exact solution in the regime where one exists—the rapid-distortion limit, tested analytically and in the solver below. *Validation* asks whether the model reproduces the behaviour of real strained turbulence outside that limit; the spectral validation, against the strained-box DNS with its exact-RDT companion, is the subject of § 4, and this section closes with the moment-level validation against the strained-turbulence DNS of Lee & Reynolds (1985).

3.1. Canonical strains and the exact onset rate

Two irrotational homogeneous strains admit exact rapid-distortion solutions and isolate the physics of interest. *Plane strain*,

$$\mathbf{A} = \text{diag}(a, -a, 0), \quad a > 0, \quad (3.1)$$

is the strain of a two-dimensional stagnation region: with x_1 tangential to the surface, x_2 normal to it and x_3 spanwise, the component compressed by the strain is the wall-normal (upwash) component u_2 , whose amplification is the pre-impact distortion relevant to leading-edge noise. *Axisymmetric strain*,

$$\mathbf{A} = \text{diag}(-\tfrac{1}{2}\gamma, -\tfrac{1}{2}\gamma, \gamma), \quad \gamma > 0, \quad (3.2)$$

is the strain of a contraction or a three-dimensional stagnation flow and provides an independent test in which two components are amplified and one suppressed. For both, $\mathbf{A}$ is symmetric, so the principal-axis alignment assumed in § 2.5 holds; the line is taken along x_2 .

For turbulence isotropic at the onset of straining the continuous evolution (2.9) gives, by (2.11) and (2.15),

$$\left. \frac{dR_{ij}}{dt} \right|_0 = \mathcal{P}_{ij} + \Pi_{ij}^{(r),\text{iso}} = \tfrac{2}{5} \mathcal{P}_{ij} = -\tfrac{8}{15} k_t S_{ij}, \quad (3.3)$$

the exact rapid-distortion rate for initially isotropic turbulence, with no adjustable constant. This is the sharpest verification available: a closed-form prediction the model must satisfy identically, not approximately. Evaluated for the two canonical strains it gives the component rates of table 1; for plane strain the predicted upwash growth rate $+\tfrac{8}{15}k_t a$ is to be compared with the $\tfrac{4}{3}k_t a$ that production alone would give—the factor- $\tfrac{5}{2}$ overstatement anticipated in § 2.4. A model that passes (3.3) for both strains has its production and rapid terms verified jointly; failure would indicate an error in the operator $\mathcal{A}_{ij}$ or its implementation, isolated from any closure or cascade effect.

3.2. Finite total strain, and the necessity of the rapid term

Beyond onset the turbulence becomes anisotropic and the rapid term (2.13) ceases to be a function of the Reynolds stress alone, so the verification becomes a test of the spectral closure rather than of the construction. We integrate the model moment equation along each canonical strain with two closures: the baseline isotropic closure (2.14) (equivalently,

	$dR_{11}/dt _0$	$dR_{22}/dt _0$	$dR_{33}/dt _0$
Plane strain (3.1)	$-\frac{8}{15}k_t a$	$+\frac{8}{15}k_t a$	0
Axisymmetric (3.2)	$+\frac{4}{15}k_t \gamma$	$+\frac{4}{15}k_t \gamma$	$-\frac{8}{15}k_t \gamma$

Table 1. Exact onset rates of the component energies from (3.3) for initially isotropic turbulence. The component compressed by the strain is amplified; the stretched component is suppressed.

isotropisation of production with $C_2 = 3/5$; IP) and the quasi-isotropic model of [Launder et al. \(1975\)](#) (LRR),

$$\Pi_{ij}^{(r)} = C_2 k_t S_{ij} + C_3 k_t (b_{ik} S_{jk} + b_{jk} S_{ik} - \frac{2}{3} b_{mn} S_{mn} \delta_{ij}) + C_4 k_t (b_{ik} W_{jk} + b_{jk} W_{ik}), \quad (3.4)$$

with anisotropy $b_{ij} = R_{ij}/2k_t - \frac{1}{3}\delta_{ij}$, mean rotation $W_{ij} = \frac{1}{2}(A_{ij} - A_{ji})$ (zero for the irrotational strains considered), and $C_2 = 4/5$, $C_3 = 7/4$, $C_4 = 131/100$; the value $C_2 = 4/5$ is fixed by requiring that (3.4) reduce at isotropy to the exact $\frac{4}{5}k_t S_{ij}$ of (2.15), so LRR shares the constant-free onset slope. The benchmark is the exact rapid-distortion solution, obtained by integrating the spectral equations (2.8) over a Gauss–Legendre ensemble of initial wavevector directions on the unit sphere: for initially isotropic turbulence the modal dynamics depend only on direction, so the Reynolds stress is the spherical average of the evolved modal covariance. This integration is independent of the moment models and reproduces the onset slope (3.3) to within 10^{-16} , verifying the rapid-distortion machinery itself.

Figure 1 reports the component-energy fractions against total strain. Three observations follow. First, at onset all curves for each component are tangent—the constant-free identity (3.3) is satisfied—and through $e \lesssim 1$ the closures are nearly indistinguishable from the exact solution. Second, as anisotropy accumulates the closures depart, LRR uniformly less than IP: for axisymmetric strain at $e = 4$ the lateral fraction is 0.498 (exact), 0.468 (LRR), 0.450 (IP), and the axial fraction 0.003, 0.063, 0.100. This gap is the spectral-closure error in isolation—the eddy events are inactive in the rapid limit—and it bounds the accuracy attainable from any single rapid operator. Third, and most important to state plainly, for plane strain both closures miss a *qualitative* feature: in the exact solution the neutral spanwise component grows monotonically, overtakes the upwash near $e \approx 3.3$, and is the largest component by $e = 4$ ($\overline{u_3^2}/2k_t = 0.51$ against 0.47), whereas both closures predict it to decay. This redistribution into the neutral direction is governed by the orientation distribution of the spectral tensor and cannot be recovered from single-point quantities; it is the moment-level signature of precisely the transverse-wavenumber information a single line does not carry. Section 5.6 returns to it with the strained-box DNS in hand and localises the deficiency in the closure’s b -dependent term.

The upwash component, whose spectrum is the input to the leading-edge response, makes the case for the rapid term quantitative (figure 2). Production acting alone drives the upwash fraction from $1/3$ to 0.98 by $e = 4$ —almost the entire kinetic energy forced into one component, an essentially non-realisable state—and overstates the initial amplification rate by the factor $\frac{5}{2}$. The rapid term removes three fifths of this at onset and keeps the fraction physical: the exact solution peaks at 0.49 near $e \approx 2.4$ and then relaxes as energy passes to the spanwise component. Both closures reproduce the amplification itself—the




Figure 1. Component energy fractions against total strain for initially isotropic turbulence under (a) plane strain and (b) axisymmetric strain, normalised to $S = 1$: exact rapid-distortion theory (wavevector-ensemble integration of (2.8)) against the model moment equation with the LRR closure (3.4) and the IP closure (2.14). For axisymmetric strain the two lateral components coincide and only $\overline{u_1^2} = \overline{u_2^2}$ (lateral) and $\overline{u_3^2}$ (axial) are shown. All closures share the exact onset slope; LRR tracks the exact solution more closely at finite strain.




Figure 2. Upwash energy fraction under plane strain for initially isotropic turbulence: exact rapid-distortion theory, the model with the LRR and IP closures, and production alone ($B_{ij} = 0$). Production without the rapid term overstates the onset amplification rate by $\frac{5}{2}$ and forces almost all the energy into the upwash component; the rapid term restores the physical, bounded amplification.

effect that the frozen, isotropic input of classical leading-edge-noise prediction omits— while overpredicting it at large strain (LRR 0.60, IP 0.65 at $e = 4$, against the exact 0.47). The rapid pressure–strain is therefore not a refinement but a necessity, and the accuracy of the method for a given configuration is controlled by the total strain accumulated along the stagnation streamline.

3.3. Verification of the implementation

The preceding checks verify the *formulation*. They do not test the *implementation*, in which the operator is not prescribed but reconstructed at every substep: the line Reynolds




Figure 3. Verification of the strain-coupled implementation under plane strain, eddy events suppressed and zero molecular diffusion. (a) Component energy fractions; (b) kinetic-energy amplification $k_t(e)/k_t(0)$. Symbols: solver, length-weighted line averages. Dashed: the LRR closure integrated as a moment equation. Solid: exact rapid-distortion theory. The solver reproduces the closure to 10^{-4} in the fractions and 0.3% in the energy; the offset from the exact solution is the closure error of § 3.2, as it must be.

stress is formed from the resolved field, the closure supplies $\Pi_{ij}^{(r)}$, the Lyapunov equation (2.16) is solved for B_{ij} , and $\mathcal{A}_{ij} = -A_{ij} + B_{ij}$ is applied pointwise. To isolate this chain the solver is run in the rapid-distortion limit—plane strain, eddy events suppressed, zero molecular diffusion—from a statistically isotropic resolved initial field, and the component energies are evaluated as length-weighted line averages. With the events inactive the line’s component-energy evolution must coincide with the closed moment equation across the whole strain range, not merely at onset.

Figure 3 shows that it does: the solver output lies on the LRR trajectory over $0 \leq e \leq 4$ to within 10^{-4} in each component fraction, so the per-substep stress reconstruction, Lyapunov solve and operator assembly are correct at every state of anisotropy. The comparison must be read correctly: the solver reproduces the *closure*, and it should—with the events suppressed the line carries no information beyond the single-point statistics the closure itself uses. The offset from the exact solution is the single-point closure error of § 3.2, not an implementation error; agreement with the exact solution here would indicate a fault. The fractions test only the anisotropy; the kinetic energy, which the traceless rapid term cannot change directly and which grows through the production integral $d \ln k_t / de = f_{22} - f_{11}$, provides the independent magnitude check, and the solver reproduces the LRR amplification to within 0.3%. Reaching that figure required a resolved spectral initialisation (a white-noise start places energy at the grid scale, where mesh-adaption interpolation damps it) and a strain-resolved timestep bound in the inviscid limit; both effects act nearly isotropically, so the energy panel is what exposes them.

3.4. Moment-level validation against the strained-turbulence DNS of Lee & Reynolds

The DNS of Lee & Reynolds (1985) is retained as a benchmark for one specific reason: it was deliberately run at a turbulence Reynolds number low enough ($q^4/\nu\epsilon \lesssim 100$) to be fully resolvable on a single line, and it reports the Reynolds-stress anisotropy against total strain for several irrotational strain modes alongside the analytic rapid-distortion prediction—a graded, moment-level reference that the spectral benchmarks of § 4 do not provide. The comparison quantity is the anisotropy b_{ij} against the deformation ratio $c = \exp \int S dt$ (so $\ln c$ is the total strain e of the present notation), computed from the single-point component




Figure 4. Reynolds-stress anisotropy under plane strain: present model (lines; ensemble mean over 1000 realisations, standard-error band) against the DNS of [Lee & Reynolds \(1985\)](#) (symbols, digitised from their figure 5.4). The strained components b_{22} and b_{33} are reproduced quantitatively; the unstrained b_{11} is strain-rate-dependent in the DNS (shaded band), with the model's near-zero value at the slow-strain edge.

variances along the line, ensemble-averaged over 1000 realisations; no spectral transform is involved. Two of their strain modes are run: plane strain and axisymmetric contraction, the line along a compressed direction in each case, to the total strains of the DNS ($c = 4$ and 3.32); the measured line compression agrees with $\exp(A_{22}t_f)$ to better than 0.3%, confirming the kinematic fidelity of the dilatation. One parameter distinction matters for interpretation: [Lee & Reynolds](#) characterise each run by $S^* = Sq^2/\varepsilon = 2\chi$, and find the axisymmetric anisotropy essentially independent of S^* —a clean, parameter-free target—whereas in plane strain the unstrained component b_{11} is strongly S^* -dependent, spanning $b_{11} \approx 0$ (slow) to +0.12 (rapid) in the DNS.

Under plane strain (figure 4) the model reproduces the two strained components quantitatively across the whole range: $b_{22} = +0.21$ at $c = 3.86$ against $\approx +0.20$ in the DNS, and $b_{33} = -0.20$ against ≈ -0.26 , capturing the trend and the bulk of the magnitude. These are exactly the components the DNS identifies as strain-rate-independent, so they constitute the robust part of the comparison. Under axisymmetric contraction (figure 5) the model reproduces the sign structure and the monotonic approach to a strained state while under-predicting the anisotropy magnitude by roughly 25% ($b_{11} = -0.21$ against ≈ -0.29 at $c = 3.16$); the dominant contribution is the spectral-closure gap isolated in § 3.2, common to the whole second-moment family, with a small additional artefact specific to the line: lying along one of the two equivalent transverse directions, it breaks the $x_2 \leftrightarrow x_3$ symmetry and returns a residual split $b_{22} = +0.102$ against $b_{33} = +0.113$ at $c = 3.16$, which the symmetry-respecting mean $\frac{1}{2}(b_{22} + b_{33})$ removes.

The one component the model does not reproduce, the plane-strain unstrained anisotropy b_{11} , is not a deficiency specific to the model, and figure 6 locates it exactly. Exact linear rapid distortion, evaluated by the wavevector-ensemble integration of § 3.2, gives b_{11} rising to +0.114 at $c = 4$: although $A_{1k} = 0$, the rapid pressure-strain feeds the neutral direction—the same orientation-carried redistribution as the spanwise growth in § 3.2. (A tempting kinematic argument, that $\langle u_1^2 \rangle$ is conserved because the component is unstrained, holds for production but not for the rapid term, and gives the wrong sign; an earlier version of this comparison used it.) The fast-strain DNS therefore tracks exact RDT, as [Lee &](#)




Figure 5. Anisotropy under axisymmetric contraction: present model (lines; 1000 realisations) against [Lee & Reynolds \(1985\)](#) (symbols, digitised from their figure 5.8). The shaded spread is the residual $b_{22} \neq b_{33}$ split from the line's breaking of the transverse symmetry.

[Reynolds](#) themselves observed, while both single-point closures give $b_{11} \approx 0$, slightly negative: the closure family misses the neutral-direction feed, which is carried by the orientation distribution that single-point models discard—the same deficiency measured spectrally, and localised in the closure's b -dependent term, in § 5.6. The present model coincides with the closure on which its operator is built, so its near-zero b_{11} is correct for its formulation and matches the DNS at slow strain rate, where the nonlinear scrambling erases the neutral-direction feed; at fast strain it inherits the closure family's known limitation rather than any line-specific artefact. The components that govern the upwash spectrum fed to the leading-edge response are precisely those the model reproduces quantitatively. Finally, [Lee & Reynolds's](#) own transport-budget analysis finds that in plane strain the intercomponent transfer is carried almost entirely by the rapid term—independent support for the rapid/slow decomposition on which § 5 rests.

4. The linear reference and the distorted spectrum

4.0.1. Compression without cascade: the model's kinematic reference

With the eddy events suppressed and zero molecular viscosity, the mean strain is the only process acting on the resolved field, and its effect on the line is purely kinematic. The line length contracts as $\mathcal{L}(e) = \mathcal{L}_0 e^{A_{22}e}$, so every resolved wavenumber is rescaled as $k(e) = k(0) e^{-A_{22}e}$ —the wavevector kinematics it shares with rapid-distortion theory ([Batchelor & Proudman 1954](#); [Hunt & Carruthers 1990](#))—while the continuous redistribution operator of § 2.4 acts uniformly on every resolved wavenumber. The combination transfers no energy between scales: a rigid translation of the line spectrum toward higher wavenumber. We emphasise at the outset that this rigid translation is a property of the strain-coupled line—uniform dilatation composed with a wavenumber-uniform operator—and not of linear distortion theory itself; the correct linear reference for the projected spectrum is constructed in § 4.0.2 below. As a check on the implementation, however, the rigid translation is an exact, parameter-free target.




Figure 6. The unstrained-direction anisotropy b_{11} under plane strain: present model (solid, 1000 realisations, standard-error band) against exact linear rapid distortion (dotted, wavevector-ensemble integration), the IP and LRR-QI closures, and the DNS of Lee & Reynolds (1985) (symbols and shaded S^* range). The fast-strain DNS tracks exact RDT, whose rapid pressure–strain feeds the neutral direction; the single-point closures—and the model built on them—miss this orientation-carried redistribution, the deficiency measured and localised in § 5.6.

Figure 8 confirms it. The component spectra translate toward higher k without change of shape—the peak neither broadens nor develops an inertial range—and the spectral centroid follows $e^{-A_{22}e}$, reaching 7.03 at $e = 3.9$ against the exact 7.03, with the three components migrating identically; the domain length follows $e^{A_{22}e}$ to four digits. The single-point moments are unchanged from the no-dilatation case, since a uniform rescaling of position leaves the length-weighted velocity statistics invariant. Compression therefore acts entirely on the spectrum and reproduces the model’s kinematic prediction exactly. It is the kinematic reference (the dashed line in all centroid figures below) against which the full model is measured.

4.0.2. The linear reference for the projected spectrum: exact RDT

The rigid translation above must not be mistaken for the prediction of linear rapid-distortion theory for the quantity the line actually carries. In the full linear theory the amplitude of each Fourier mode evolves at a rate that depends on the orientation of its three-dimensional wavevector (equation 2.8); the one-dimensional component spectra are projections—integrals over the transverse wavenumber plane—and there is no reason for a projection of orientation-dependent amplification to translate rigidly. Whether it does is a question of calculation, and the answer is that it does not. Using the closed-form Cauchy solution of the linear problem applied to isotropic initial fields and projected onto a line (the exact-RDT companion calculations described in § 4.2), we define the shape ratio $R_{nn}(\kappa_2)$ as the projected component spectrum at strain e , normalised by its integral, divided by the rigidly translated initial spectrum normalised likewise; $R_{nn} \equiv 1$ at every κ_2 if and only if the translation is rigid. At $e = 1$ under plane strain the upwash ratio R_{22} falls monotonically from 1.60 at half the initial spectral centroid to 0.38 at twelve times it: exact linear theory redistributes the projected upwash spectrum toward low wavenumber




Figure 7. Exact linear theory does not translate the projected spectra rigidly. Left: shape ratio $R_{nn}(\kappa_2)$ of the exactly distorted, line-projected component spectra to a rigid translation, at $e = 1$ under plane strain (closed-form Cauchy solution, isotropic initial fields, four realisations). Right: the transverse splitting $\phi_{11}/\phi_{33} - 1$ versus wavenumber in exact linear theory and in the strained-box DNS of § 4.2.




Figure 8. Compression without cascade (eddy events suppressed, inviscid). (a) Component line spectra $E_i(\kappa_2)$ at total strains $e = 0, 1, 2, 3, 3.9$ (lighter to darker); the spectra translate to higher wavenumber without changing shape. (b) Spectral-centroid migration $\bar{\kappa}(e)/\bar{\kappa}(0)$ for each component (symbols) against the kinematic prediction $e^{-A_{22}e}$ (dashed). The centroid follows the prediction and the three components coincide, confirming purely geometric compression of the line.

by a factor of four across the resolved range, because modes aligned with the line ($\kappa \rightarrow \kappa_2$) are amplified least. The transverse components are affected more weakly (R_{11} from 1.27 to 0.75; R_{33} within ten per cent of unity), so the projected anisotropy of linear theory is itself scale-dependent. Figure 7 shows the ratios.

Two consequences run through the remainder of the paper. First, the comparison of the full model against the rigid translation (§ 4.0.3) measures the departure of the model from its own kinematics; the departure of the *flow* from linear theory must be measured against the exact projected reference, and the two differ already at the linear level. Second, the model's rigid spectral kinematics is itself a structural approximation—linear theory deepens the projected anisotropy with wavenumber, which a wavenumber-uniform operator cannot represent—and § 4.2.1 quantifies the resulting deficiency against both the exact reference and DNS.

4.0.3. The distorted spectrum: strain on against strain off

The central model-internal comparison isolates what the strain does to the spectrum at fixed Reynolds number, holding the turbulence otherwise identical. Two ensembles of 128 realisations are run at $\nu = 10^{-5}$: the full model (eddies and dilatation active) and a baseline identical in every respect except that the strain machinery is switched off, so the line does not compress. Because the two differ only in the presence of the mean strain, the difference between them is the distortion, cleanly separated from the cascade and dissipation the eddies produce on their own.

Figure 9 shows the result, and it departs from the kinematic reference in a specific and physically meaningful way. Consider the centroid first. The no-strain baseline, after the brief initialisation transient, relaxes monotonically to $\bar{\kappa}/\bar{\kappa}_0 < 1$: with no compression, the eddies merely redistribute energy and the centroid drifts below its initial value. The full model is lifted clearly above this baseline—compression does push the spectrum toward higher wavenumber—yet it settles at $\bar{\kappa}/\bar{\kappa}_0 \approx 1.6$ – 2.1 , far below the rigid-translation line, which rises to 7.4 at $e = 4$. The reason is visible in the spectra: the full-model spectrum is not a rigid translation of the baseline but a *broadband* distortion. It extends to substantially higher wavenumber than the baseline—energy is genuinely carried to small scales by compression and cascade—but the bulk of the energy remains at the energetic scales, and what reaches high wavenumber is dissipated there rather than accumulated.

Two distinct effects are mixed in this departure, and § 4.0.2 requires that they be separated. The first is the eddy cascade and dissipation, which the strain-on/strain-off construction isolates cleanly at the level of the model. The second is that exact linear theory itself, once projected onto the line, is already broadband (figure 7): part of what a rigid-translation reference attributes to nonlinearity is present in the linear physics. The strain-on/strain-off comparison therefore measures what the mean strain does *within the model*; how the model’s distorted spectrum stands against exact linear theory and against DNS on the same footing is answered by the three-way comparison of § 4.2.1. What survives either way is the qualitative point that a frozen, rigidly strained upwash spectrum misrepresents the distorted inflow, which motivates supplying a distorted spectrum to Amiet’s theory in place of it, in the spirit of the rapid-distortion-modified inflow spectra of [de Santana et al. \(2016\)](#) but without the frozen-shape assumption.

4.0.4. The physical operating point

The molecular viscosity is, in these units, the inverse integral-scale Reynolds number, and it is a physical property of the represented inflow, not a tunable parameter: a sweep over $\nu = 10^{-4}$ – 10^{-7} shows the cascade deepening monotonically, with the resolved cases settling below the linear-RDT centroid line while the lowest viscosities climb toward it only by piling energy into an under-resolved grid-scale shelf. The operating point is therefore the viscosity matching the target inflow, resolution-checked. For the reference configuration—a NACA0012 at $U_\infty = 20 \text{ m s}^{-1}$ in grid turbulence with intensity $u'/U_\infty \approx 6\%$ and integral scale $\mathcal{L} \approx 55 \text{ mm}$ at the leading-edge plane—the integral-scale Reynolds number is $Re_{\mathcal{L}} \approx 4.4 \times 10^3$, which maps through the initial-condition normalisation to $\nu \approx 3 \times 10^{-6}$, run on a refined grid (16000 cells) on which the dissipation rolloff is resolved with margin; the ensemble comprises 200 realisations.

Figure 10 shows the result. The component spectra fall smoothly across four decades with no grid-scale shelf, the upwash component E_2 amplified above the streamwise E_1 —the spectral signature of the plane strain—and no extended $k^{-5/3}$ range, as expected physically at this Reynolds number ([Kolmogorov 1991](#); [Pope 2000](#)). The centroid settles after the initialisation transient and migrates to ≈ 2.4 by $e = 4$, far below the rigid-translation value of 7.4 : the broadband, dissipative distortion of § 4.0.3 at the physical operating point. The




Figure 9. Full model against the matched no-strain baseline at $\nu = 10^{-5}$ (ensemble means over 128 realisations, geometric $\pm\sigma$ bands). (a) Final-strain total spectrum $E(\kappa_2)$: the full model (strain on) extends to higher wavenumber than the baseline (strain off) but remains broadband, not a rigid translation. (b) Centroid migration: the baseline relaxes below unity, the full model is lifted above it by the compression, yet stays far below the rigid-translation line (dashed). The gap between the two solid curves is the distortion attributable to the mean strain at this Reynolds number.

regime is quantified with an estimator-free dissipation: because the eddy events conserve energy and the production is exact, molecular dissipation is the only sink, so it is fixed by the resolved energy budget

$$\varepsilon = \mathcal{P} - \frac{dk_t}{dt} = -A_{ij}R_{ij} - \frac{dk_t}{dt}, \quad (4.1)$$

every term available on the line. By this budget the strain-rate parameter settles at $\chi = Sk_t/\varepsilon \approx 0.8$ over the strained trajectory, corroborated within a few tens of per cent by the direct gradient estimate $\langle 2\nu(\partial u_i/\partial y)^2 \rangle$: the operating point sits at $\chi = O(1)$, the regime the leading-edge stagnation flow occupies, where neither the rapid-distortion nor the near-equilibrium limit applies and where beating linear theory is meaningful. The spectral *peak* corroborates the regime reading independently of the dissipation estimator: the energy-containing wavenumber itself migrates far less than the rigid law, so the eddy relaxation holds back the energy-containing range, not merely the dissipation tail—linear theory misplaces the energy-containing wavenumber, not only the spectral shape at high κ . The cost is the practical justification of the approach: the operating-point ensemble completes in a few hundred core-hours, roughly two orders of magnitude below a strained DNS ensemble at comparable Reynolds number, which is what makes parametric sweeps over inflow condition feasible.

4.1. Validation against the strained-turbulence experiment of Chen, Meneveau & Katz

The strain-on/strain-off comparison is model-internal; an external benchmark must confirm that both the rapid-distortion limit and the departure from it are physical. We use the plane-strain experiment of [Chen et al. \(2006\)](#) (CMK), in which initially near-isotropic turbulence at $R_\lambda \approx 400$ is subjected to a controlled planar straining cycle and the one-dimensional component spectra are compared with rapid-distortion theory. We reproduce the *straining* phase of their cycle (the destraining phases involve a strain reversal outside the present monotonic configuration); its end corresponds to a total deformation $D = \exp(\frac{1}{2} \int S dt) \approx 3$, i.e. $e \approx 2.20$ in the present notation.




Figure 10. The physical operating point ($\nu \approx 3 \times 10^{-6}$, $Re_{\mathcal{L}} \approx 4.4 \times 10^3$, 16000 cells; ensemble over 200 realisations, geometric $\pm\sigma$). (a) Final-strain component spectra: smooth rolloff with no grid-scale shelf; the upwash E_2 amplified above the streamwise E_1 ; dotted reference slope $\kappa^{-5/3}$. (b) Total-spectrum centroid migration against the rigid-translation law (dashed): the broadband distortion that linear theory cannot represent, at the operating point and with the noise averaged out.




Figure 11. Rapid-distortion limit (eddy events suppressed, inviscid) against the analytic reference of [Lee et al. \(1986\)](#) used by [Chen et al. \(2006\)](#). (a) Component spectral-centroid migration follows the rigid-translation law, reaching the deformation ratio $D = 3$ at the end of the straining phase. (b) Kinetic-energy amplification: the solver follows the closure trajectory, within 2% of the exact rapid-distortion value at $e = 2.20$.

In the rapid-distortion limit (eddy events suppressed, inviscid) the model must reproduce the analytic reference of [Lee et al. \(1986\)](#) that CMK use. Figure 11 shows that it does: the component spectral centroids follow the rigid-translation law, reaching the deformation ratio 3.00 at $e = 2.20$ with the components coincident, and the kinetic energy reaches 1.72 against the exact rapid-distortion amplification 1.69—the residual being the moment closure’s, not the implementation’s (§ 3.3), since with the events suppressed the solver evolves the closure trajectory (1.72) exactly. The truly inviscid setting is required: any nonzero viscosity dissipates the compressed small scales and degrades the rigid translation, an effect we verified directly.

With the eddy events and viscosity active the model is no longer constrained to rigid translation, and the strain drives a measurable anisotropy between the components.




Figure 12. Full model (eddies and viscosity active; ensembles over 128 realisations at $\nu = 10^{-5}$): strain-induced migration of the streamwise and upwash component centroids, each as the ratio of the strained ensemble to the matched no-strain baseline, against the rigid-translation law (dashed). The components diverge under strain—the upwash held back most—reproducing the qualitative departure-from-RDT mechanism of [Chen *et al.* \(2006\)](#). The comparison is qualitative; see text.

Figure 12 shows the strain-induced migration of the streamwise and upwash component centroids, each measured against the matched no-strain baseline (the per-component form of the strain-on/strain-off construction, which cancels the initialisation transient and the eddies' own relaxation): the components diverge progressively under continued strain—the streamwise migration reaching 1.9 at the end of the straining phase against 1.5 for the upwash, both far below the rigid value 3—so the upwash component is held back most. This is the qualitative behaviour CMK report—the strained spectrum departs from the isotropic-translation expectation with the components responding differently—and it is the spectral signature of the broadband distortion of § 4.0.3. The comparison is restricted to its qualitative content, for three reasons stated plainly: the band-limited initial spectrum differs from CMK's, so absolute levels are not comparable; the line-resolved one-dimensional spectra and the experiment's transverse/longitudinal projections are not the same projection of the spectrum tensor; and the scale separation achieved here ($\mathcal{L}/\eta \approx 250$) is below the experimental ≈ 930 , whose compressed dissipation range exceeds feasible line resolution. Within these limits the model reproduces the rapid-distortion reference and the qualitative departure mechanism of the benchmark; the quantitative spectral assessment is carried out against the strained-box DNS below, where the projection and the initial spectrum are matched by construction.

4.2. Validation against a strained-box DNS with an exact-RDT companion

The comparisons above are either model-internal (§ 4.0.3) or, where external, qualitative by construction (§ 4.1) or at the Reynolds-stress level (§ 3.4). The referent quantity for the leading-edge application is spectral, and the objection that the emergent-spectrum results compare the model only with itself must be met with an independent, projection-matched spectral benchmark under the same strain. To that end we performed direct numerical simulations of plane-strained homogeneous turbulence in a deforming-frame (Rogallo-type) pseudo-spectral formulation: a 128^3 box, decaying isotropic precursor, then plane strain $A = S \text{diag}(\frac{1}{2}, -\frac{1}{2}, 0)$ applied at strain-rate parameters $Sk_t/\varepsilon \in \{0.8, 16\}$ at onset, carried to total strain $e = 1$, four independent realisations ($\kappa_{\max}\eta \approx 0.76$ at $e = 1$: a pilot resolution, so small-scale statements below are qualitative). Alongside every DNS

$\Delta b_{22}(\kappa_2)$, band κ_2/κ_c :	0.5–0.75	0.75–1.2	1.2–1.9	1.9–3	3–4.8	4.8–7.6	7.6–12
exact projected RDT	+0.146	+0.008	–0.020	–0.041	–0.050	–0.054	–0.056
DNS	+0.032	–0.033	–0.029	–0.032	–0.036	–0.040	–0.046
strain-coupled ODT	+0.100	+0.101	+0.114	+0.111	+0.127	+0.099	+0.147

Table 2. Strain-induced upwash anisotropy per wavenumber band at $e = 1$, $Sk_t/\varepsilon = 0.8$, each system referenced to its own unstrained state. Bands below $0.75 \kappa_c$ rest on a few box modes and are indicative only.

realisation runs an *exact-RDT companion*: the same initial field evolved under the closed-form Cauchy solution of the linearised equations and projected onto lines identically to the DNS and to ODT. The companion is what § 4.0.2 used to construct the linear reference; here it completes a three-way comparison in which model, linear theory and DNS are reduced to the same one-dimensional observables. On the ODT side the ensembles use an unstrained precursor: the line is evolved as ordinary ODT until the initialisation transient has relaxed and the eddy population is statistically active, and the strain is switched on only then, so that the strained evolution starts from a physically conditioned state rather than from the synthetic initial condition. All ODT statistics in this subsection are medians over 1024 realisations per case with bootstrap confidence intervals; median statistics are used because the strained band energies are strongly intermittent across realisations, and ensemble means of spectral ratios are dominated by rare events at any practical sample size.

4.2.1. Three-way spectral anisotropy: ODT, exact RDT, DNS

The comparison quantity is the strain-induced change of the spectral component anisotropy, $\Delta b_{nn}(\kappa_2) = \phi_{nn}/\sum_m \phi_{mm} - \frac{1}{3}$ at strain e , minus the same quantity evaluated in the system's own unstrained state with each mode mapped to its strained wavenumber. Referencing each system to its own $e = 0$ state cancels viscous decay, which is component-blind, and the longitudinal–transverse structure of the isotropic line spectra, which differs between a three-dimensional box and the component-symmetric ODT line. Figure 13 and table 2 give the result at $e = 1$, $Sk_t/\varepsilon = 0.8$, on common bands of $\kappa_2(e)/\kappa_c(0)$ with κ_c the initial spectral centroid.

The structure of the table is the finding. Exact linear theory concentrates the strain-induced upwash anisotropy at the energy-containing scales and *removes* it from the small scales, because modes aligned with the line are amplified least; the DNS shows the same shape, weakened at the large scales by the nonlinear return. The strain-coupled ODT distributes its moment-level anisotropy essentially uniformly over wavenumber—the signature, anticipated in § 4.0.2, of a wavenumber-uniform redistribution operator acting on a uniformly dilated line. The integrated anisotropy is nevertheless correct (the model tracks the closure at the moment level, § 3.4), so the deficiency is one of allocation across scales, not of amount. For the leading-edge application the operative consequence is that the high-wavenumber anisotropy of the upwash spectrum, which the Amiet response weights, has the wrong sign in the present model; the closure element that must carry the correction—a wavenumber-dependent rapid operator in place of the uniform one—is exactly the spectral kernel formulation of § 5, whose exercise is thereby motivated by a measured deficit rather than by completeness.

4.2.2. Distance from linear theory as a function of strain rapidity

The three-way comparison resolves the anisotropy by wavenumber at one strain-rate parameter. A complementary scalar measure tracks the *whole* spectral shape across strain




Figure 13. Three-way comparison at $Sk_t/\varepsilon = 0.8$: strain-induced anisotropy of the line spectra relative to each system's own unstrained state; strain-coupled ODT (1024 realisations) against exact projected RDT and DNS (128³, four realisations); $e = 1$ solid, $e = 0.5$ dashed.

and strain rapidity: at each total strain the median ODT component line spectra are fitted, jointly in the upwash and transverse components, by the exactly distorted spectrum of linear theory—the closed-form Cauchy distortion of an isotropic von Kármán field, projected onto the line, with only the pre-distortion amplitude and energy wavenumber free. The root-mean-square log-residual of that two-parameter fit is a scale-resolved distance between the model's strained spectra and exact linear theory. Figure 14 shows it for precursor-conditioned ensembles at $Sk_t/\varepsilon \approx 0.4$ and ≈ 8 at onset (1024 realisations each, identical precursor state: the two fits coincide at $e = 0$ at thirteen per cent, the residual shape mismatch of a relaxed ODT line against the von Kármán form).

The distance grows with strain in both cases, but *faster at rapid strain*: to approximately 22 per cent at $e = 2$ for $Sk_t/\varepsilon \approx 0.4$ against approximately 24 per cent, saturating early, at $Sk_t/\varepsilon \approx 8$, with the separation established already by $e = 0.5$. If the model's rapid response were spectrally faithful, the rapid curve is the one that should approach zero: eddy events are subdominant there by construction. That the deviation is largest precisely where the eddies matter least localises the deficiency in the rapid kinematics themselves—the wavenumber-uniform operator and the rigid dilatation—and not in the eddy or relaxation statistics, in quantitative agreement with the three-way comparison. This measured validity envelope accompanies the model into any application: predictions inherit a spectral-shape uncertainty of order the plotted distance at the operating point's strain and rapidity.

5. Closure of the rapid pressure–strain term on a single line

The results of § 4 establish what the strain-coupled model produces and where it stands against exact linear theory and DNS. The rapid pressure–strain redistribution that drives that evolution was evaluated throughout by the algebraic moment closure of § 2 (Launder *et al.* 1975), applied component by component on the resolved line. That closure is a modelling input, and the redistribution it represents is, in its exact form, a *nonlocal, three-dimensional* object: the rapid pressure at a point is a volume integral over the entire fluctuating field, and the pressure–strain correlation is an integral over the full three-dimensional spectrum. What a single one-dimensional line of turbulence can say about such an object is the analytical question of this paper, and this section answers it in a form that is deliberately




Figure 14. Scale-resolved distance between the strained ODT line spectra and exact linear theory: rms log-residual of the two-parameter exact-RDT-distorted von Kármán fit, against total strain, for slow and rapid onset strain-rate parameters (medians over 1024 realisations; identical precursor state at $e = 0$).

more modest, and more defensible, than a closure claim: we state exactly what obstructs a single-line representation, we derive sharp bounds on the rapid term from line data alone, we *measure* rather than assume the axisymmetry hypothesis under which the term collapses, and we identify—by measurement—which physical content the moment closure discards.

The section proceeds as follows. The exact term and the obstruction are stated first (§ 5.1–§ 5.2); the axisymmetric collapse follows, together with the statement of what it does *not* deliver (§ 5.3); the bounds that line data do deliver, with their falsifiers, come next (§ 5.4); the axisymmetry assumption is then measured directly in the strained-box DNS, including a control experiment that isolates its failure mode (§ 5.5); the component-ordering deficiency raised by the exact solution is measured and localised (§ 5.6); the reduction to the moment closure and the disjointness from the eddy redistribution close the argument (§ 5.7–§ 5.8).

5.1. The exact rapid pressure–strain term

The starting point is the Poisson equation (2.7) for the fluctuating pressure, whose source splits into a part linear in the fluctuations and the mean velocity gradient (the *rapid* part) and a part quadratic in the fluctuations (the *slow* part) (Pope 2000). The rapid part is linear in $\mathbf{u}'$ and proportional to the mean gradient, which for the present homogeneous plane strain is the constant tensor A_{kl} . Writing p'_r for the rapid pressure and inverting the Laplacian with the free-space Green’s function $G(\mathbf{x}) = -1/4\pi|\mathbf{x}|$,

$$p'_r(\mathbf{x}) = \frac{\rho}{2\pi} A_{kl} \int \frac{1}{|\mathbf{x} - \mathbf{x}'|} \frac{\partial u'_l}{\partial x'_k}(\mathbf{x}') d\mathbf{x}'. \quad (5.1)$$

The rapid pressure–strain correlation that redistributes energy among the Reynolds-stress components is $\Pi_{ij}^{(r)} = \overline{(p'_r/\rho) (\partial u'_i/\partial x_j + \partial u'_j/\partial x_i)}$. Substituting (5.1) and using homogeneity to pass to the velocity spectrum tensor $\Phi_{mn}(\boldsymbol{\kappa})$ (the Fourier transform of $u'_m(\mathbf{x}) u'_n(\mathbf{x} + \mathbf{r})$), one obtains the standard exact result (Batchelor & Proudman 1954;

Crow 1968; Hunt & Carruthers 1990)

$$\Pi_{ij}^{(r)} = A_{kl} M_{ijkl}, \quad M_{ijkl} = \int_{\mathbb{R}^3} \left[\frac{\kappa_j \kappa_k}{\kappa^2} \Phi_{il}(\boldsymbol{\kappa}) + \frac{\kappa_i \kappa_k}{\kappa^2} \Phi_{jl}(\boldsymbol{\kappa}) \right] d\boldsymbol{\kappa}, \quad (5.2)$$

where $\kappa = |\boldsymbol{\kappa}|$. Equations (2.7)–(5.2) are exact; no modelling has entered. The tensor M_{ijkl} is a linear functional of the *full three-dimensional* spectrum $\Phi_{mn}(\boldsymbol{\kappa})$ through the nonlocal projection operator $\kappa_a \kappa_b / \kappa^2$, which is the wavenumber-space image of the inverse Laplacian in (5.1). This nonlocality and three-dimensionality are the obstruction that the single-line model must confront.

5.2. The obstruction: what an ODT line resolves

ODT evolves a velocity vector $\mathbf{u}'(\mathbf{y})$ on a single line aligned with the resolved direction, here taken along the contraction axis of the strain, $y \equiv x_2$. From it the model has access to the one-dimensional spectra

$$\phi_{mn}(\kappa_2) = \int_{\mathbb{R}^2} \Phi_{mn}(\boldsymbol{\kappa}) d\kappa_1 d\kappa_3, \quad (5.3)$$

the one-dimensional spectra obtained by integrating Φ_{mn} over the two unresolved wavenumber components. The integrand of (5.2), by contrast, carries the *direction-resolved* weight $\kappa_a \kappa_b / \kappa^2$, which cannot be reconstructed from $\phi_{mn}(\kappa_2)$ alone: integrating out κ_1, κ_3 as in (5.3) destroys exactly the angular information that the projection operator acts on. The reduction $M_{ijkl} \rightarrow$ (functional of ϕ_{mn}) is therefore not available without an assumption about how the energy is distributed over the unresolved wavevector directions. This is the crux of the present question, and we make the required assumption explicit rather than implicit.

The obstruction just stated does not disappear under any symmetry assumption short of full isotropy. As § 5.3 shows, axisymmetry about the line reduces the unknown spectrum to two scalar functions of $(\kappa_2, \kappa_\perp)$ —but the line data are two weighted $\kappa_\perp$ -integrals of those functions per resolved wavenumber, so the map from what the line carries to what the rapid term requires retains an infinite-dimensional null space. Stating this null space exactly, and extracting what survives it, is the correct formulation of the single-line question.

5.3. The closure assumption and the collapsed form

The spectrum tensor of the incoming turbulence, before the rapid distortion acts, is axisymmetric about the resolved direction $\mathbf{e}_2$: $\Phi_{mn}(\boldsymbol{\kappa})$ depends on $\boldsymbol{\kappa}$ only through κ_2 and $\kappa_\perp = (\kappa_1^2 + \kappa_3^2)^{1/2}$, and its polarisation is invariant under rotation about $\mathbf{e}_2$.

This is a statement about the *turbulence*, not about the strain. It is justified for the present configuration because the inflow is grid turbulence, which is closely isotropic at the grid and is subsequently contracted along a single axis as it approaches the stagnation plane; isotropy is the special case of A1, and uniaxial contraction preserves axisymmetry about the contraction axis. The plane strain $A_{kl} = \text{diag}(\frac{1}{2}, -\frac{1}{2}, 0)$ acting at the leading edge is itself *not* axisymmetric, so A1 is imposed on the state on which the rapid operator acts, consistent with the rapid-distortion ordering in which $\Pi^{(r)}$ is evaluated on the current (here, axisymmetric to leading order) field. The error incurred when the accumulated distortion has driven the spectrum away from axisymmetry is quantified in § 5.5.

Under A1 the angular integral in (5.2) can be carried out. Writing $\boldsymbol{\kappa} = (\kappa_\perp \cos \theta, \kappa_2, \kappa_\perp \sin \theta)$ and using that, by A1, Φ_{mn} is independent of θ , the azimuthal averages of the projection

weights reduce to

$$\left\langle \frac{\kappa_1^2}{\kappa^2} \right\rangle_\theta = \left\langle \frac{\kappa_3^2}{\kappa^2} \right\rangle_\theta = \frac{1}{2} \frac{\kappa_\perp^2}{\kappa^2}, \quad \left\langle \frac{\kappa_2^2}{\kappa^2} \right\rangle_\theta = \frac{\kappa_\perp^2}{\kappa^2}, \quad \left\langle \frac{\kappa_1 \kappa_3}{\kappa^2} \right\rangle_\theta = 0, \quad (5.4)$$

with $\kappa^2 = \kappa_2^2 + \kappa_\perp^2$, and all weights mixing a resolved and an unresolved index (e.g. $\kappa_1 \kappa_2 / \kappa^2$) averaging to zero. Substituting (5.4) into (5.2) removes the azimuthal dependence and leaves a two-dimensional integral over $(\kappa_2, \kappa_\perp)$ of the axisymmetric spectrum. Defining the diagonal one-dimensional spectra along the line,

$$E_n^\parallel(\kappa_2) = \int_0^\infty \Phi_{nn}(\kappa_2, \kappa_\perp) 2\pi \kappa_\perp d\kappa_\perp \quad (\text{no sum}), \quad (5.5)$$

the rapid pressure–strain term reduces to a two-dimensional integral over the half-plane $(\kappa_2, \kappa_\perp)$ of the axisymmetric spectrum. The axisymmetric, solenoidal spectrum is fixed by two scalar functions of $(\kappa_2, \kappa_\perp)$ (Batchelor 1946; Chandrasekhar 1950); we denote them $a(\kappa_2, \kappa_\perp)$ and $c(\kappa_2, \kappa_\perp)$, where a is the isotropic-like scalar (the sole survivor when the turbulence is isotropic) and c measures the axisymmetric departure from isotropy. Carrying out the azimuthal average (5.4) and specialising to the plane strain $A = \text{diag}(\frac{1}{2}, -\frac{1}{2}, 0)$, the diagonal rapid pressure–strain components are

$$\Pi_{nn}^{(r)} = \int_0^\infty \int_0^\infty g_n(x) [a(\kappa_2, \kappa_\perp), c(\kappa_2, \kappa_\perp)] 2\pi \kappa_\perp d\kappa_\perp d\kappa_2 \quad (\text{no sum on } n), \quad (5.6)$$

with $x \equiv \kappa_2^2 / \kappa^2$, $\kappa^2 = \kappa_2^2 + \kappa_\perp^2$, and the kernels, linear in the two spectral scalars,

$$g_1(x) = a\left(\frac{1}{8} + \frac{3}{4}x - \frac{7}{8}x^2\right) + c \frac{7}{8}x(1-x)^2, \quad (5.7)$$

$$g_2(x) = -\frac{3}{2}x [a(1-x) + c(1-x)^2], \quad (5.8)$$

$$g_3(x) = a\left(-\frac{1}{8} + \frac{3}{4}x - \frac{5}{8}x^2\right) + c \frac{5}{8}x(1-x)^2. \quad (5.9)$$

Equations (5.6)–(5.9) constitute the axisymmetric collapse: *under A1, the rapid pressure–strain term is a functional of the axisymmetric spectral scalars a, c on the resolved half-plane and the mean strain*, with no reference to the unresolved azimuthal structure beyond what A1 fixes. The kernels satisfy two exact constraints that verify the reduction. First, energy conservation: pressure–strain redistributes without changing the trace, and indeed $g_1(x) + g_2(x) + g_3(x) = 0$ identically for all x and for both scalars, so $\Pi_{11}^{(r)} + \Pi_{22}^{(r)} + \Pi_{33}^{(r)} = 0$. Second, the isotropic limit: setting $c = 0$ with a a function of κ alone, the integral of g_3 over the half-plane vanishes, so $\Pi_{33}^{(r)} = 0$ —as it must, since the plane strain has no component along x_3 and the isotropic rapid term is proportional to S_{ij} (Crow 1968). Both constraints hold identically in the derived kernels, confirming the tensor reduction.

The status of (5.6)–(5.9) must be stated precisely, because it is here that the closure question is decided. Under A1 the rapid term is a linear functional of the two axisymmetric scalars $a(\kappa_2, \kappa_\perp)$ and $c(\kappa_2, \kappa_\perp)$ on the resolved half-plane—a *two-dimensional* representation, not yet a single-line closure, since the line determines only the $\kappa_\perp$ -integrals (5.5) of fixed combinations of a and c and not the functions themselves. Equations (5.6)–(5.9) are therefore the *collapsed form* on which any single-line statement must be built; the single-line content itself—what the line data pin down, and how sharply—is established next.

e	$Sk_t/\varepsilon = 0.8$					$Sk_t/\varepsilon = 16$				
	b_{22}	γ_{eff}	$m=2$	$\frac{\phi_{11}}{\phi_{33}} - 1$		b_{22}	γ_{eff}	$m=2$	$\frac{\phi_{11}}{\phi_{33}} - 1$	
0	0.003	0.10	0.133	+0.06		0.003	0.10	0.133	+0.06	
0.5	0.064	0.79	0.166	-0.20		0.070	1.28	0.247	-0.21	
1	0.125	1.44	0.189	-0.32		0.124	4.22	0.396	-0.46	

Table 3. A1 measured under plane strain (128³ DNS, four realisations): upwash anisotropy, effective polarisation, azimuthal $m = 2$ residue of Φ_{22} (isotropic floor 0.133), and the on-line falsifier T1.

5.4. What line data determine: sharp bounds on the rapid term

Under A1 the solenoidal spectrum tensor takes the two-scalar form $\Phi_{mn} = a P_{mn} + c \xi_m \xi_n$, with P_{mn} the solenoidal projector and ξ_m the projected axis direction; realisability is the pair of pointwise conditions $a \geq 0$ and $a + (1 - x)c \geq 0$ with $x = \kappa_2^2/\kappa^2$. The line carries $\phi_{11}(\kappa_2) = \phi_{33}(\kappa_2)$ and $\phi_{22}(\kappa_2)$: per resolved wavenumber, two weighted $\kappa_\perp$ -integrals of the two unknown functions. The rapid term (5.6) is linear in (a, c) , so its extreme values over everything consistent with the line data, solenoidality and realisability are obtained from a family of linear programs, one per κ_2 , yielding sharp bounds $[I_{nn}^-, I_{nn}^+]$ and a kinematic indeterminacy $\mathcal{I}_n = (I_{nn}^+ - I_{nn}^-)/2k_t S$ that quantifies exactly how much the null space costs. The forward map from (a, c) to the line spectra and the derivation of the bound kernels are given in Appendix A.

Calibrated on an isotropic von Kármán spectrum (with a verification suite pinning tracelessness, the isotropic angular averages $+\frac{1}{5} : -\frac{1}{5} : 0$, the longitudinal–transverse relation, and the exact $\frac{4}{5}k_t S_{ij}$), the indeterminacy is

constraint level	$\mathcal{I}_{11}$	$\mathcal{I}_{22}$	$\mathcal{I}_{33}$
line data + solenoidality + realisability	0.77	1.28	0.86
+ log-Lipschitz slope cap ($\lambda = 4$)	< 5% narrower		
+ polarisation cap $ c \leq \gamma a$, $\gamma = 0$	0.34	0.58	0.24

Two features of this table matter. Smoothness assumptions buy almost nothing: the null space is not a roughness artefact. The polarisation cap is the lever—but § 5.5 shows that under strain the effective polarisation reaches order one, so the honest bound in the strained regime is the uncapped one.

The representation also supplies two *free falsifiers* that the line itself reports: under A1, $\phi_{11}(\kappa_2) = \phi_{33}(\kappa_2)$ at every wavenumber (T1) and the velocity cross-spectra vanish (T2). Any measured transverse splitting is a wavenumber-resolved violation of A1; rather than assuming the hypothesis, the line monitors it.

5.5. The axisymmetry hypothesis, measured

A1 is exact for the isotropic inflow and is degraded by the plane strain itself. Rather than estimating this error perturbatively, we measure it in the strained-box DNS of § 4.2, where the full spectrum is available: the exact azimuthal decomposition of Φ_{22} yields the true scalars (a, c) , the discarded $m = 2$ azimuthal residue (the direct A1-violation measure, with isotropic sampling floor 0.133 in this box), and the effective polarisation $\gamma_{\text{eff}} = |c|/a$.

Three conclusions follow (table 3). First, the A1 error is *not* perturbatively small in the accumulated anisotropy: the residue grows from its floor to 0.19 at $Sk_t/\varepsilon = 0.8$ and to 0.40 at 16 by $e = 1$ —larger at rapid strain, because at moderate strain rate the nonlinear dynamics restore axisymmetry as fast as the strain destroys it. An $O(b^2)$ error estimate, which the symmetric structure of the discarded harmonic suggests, is superseded by this




Figure 15. The A1 control experiment at $Sk_t/\varepsilon = 16$ (128³, four realisations): axisymmetric contraction (A1 exact by symmetry) against plane strain at matched A_{22} . Left: azimuthal $m = 2$ residue of Φ_{22} . Right: transverse splitting and b_{22} .

measurement. Second, the polarisation cap that narrows the isotropic bounds is illegitimate under strain: γ_{eff} passes unity by $e \approx 0.5$. Third, correspondingly, the uncapped bound brackets the exact rapid term (evaluated by direct integration in the DNS) up to $e \approx 0.25$ – 0.5 and fails beyond, and the failure is A1’s, not the estimator’s.

The decisive evidence for that attribution is a control experiment: the same box under *axisymmetric contraction* with matched A_{22} , for which A1 is exact by symmetry. There the $m = 2$ residue stays at the sampling floor and the transverse splitting at zero at every strain up to $e = 1$ even at $Sk_t/\varepsilon = 16$, while under plane strain both grow (figure 15). The single-line closure is therefore *exact for axisymmetric distortion*; its plane-strain error is a strain-geometry property, now quantified, rather than a defect of the line representation. For the applications this maps cleanly: a rounded, three-dimensional stagnation region imposes an axisymmetric strain (closure exact); a two-dimensional leading edge imposes plane strain (closure carries the measured correction). The acoustically relevant amplification $b_{22}(e)$, finally, is indistinguishable between the two geometries (0.124 against 0.117 at $e = 1$, $Sk_t/\varepsilon = 16$): the quantity the leading-edge response consumes is robust to precisely the geometry that A1 is sensitive to.

5.6. Component ordering under plane strain

The exact solution of the rapid problem under plane strain has the neutral spanwise component eventually overtaking the upwash; algebraic closures predict the opposite. The exact rapid term evaluated in the DNS settles the matter: at $e = 1$, $\Pi_{33}^{(r)}/k_t S = 0.14$ at $Sk_t/\varepsilon = 0.8$ and 0.27 at 16, exceeding $\Pi_{11}^{(r)} = 0.23$ in the rapid case—the overtaking is real and sets in at rapid strain. LRR-QI, whose spanwise component is carried by its b -dependent term, matches at $Sk_t/\varepsilon = 0.8$ ($\Pi_{33}^{(r)} = 0.143 k_t S$) but delivers half the exact value at 16; IP gives zero identically. The deficiency thus lives in the moment closure’s b -term, not in the line model: the ODT moment-level b_{ij} follows the closure to $\sim 2\%$, and its b_{33} stays within 0.006 of zero in every run. In the spectral representation the ordering is carried by the c -scalar through the kernels (5.7)–(5.9); the moment reduction discards it. This answers, by measurement, where the known component-ordering error enters and what it would take to remove it.

5.7. Reduction to the moment closure

The closed form (5.6)–(5.9) operates on the resolved spectrum, whereas the distortion results of § 4 were obtained with the algebraic moment closure of § 2.4 (Launder *et al.* 1975). For the two to be consistent, the spectral kernel must reduce to that moment closure in the limit where the moment closure is exact—isotropic turbulence. We show that it does, which both justifies the closure used in the simulations and fixes the prefactor in an explicit convention.

Figure 16 plots the kernels g_1, g_2, g_3 of (5.7)–(5.9) against the angular variable $x = \kappa_2^2/\kappa^2$. Two features are exact and hold for both spectral scalars. The kernels sum to zero at every x ,

$$g_1(x) + g_2(x) + g_3(x) = 0 \quad \forall x, \quad (5.10)$$

so the rapid term conserves turbulent kinetic energy pointwise in wavenumber, not merely in the integral; and each kernel vanishes at $x = 1$, where the wavevector aligns with the line and the projection $\kappa_a \kappa_b / \kappa^2$ degenerates. The signs are physically ordered: $g_1 > 0$ (energy fed to the streamwise component E_1), $g_2 < 0$ (energy removed from the line/upwash component E_2) over the bulk of the angular range, with g_3 changing sign, consistent with the plane strain stretching x_1 and compressing x_2 .

For *isotropic* turbulence the anisotropy scalar vanishes ($c = 0$) and the isotropic-like scalar a depends only on κ . Averaging the kernels over solid angle—equivalently, integrating $g_n(x)$ against the isotropic distribution of x on the sphere—gives

$$\langle g_1 \rangle : \langle g_2 \rangle : \langle g_3 \rangle = +\frac{1}{5} : -\frac{1}{5} : 0, \quad (5.11)$$

in the ratio $+1 : -1 : 0$. This is exactly the structure of the plane strain $S_{ij} = \text{diag}(\frac{1}{2}, -\frac{1}{2}, 0)$: the rapid term is proportional to S_{ij} , with $\Pi_{33}^{(r)} = 0$ because the strain has no component along x_3 . The spectral kernel therefore collapses, under isotropy, to the isotropic-production form of the moment closure,

$$\Pi_{ij}^{(r)} = \frac{4}{5} k_t S_{ij} \quad (\text{isotropic limit}), \quad (5.12)$$

the result of Crow (1968) that underlies the isotropisation-of-production (IP) rapid term in the closure of Launder *et al.* (1975). We note a convention: with the symmetrised definition of the pressure–strain correlation adopted here and the factor of two carried in the Poisson source (2.7), the kernels (5.7)–(5.9) yield the coefficient $\frac{2}{5}k_t$ for the 11-component directly (since $S_{11} = \frac{1}{2}$), equivalent to the standard $\frac{4}{5}k_t S_{ij}$ of (5.12); the ratio (5.11) and the tracelessness (5.10) are independent of this convention.

The consequence is that the algebraic closure used to obtain the distortion of § 4 is not an independent assumption but the isotropic-limit reduction of the line-consistent spectral term derived here. The spectral form (5.6) carries, in addition, the angular (x -dependent) and anisotropy (c -scalar) information that the moment closure discards—and both discards are now measured: the wavenumber dependence of the linear response in the three-way comparison of § 4.2.1, and the component ordering in § 5.6. The moment closure and the collapsed form thus agree to leading order, and the terms in which they depart are exactly the measured deficiencies of § 4.




Figure 16. Rapid pressure–strain closure kernels g_1, g_2, g_3 (eqs. (5.7)–(5.9)) against the angular variable $x = \kappa_2^2/\kappa^2$. Solid curves: coefficient of the isotropic-like spectral scalar a ; faint dashed curves: coefficient of the anisotropy scalar c . The kernels sum to zero at every x (energy conservation, eq. (5.10)) and vanish at $x = 1$. Averaged over an isotropic spectrum they give the ratio $+\frac{1}{5} : -\frac{1}{5} : 0$ (eq. (5.11)), recovering the isotropisation-of-production form $\Pi_{ij}^{(r)} = \frac{4}{5} k_t S_{ij}$ (Crow 1968) that underlies the moment closure used in § 4.

5.8. Consistency with the eddy redistribution: avoiding double counting

A closed rapid term is necessary but not sufficient: it must not duplicate redistribution already effected by the eddy events. The pressure–strain correlation decomposes canonically into rapid and slow parts, $\Pi_{ij} = \Pi_{ij}^{(r)} + \Pi_{ij}^{(s)}$, corresponding to the two source terms in (2.7). The slow part $\Pi_{ij}^{(s)}$ is the return-to-isotropy mechanism, quadratic in the fluctuations and independent of the mean gradient; the eddy events in ODT, which act on the fluctuating field without reference to A_{kl} , are the model’s representation of exactly this slow redistribution. The rapid part $\Pi_{ij}^{(r)}$, proportional to A_{kl} and linear in the fluctuations, is by construction absent from the eddy mechanism, which carries no dependence on the mean strain. The two therefore occupy disjoint terms of the canonical decomposition, and adding (5.6) as an explicit mean-strain-driven source alongside the eddies does not double-count, provided the eddy events are calibrated to reproduce the slow return-to-isotropy in the absence of mean strain—a condition that the no-strain baseline of § 4.0.3 verifies directly, since there the centroid relaxes under the eddies alone with no spurious rapid contribution. This argument establishes consistency at the level of the decomposition; the no-strain baseline of § 4.0.3 provides the direct numerical confirmation, the centroid there relaxing under the eddy mechanism alone.

The disjointness argument is now also a measured statement, from both sides. With the strain switched off the eddy events generate no spurious strain-like signal (the nominal-turbulence null of Appendix B). Conversely, in the rapid limit ($Sk_t/\varepsilon = 16$, under one eddy per line in $e \leq 1$) every kernel-energy allocation rule tested—the isotropic kernel, a strain-biased generalisation, and discrete eddy types—reproduces the exact rapid-distortion evolution of $b_{22}(e)$ to ± 0.003 : the rapid term cannot be reached from inside the eddy events, and the continuous operator carries it alone. Rapid and slow redistribution occupy disjoint terms by construction, and now by measurement.

5.9. Summary: what the single-line treatment establishes

1. The rapid term is exactly $A_{kl}M_{ijkl}$; its single-line representation is obstructed by an azimuthal null space that is stated exactly and survives the axisymmetric collapse at the level of the two spectral scalars.
2. Under A1 the term collapses to the kernel form (5.6)–(5.9); A1 is exact for axisymmetric distortion and is *measured* under plane strain, where its violation grows with accumulated strain and with strain rapidity (table 3).
3. Independently of A1's validity, line data bound the rapid term sharply; the bound is honest—and verified to bracket the exact term—up to $e \approx 0.25$ – 0.5 under plane strain, and at all computed strains under axisymmetric distortion.
4. The moment closure used in the simulations is the isotropic reduction of the collapsed form (verified, including the exact $\frac{4}{3}k_t S_{ij}$); what it discards is the wavenumber dependence of the linear response and the component ordering, both measured in § 4 and § 5.6.
5. Rapid and slow redistribution are disjoint: verified by the strain-off null and by the rapid-limit invariance of every kernel-energy allocation rule.
6. The acoustically relevant upwash amplification $b_{22}(e)$ is linear to $e = 1$ and robust to the strain geometry that controls A1's validity.

6. Measured limits, and scale-local relaxation

The envelope of § 4.2.2 localises the model's spectral deficiency in its wavenumber-uniform rapid kinematics: the strain-induced anisotropy is allocated uniformly across scales where linear theory and DNS concentrate it at the energy-containing range and remove it from the fine scales. This section asks the constructive question: which mechanism available to the model can supply the missing scale dependence? We answer it experimentally, by a scale-resolved diagnostic and a controlled family of interventions in the eddy mechanism, each tested at 1024 realisations with seeds paired to a common baseline, and each either falsified or confirmed by the same measurement. The outcome is a single measured statement: what is missing is *scale-local relaxation*, and supplying it—at the scales of the imbalance—is both necessary and, at moderate strain, sufficient to move the fine-scale anisotropy.

6.1. A transmission diagnostic

The configuration is a scale-diagnostic variant of the strained line: unit strain rate (so $e = t$), an initial spectrum peaked at a wavenumber κ_p resolved by more than a decade of smaller scales, and two measurement bands, $\kappa_2 \in [0.6, 2] \kappa_p$ (the energy-containing band) and $[6, 16] \kappa_p$ (roughly two triplet-map generations below it, since each map compresses by a factor of three). On each band the component imbalance is $\mathcal{A}_b = E_2^{\parallel}(b)/\frac{1}{2}[E_1^{\parallel}(b)+E_3^{\parallel}(b)]$, the band-integrated upwash energy over the transverse mean; all statistics are medians over realisations with bootstrap confidence intervals, for the intermittency reason given in § 4.2. The model's eddy and kernel mechanisms are component-symmetric, so $\mathcal{A} = 1$ is their fixed point; the strain drives $\mathcal{A}$ up at the scales where it deposits anisotropy, and the ratio $\mathcal{A}_{hi}/\mathcal{A}_{lo}$ measures how much of the large-scale imbalance survives transmission down the cascade.

The baseline (figure 17, black) exhibits the deficiency of § 4.2.1 in transmission form: $\mathcal{A}_{hi}/\mathcal{A}_{lo}$ stays near unity out to $e = 3$ (1.02, 0.95, 0.89 at $e = 1, 2, 3$)—the

anisotropy injected at the energy-containing scales crosses two map generations essentially undiminished—and drops only at the deepest strain. The mechanism is direct: the high band is populated by the maps of *large* eddies (one generation takes κ_p to $3\kappa_p$, a second to $9\kappa_p$, inside the band), so fine-scale anisotropy is delivered by large-eddy events, not by a chain of small ones.

6.2. Interventions: what fails and what works

Table 4 summarises the intervention family. Two families fail for instructive reasons. Acceptance gating—rejecting accepted eddies whose post-event state remains anisotropic, at thresholds spanning 45 to 98 per cent rejection—is scale-blind: at mild thresholds the surviving eddy population reproduces the baseline anisotropy budget in every band (the gate self-selects eddies whose kernels were already going to do the relaxing, and the unrelaxed field keeps generating candidates), while at the harshest threshold the model merely drifts toward its eddy-free limit. Restricting the gate to sub-band-scale eddies sharpens the negative result into the mechanism statement above: removing up to 97 per cent of small-eddy events leaves both the fine-scale anisotropy *and* the fine-scale energy unchanged, so no acceptance-gating scheme can control transmission at this Reynolds number. Reweighting the triplet-map images (a middle image of relative size $\frac{2}{3}$, which halves the compression of the map’s dominant central channel) moves one-point and energy-containing statistics— $\mathcal{A}_{10}$ falls significantly and the upwash energy fraction with it—but not the fine scales: with the image fractions summing to one, slowing the middle channel necessarily speeds the outer ones, and one outer image maps the energy peak directly into the measurement band.

What works is relaxation applied *at the scale of the imbalance*. After each accepted eddy’s map and kernels, the same energy-equalising kernel procedure (§ 2) is applied to each of the three map images—the sub-interval’s own triplet-map displacement shape as the kernel, coefficients from the sub-interval’s own integrals, so that each sub-application conserves the momentum components and the total kinetic energy exactly, with no map applied and no random numbers drawn—and then recursively to 9 and 27 sub-regions. One level ($\ell/3$ for an eddy of size ℓ) is null: for the large eddies that feed the band, $\ell/3$ still sits above it. Two levels produce the first statistically significant fine-scale isotropisation of the family ($\mathcal{A}_{hi}$ from 1.43 to 1.27 at $e = 1$, both bands’ paired contrasts individually significant), and three levels extend the effect to deep strain: the levels that work, $\ell/9$ and $\ell/27$, are exactly the first that reach the measured band. The bookkeeping is clean throughout—band energies move by a few per cent, the upwash energy fraction and the intermittency tails are unchanged—and the pre-strain fine-scale floor moves toward component equality immediately ($\mathcal{A}_{hi}$ from 0.83 to 0.95 at $e = 0$).

Four further variants separate the candidate explanations for the one remaining limitation, the fading of the gains beyond $e \approx 3$. Tying the recursion depth to the eddy size (subdividing while the sub-interval remains above a near-dissipative floor, so that large eddies do proportionally more sub-scale work) gives the gains of depth with no reversal at the deepest strain: both bands significant at $e = 1$ and the effect retaining its sign through $e = 3$. Iterating the two-level pass three times within each event answers the amplitude hypothesis in the negative: it is a stronger early isotropiser but overshoots at deep strain. Delocalising the relaxation—after each eddy, applying the kernel isotropisation in N randomly placed intervals of the same size, overlaps allowed—tests whether the *location* of the events is the limiter: $N = 1$ is null, and $N = 3$ is significant in both bands at $e = 1$ but acts mainly at the energy-containing scales, since the sampled intervals inherit the eddy-size distribution and full-interval isotropisation acts at that scale. Stacking the two ideas—the

intervention (knob)	fine-scale effect	reading
acceptance gate on post-event anisotropy (45–98% rejection)	null at all thresholds; harsh-est drifts to eddy-free limit	gate is scale-blind: modulates eddy rate, not the scale allocation
gate on sub-band eddies only (42%, 97% rejection)	null, including band <i>energy</i>	fine scales are fed by large-eddy maps directly
unequal map images (middle $\frac{2}{3}$)	null (one-point statistics move: $\mathcal{A}_{l_0}$ 2.85 $\rightarrow$ 2.50 at $e = 3.9$)	outer images feed the band in one step; channels trade off
sub-scale kernels, 1 level ($\ell/3$)	null within CIs	$\ell/3$ sits above the measured band
sub-scale kernels, 2 levels ($\ell/9$)	$\mathcal{A}_{hi}$: 1.43 $\rightarrow$ 1.27 at $e = 1$ (significant, both bands)	first significant fine-scale isotropi-sation
sub-scale kernels, 3 levels ($\ell/27$)	significant at $e = 1$ and $e = 3$ (–12% paired)	deepest levels reach the band
depth tied to eddy size	significant in both bands at $e = 1$; no late-strain reversal	best of the family: gains without the deep-strain overshoot
2 levels iterated $3\times$ per event	strongest at $e = 1$ (1.29); overshoots at deep strain	per-event amplitude is not the limiter
concurrent relaxation: N same-size intervals, randomly placed ($N = 1, 3$)	$N = 1$ null; $N = 3$ significant in both bands at $e = 1$, strongest at the energy-containing scales; fade persists	the sampled intervals act at eddy scale; delocalising the events alone is not the lever
concurrent + depth ($N = 3$, adaptive hierarchy in each interval)	strongest early effect of the family ($\mathcal{A}_{hi}$ 1.43 $\rightarrow$ 1.26); low band significant even at $e = 3.9$; fine scales overshoot depth-only at deep strain	dosage helps early and at large scales, and harms deep-strain fine scales wherever it is applied

Table 4. The intervention family on the transmission diagnostic (1024 paired-seed realisations per case; effects quoted as median paired same-seed contrasts, “significant” meaning the 95% bootstrap interval excludes zero).

adaptive hierarchy inside each sampled interval—is the strongest early isotropiser of the family ($\mathcal{A}_{hi}$ from 1.43 to 1.26 at $e = 1$) and delivers the only significant deep-strain effect, in the energy-containing band at $e = 3.9$; yet at the fine scales and deep strain it overshoots the depth-only variant significantly. The ledger is then complete: depth sets *where* the relaxation acts and is necessary; dosage and delocalisation raise the early gains in proportion to the total relaxation delivered; and no combination tested—more depth, more dosage, more events, other locations—prevents the fine-scale fade beyond $e \approx 3$. The deep-strain fine scales are resupply-limited, not implementation-limited: relaxation acts at the event rate while the strain resupplies the imbalance continuously, and at these total strains no per-event scheme keeps pace. Whether that saturation is a model artefact or the physical statement that sustained rapid distortion outruns any cascade-mediated return to isotropy is not decided by these experiments.

6.3. The measured missing ingredient

The arc closes the loop opened by the three-way comparison. The deficiency measured there—anisotropy allocated uniformly over wavenumber—is confirmed here in transmission form, is unfixable by gating or map reweighting, and is corrected, at moderate strain, by exactly one ingredient: relaxation applied at the scale of the imbalance. At the formulation level this ingredient has a natural continuous home: the spectral kernel form




Figure 17. Scale-local relaxation on the transmission diagnostic: (a) energy-containing-band imbalance $\mathcal{A}_{lo}$, (b) fine-scale-band imbalance $\mathcal{A}_{hi}$, (c) transmission $\mathcal{A}_{hi}/\mathcal{A}_{lo}$, (d) upwash energy fraction, against total strain: standard ODT, hierarchical sub-scale kernels at fixed depth (1–3 levels), depth tied to eddy size, the per-event iterated variant, concurrent relaxation ($N = 3$ randomly placed same-size intervals per event), and the stacked variant (concurrent intervals each running the adaptive hierarchy). Medians over 1024 paired-seed realisations, 95% bootstrap intervals.

(5.6)–(5.9) is precisely a wavenumber- and angle-dependent rapid operator, and replacing the wavenumber-uniform moment closure of § 2 by a discretisation of it is the formulation-level analogue of the sub-scale kernels’ event-level correction. The present results bound what such an upgrade can and cannot deliver: it can re-allocate the rapid redistribution across scales (the measured deficit of § 4.2.1), and it cannot, any more than the event-level mechanisms, outrun sustained resupply at deep strain.

7. Consequence for leading-edge noise

[M5: detachable LE section — decision D1.]

8. Conclusions

[M4: conclusions rewrite.]

Appendix A. The forward map and the bound kernels

This appendix derives the three ingredients that § 5 uses: the two-scalar representation of the axisymmetric solenoidal spectrum with its realisability cone, the forward map from the two scalars to the line spectra (with the falsifiers T1–T2), and the closure kernels (5.7)–(5.9), together with the linear programs that turn these linear relations into the sharp bounds of § 5.4.

A.1. The axisymmetric representation and realisability

The spectrum tensor of statistically homogeneous turbulence is Hermitian, positive semi-definite at each κ , and solenoidal, $\kappa_m \Phi_{mn}(\kappa) = 0$; it therefore has rank at most two, supported on the plane normal to κ . Under A1 (axisymmetry about e_2 , mirror-symmetric, helicity-free) it is fixed by two scalar functions of $(\kappa_2, \kappa_\perp)$ (Batchelor 1946; Chandrasekhar 1950; Lindborg 1995):

$$\Phi_{mn}(\kappa) = a(\kappa_2, \kappa_\perp) P_{mn}(\kappa) + c(\kappa_2, \kappa_\perp) \xi_m \xi_n, \quad \xi = e_2 - \mu \frac{\kappa}{\kappa}, \quad \mu = \frac{\kappa_2}{\kappa}, \quad (\text{A } 1)$$

with $P_{mn} = \delta_{mn} - \kappa_m \kappa_n / \kappa^2$ the solenoidal projector, so that $\kappa_m \xi_m = 0$ and $|\xi|^2 = 1 - \mu^2 = 1 - x$ with $x = \kappa_2^2 / \kappa^2$: solenoidality is exact by construction, which is how the third would-be scalar of a general axisymmetric tensor is eliminated. For a unit vector v normal to κ , $\Phi_{mn} v_m v_n = a + c (\xi \cdot v)^2$, so the two eigenvalues of the polarisation matrix on that plane are a (attained for $v \perp \xi$) and $a + (1 - x) c$ (for $v \parallel \xi$). Positive semi-definiteness is therefore the pointwise *realisability cone*

$$a(\kappa_2, \kappa_\perp) \geq 0, \quad a(\kappa_2, \kappa_\perp) + (1 - x) c(\kappa_2, \kappa_\perp) \geq 0, \quad (\text{A } 2)$$

quoted in § 5.4. Isotropy is the special case $c = 0$ with a a function of κ alone.

A.2. The forward map to the line spectra

Writing $\kappa = (\kappa_\perp \cos \theta, \kappa_2, \kappa_\perp \sin \theta)$, the azimuthal moments at fixed $(\kappa_2, \kappa_\perp)$ are $\langle \kappa_1^2 \rangle_\theta = \langle \kappa_3^2 \rangle_\theta = \frac{1}{2} \kappa_\perp^2$, $\langle \kappa_1^4 \rangle_\theta = \langle \kappa_3^4 \rangle_\theta = \frac{3}{8} \kappa_\perp^4$, $\langle \kappa_1^2 \kappa_3^2 \rangle_\theta = \frac{1}{8} \kappa_\perp^4$, and every odd moment vanishes. Applied to (A 1) the diagonal weights average to

$$\langle P_{11} \rangle_\theta = \langle P_{33} \rangle_\theta = \frac{1}{2} (1 + x), \quad P_{22} = 1 - x, \quad \langle \xi_1^2 \rangle_\theta = \langle \xi_3^2 \rangle_\theta = \frac{1}{2} x (1 - x), \quad \xi_2^2 = (1 - x)^2, \quad (\text{A } 3)$$

and every mixed weight to zero. Substituting into the definition (5.3) of the line spectra, $\phi_{mn}(\kappa_2) = \int_0^\infty \langle \Phi_{mn} \rangle_\theta 2\pi \kappa_\perp d\kappa_\perp$, gives the **forward map** from the two scalars to the data a line carries:

$$\phi_{11}(\kappa_2) = \phi_{33}(\kappa_2) = 2\pi \int_0^\infty \left[a \frac{1+x}{2} + c \frac{x(1-x)}{2} \right] \kappa_\perp d\kappa_\perp, \quad (\text{A } 4)$$

$$\phi_{22}(\kappa_2) = 2\pi \int_0^\infty \left[a (1 - x) + c (1 - x)^2 \right] \kappa_\perp d\kappa_\perp, \quad (\text{A } 5)$$

$$\phi_{mn}(\kappa_2) = 0 \quad (m \neq n). \quad (\text{A } 6)$$

Equations (A 4)–(A 6) contain the two free falsifiers quoted in § 5.4: under A1 the two transverse line spectra coincide at every resolved wavenumber (T1), and the cross-spectra vanish (T2); both are properties the line itself reports, scale by scale, with no reference to unresolved data. The ϕ_{22} bracket is the same combination of a and c that appears in the kernel g_2 below—the conventions mesh, as they must, because ϕ_{22} and the 22-component of the rapid term weight the spectrum through the same projection geometry.

They also explain why the isotropic intuition about line data fails under A1. For isotropic turbulence ($c = 0$, $a = a(\kappa)$) the map is over-determined: one unknown function of one variable against two measured functions, with ϕ_{22} determining a by the classical Abel-type inversion and the transverse spectrum slaved to it by the derivative relation. Under A1 the bookkeeping reverses: *two unknown functions of two variables* against two measured functions of one variable. At each fixed κ_2 the data supply exactly two numbers—the two weighted $\kappa_\perp$ -integrals (A 4)–(A 5)—about two unknown functions of $\kappa_\perp$; any perturbation $(\delta a, \delta c)(\kappa_\perp)$ annihilating both integrals while respecting (A 2) is invisible to the line. This infinite-dimensional null space is the quantitative content of the obstruction of § 5.2 after the axisymmetric collapse.

880

A.3. Derivation of the closure kernels

For the diagonal components the exact result (5.2) reads

$$\Pi_{nn}^{(r)} = 2 A_{kl} \int_{\mathbb{R}^3} \frac{\kappa_n \kappa_k}{\kappa^2} \Phi_{nl}(\boldsymbol{\kappa}) d\boldsymbol{\kappa} \quad (\text{no sum on } n). \quad (\text{A } 7)$$

Substituting (A 1), carrying out the azimuthal average with the moments above, and specialising to $A = \text{diag}(\frac{1}{2}, -\frac{1}{2}, 0)$ reduces each component to a $2\pi\kappa_\perp$ -weighted integral over the half-plane of a polynomial combination in x , linear in (a, c) —the kernels (5.7)–(5.9) of the main text. The upwash component illustrates the mechanics: only $k = l = 1$ and $k = l = 2$ contribute, with azimuthal averages $\langle \kappa_1 \Phi_{21} \rangle_\theta = -\frac{1}{2} \kappa_\perp^2 [a + (1-x)c]/\kappa^2$ and $\langle \Phi_{22} \rangle_\theta = (1-x)[a + (1-x)c]$, so that both enter through the same bracket and

$$\langle \text{integrand of (A 7)} \rangle_\theta \Big|_{n=2} = 2 \left[-\frac{1}{2} \cdot \frac{x(1-x)}{2} - \frac{1}{2} x(1-x) \right] [a + (1-x)c] = -\frac{3}{2} x(1-x)[a + (1-x)c], \quad (\text{A } 8)$$

881 which is $g_2(x)$. The transverse components involve in addition the fourth moments of (A 3)’s underlying table
 882 and yield g_1 and g_3 ; the full reduction has been verified symbolically against (A 7). Three identities check the
 883 algebra: $g_1 + g_2 + g_3 = 0$ at every x for both scalars (pointwise energy conservation); each kernel vanishes
 884 at $x = 1$, where the wavevector aligns with the line and the projection degenerates; and averaging over the
 885 isotropic angular measure $d\mu$ on $[0, 1]$ gives $\langle g_1 \rangle : \langle g_2 \rangle : \langle g_3 \rangle = +\frac{1}{5} : -\frac{1}{5} : 0$, reproducing $\Pi_{ij}^{(r)} = \frac{4}{5} k_i S_{ij}$
 886 (Crow 1968) as in § 5.7.

887

888

A.4. The bounds as linear programs

The target (5.6), the data constraints (A 4)–(A 5), and the cone (A 2) are all linear in (a, c) , and everything decomposes over the resolved wavenumber: writing $\Pi_{nn}^{(r)} = \int_0^\infty \pi_n(\kappa_2) d\kappa_2$ with $\pi_n = 2\pi \int_0^\infty g_n(x)[a, c] \kappa_\perp d\kappa_\perp$, the extremisation over all spectra consistent with the line data is a family of *independent* infinite-dimensional linear programs, one per κ_2 :

$$\pi_n^\pm(\kappa_2) = \max / \min_{(a,c)(\kappa_\perp)} 2\pi \int_0^\infty g_n(x)[a, c] \kappa_\perp d\kappa_\perp \quad \text{s.t.} \quad (\text{A } 4)\text{--}(\text{A } 5) \text{ hold, (A } 2) \text{ pointwise,} \quad (\text{A } 9)$$

889 whence $\Pi_{nn}^\pm = \int \pi_n^\pm d\kappa_2$ and the indeterminacy $\mathcal{I}_n = (\Pi_{nn}^+ - \Pi_{nn}^-)/2k_i S$ of § 5.4, normalised by the natural
 890 rapid-term scale so that $\mathcal{I}_n$ is comparable across strains (the isotropic value is $\Pi_{11}^{(r)} = \frac{2}{5} k_i$ per unit $S_{11} = \frac{1}{2}$).
 891 Discretised on a logarithmic $\kappa_\perp$ grid of order 10^2 points, (A 9) is a linear program with two equality constraints
 892 and the cone, solved per κ_2 in milliseconds; the bounds are global and sharp by LP duality—no fitting and
 893 no local minima. The side conditions of § 5.4 enter as further linear constraints: the log-Lipschitz slope cap
 894 $|d \ln(a, c)/d \ln \kappa_\perp| \leq \lambda$ and the polarisation cap $|c| \leq \gamma a$. The LP extremals concentrate on few-point supports
 895 in $\kappa_\perp$ (bang–bang spectra), which is why the raw bounds are conservative and why smoothness alone barely
 896 narrows them: a log-Lipschitz spectrum can still vary like $\kappa_\perp^{\pm\lambda}$, so the angular reweighting that drives the
 897 indeterminacy survives smoothing, while the polarisation cap attacks it directly. The dual variables of (A 9)
 898 certify, per resolved wavenumber, which linear combinations of the measured line spectra control each π_n ;
 899 finite-window bias on ϕ_{mn} at small κ_2 propagates linearly and can be carried as interval data in place of the
 900 equalities with no change of machinery.

901

902 Appendix B. Numerical parameters, initial conditions and statistics

903

B.1. ODT configurations

904 All line ensembles share the solver settings of the standard model (Stephens & Lignell 2021): eddy-rate
 905 constant $C = 7$, viscous-penalty parameter $Z = 450$, kernel parameter $\alpha = 2/3$, adaptive mesh with $dx \in$
 906 $[2 \times 10^{-4}, 2 \times 10^{-2}]$ on a unit domain (4000 initial cells), explicit advancement with diffusive CFL 0.5 and a
 907 strain-resolved bound of 10^{-2} on the total-strain increment per step (§ 3.3). The initial condition is the resolved
 908 spectral field of § 3.3: a band-limited isotropic superposition of 64 Fourier modes peaked at eight wavelengths
 909 per domain, which places no energy at the grid scale. Strained production ensembles use an unstrained precursor:
 910 the line evolves as standard ODT until the eddy population is statistically active and the initialisation transient
 911 has relaxed, and the strain is switched on only then (§ 4.2). The molecular viscosity is set per study: 10^{-4} for
 912 the transmission diagnostic of § 6 (a scale diagnostic, resolved with wide margin), 10^{-5} for the strain-on/off
 913 ensembles of § 4.0.3, and 3×10^{-6} on a 16000-cell grid at the operating point of § 4.0.4.

B.2. Strained-box DNS and exact-RDT companions

The benchmark of § 4.2 is a pseudo-spectral 128^3 simulation in a Rogallo deforming frame: a decaying isotropic precursor, then plane strain $A = S \text{diag}(\frac{1}{2}, -\frac{1}{2}, 0)$ at $Sk_t/\varepsilon \in \{0.8, 16\}$ at onset, carried to $e = 1$ over four independent realisations, with $\kappa_{\max}\eta \approx 0.76$ at $e = 1$ —a pilot resolution, so small-scale statements are qualitative. Every realisation has an exact-RDT companion: the same initial field evolved under the closed-form Cauchy solution of the linearised equations and projected onto lines identically. The initial Fourier fields are whitened by the *symmetric* inverse square root $C^{-1/2}$ of the sample covariance, so the initial anisotropy vanishes per realisation. This detail is load-bearing: an ordered-Cholesky whitening used initially left a bias $b_{33} = +0.0148 \pm 0.0046$, detected by the nominal-turbulence null test (strain machinery active, $A = 0$) and traced by a reversal experiment to the ordering; with the symmetric form all $|b_{ii}| \leq 0.005$ and the on-line falsifier T1 exceeds its confidence band at the chance rate. The estimator pipeline of § 5.4 passes the same null battery on JHTDB isotropic turbulence (1024^3 , 128 lines): T1 exceedances at the chance rate, cross-spectral coherences $\sim 10^{-2}$, the longitudinal–transverse derivative relation to 4%, and the LP bounds bracketing the exact isotropic value at every constraint level.

B.3. Statistics policy

Spectral band quantities of the strained line are strongly intermittent across realisations—single realisations carry 10^2 – 10^3 times the median band energy—so ensemble means of band ratios do not converge at any practical sample size. All ODT spectral statistics are therefore medians over realisations with 95% bootstrap confidence intervals of the median, and every intervention in § 6 is assessed by the median of *paired* same-seed contrasts, quoted significant only when the bootstrap interval excludes zero. Ensemble sizes: 1024 realisations for the transmission diagnostic and the distance envelope, 128–200 for the spectrum ensembles of § 4, and four seeds for the DNS and its companions, whose derived quantities carry jackknife errors over realisations.

REFERENCES

- BACHELOR, G. K. 1946 The theory of axisymmetric turbulence. *Proceedings of the Royal Society of London. Series A* **186** (1007).
- BACHELOR, G. K. & PROUDMAN, IAN 1954 The effect of rapid distortion of a fluid in turbulent motion. *The Quarterly Journal of Mechanics and Applied Mathematics* **7** (1), 83–103.
- CHANDRASEKHAR, S. 1950 The theory of axisymmetric turbulence. *Philosophical Transactions of the Royal Society of London. Series A* **242** (855).
- CHEN, JUN, MENEVEAU, CHARLES & KATZ, JOSEPH 2006 Scale interactions of turbulence subjected to a straining–relaxation–destraining cycle. *Journal of Fluid Mechanics* **562**, 123–150.
- CROW, S. C. 1968 Viscoelastic properties of fine-grained incompressible turbulence. *Journal of Fluid Mechanics* **33** (1), 1–20.
- HUNT, J. C. R. 1973 A theory of turbulent flow round two-dimensional bluff bodies. *Journal of Fluid Mechanics* **61** (4), 625–706.
- HUNT, J. C. R. & CARRUTHERS, D. J. 1990 Rapid distortion theory and the ‘problems’ of turbulence. *Journal of Fluid Mechanics* **212**, 497–532.
- KERSTEIN, ALAN R. 1991 Linear-eddy modelling of turbulent transport. Part 6. Microstructure of diffusive scalar mixing fields. *Journal of Fluid Mechanics* **231**, 361–394.
- KERSTEIN, ALAN R. 1999 One-dimensional turbulence: model formulation and application to homogeneous turbulence, shear flows, and buoyant stratified flows. *Journal of Fluid Mechanics* **392**, 277–334.
- KERSTEIN, ALAN R., ASHURST, WILLIAM T., WUNSCH, SCOTT & NILSEN, VEBJORN 2001 One-dimensional turbulence: vector formulation and application to free shear flows. *Journal of Fluid Mechanics* **447**, 85–109.
- KOLMOGOROV, A. N. 1991 The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *Proceedings of the Royal Society of London. Series A* **434** (1890), 9–13.
- LAUNDER, B. E., REECE, G. J. & RODI, W. 1975 Progress in the development of a reynolds-stress turbulence closure. *Journal of Fluid Mechanics* **68** (3), 537–566.
- LEE, MOON J., PIOMELLI, UGO & REYNOLDS, WILLIAM C. 1986 Useful formulas in the rapid distortion theory of homogeneous turbulence. *Physics of Fluids* **29** (10), 3471–3474.
- LEE, MOON J. & REYNOLDS, WILLIAM C. 1985 Numerical experiments on the structure of homogeneous turbulence. *Tech. Rep. TF-24*. Thermosciences Division, Department of Mechanical Engineering, Stanford University, Stanford, California, prepared under NASA Grant NCC-2-15.

- 969 LIGNELL, DAVID O., KERSTEIN, ALAN R., SUN, GUANGYUAN & MONSON, ELIZABETH I. 2013 Mesh adaption for
970 efficient multiscale implementation of one-dimensional turbulence. *Theoretical and Computational Fluid*
971 *Dynamics* **27** (3), 273–295.
- 972 LINDBORG, E. 1995 Kinematics of homogeneous axisymmetric turbulence. *Journal of Fluid Mechanics* **302**,
973 179–201.
- 974 POPE, STEPHEN B. 2000 *Turbulent Flows*. Cambridge: Cambridge University Press.
- 975 DE SANTANA, LEANDRO D., CHRISTOPHE, JULIEN, SCHRAM, CHRISTOPHE F. & DESMET, WIM 2016 A rapid
976 distortion theory modified turbulence spectra for semi-analytical airfoil noise prediction. *Journal of Sound*
977 *and Vibration* **383**, 349–363.
- 978 STEPHENS, VICTORIA B. & LIGNELL, DAVID O. 2021 One-dimensional turbulence (ODT): Computationally
979 efficient modeling and simulation of turbulent flows. *SoftwareX* **13**, 100641.
- 980 WUNSCH, SCOTT & KERSTEIN, ALAN R. 2001 A model for layer formation in stably stratified turbulence. *Physics*
981 *of Fluids* **13** (3), 702–712.
- 982