frontal plane elevation — ex vivo − in vivo difference, 95% CI (overlap only)

Sternoclavicular — Protraction(+)/retraction(-)

13/4 sh · 4/3 st

K = 6

not fitted yet

Sternoclavicular — Depression(+)/elevation(-)

13/4 sh · 4/3 st

K = 5

not fitted yet

Sternoclavicular — Backwards(+)/forward(-) rot

13/4 sh · 4/3 st

K = 6

not fitted yet

Acromioclavicular — Protraction(+)/retraction(-)

10/4 sh · 1/3 st

K = 4

not fitted yet

Acromioclavicular — Medial(+)/lateral(-) rot

10/4 sh · 1/3 st

K = 7

not fitted yet

Acromioclavicular — Posterior(+)/anterior(-) tilt

10/4 sh · 1/3 st

K = 8

not fitted yet

Scapulothoracic — Protraction(+)/retraction(-)

13/31 sh · 4/5 st

K = 6

not fitted yet

Scapulothoracic — Medial(+)/lateral(-) rot

13/31 sh · 4/5 st

K = 7

not fitted yet

Scapulothoracic — Posterior(+)/anterior(-) tilt

13/31 sh · 4/5 st

K = 7

not fitted yet

Glenohumeral — Plane of elevation

10/23 sh · 1/3 st

K = 7

not fitted yet

Glenohumeral — Elevation(-)/Depression(+)

10/23 sh · 1/3 st

K = 7

not fitted yet

Glenohumeral — Int(+)/ext(-) rotation

10/23 sh · 1/3 st

K = 7

not fitted yet

blue line = EX VIVO − IN VIVO, band = 95% CI · squares along the bottom = band excludes 0. Note the sign: in vivo is the reference, so this is mirrored against v2.
stars = LIKELIHOOD-RATIO TEST of H0 'no ex-vivo departure' against H1, Bonferroni-adjusted across every compared cell on the plate: *** p<0.001, ** p<0.01, * p<0.05, ns = not significant.
ex/in · studies ex/in | K, this cell's own spline df chosen by BIC | mean and max |Δ| in degrees. Curves span the full abscissa — every x any shoulder reaches is fitted, so the thinnest stretches rest on a single shoulder and the 95%