# sagittal plane elevation — ex vivo − in vivo difference, 95% CI (overlap only)

**Sternoclavicular — Protraction(+)/retraction(-)**

13/3 sh · 4/2 st

K = 4

not fitted yet

**Sternoclavicular — Depression(+)/elevation(-)**

13/3 sh · 4/2 st

K = 6

not fitted yet

**Sternoclavicular — Backwards(+)/forward(-) rot**

13/3 sh · 4/2 st

K = 6

not fitted yet

**Acromioclavicular — Protraction(+)/retraction(-)**

10/3 sh · 1/2 st

K = 7

not fitted yet

**Acromioclavicular — Medial(+)/lateral(-) rot**

10/3 sh · 1/2 st

K = 3

not fitted yet

**Acromioclavicular — Posterior(+)/anterior(-) tilt**

10/3 sh · 1/2 st

K = 5

not fitted yet

**Scapulothoracic — Protraction(+)/retraction(-)**

13/25 sh · 4/5 st

K = 7

not fitted yet

**Scapulothoracic — Medial(+)/lateral(-) rot**

13/25 sh · 4/5 st

K = 6

not fitted yet

**Scapulothoracic — Posterior(+)/anterior(-) tilt**

13/25 sh · 4/5 st

K = 8

not fitted yet

**Glenohumeral — Plane of elevation**

10/24 sh · 1/4 st

K = 10

not fitted yet

**Glenohumeral — Elevation(-)/Depression(+)**

10/24 sh · 1/4 st

K = 6

not fitted yet

**Glenohumeral — Int(+)/ext(-) rotation**

10/24 sh · 1/4 st

K = 10

not fitted yet

blue line = EX VIVO − IN VIVO, band = 95% CI · squares along the bottom = band excludes 0. Note the sign: in vivo is the reference, so this is mirrored against v2.
stars = LIKELIHOOD-RATIO TEST of H0 'no ex-vivo departure' against H1, Bonferroni-adjusted across every compared cell on the plate: *** p<0.001, ** p<0.01, * p<0.05, ns = not significant.
ex/in · studies ex/in | K, this cell's own spline df chosen by BIC | mean and max |Δ| in degrees. Curves span the full abscissa — every x any shoulder reaches is fitted, so the thinnest stretches rest on a single shoulder and the 95%