

scapular plane elevation — ex vivo – in vivo difference, 95% CI (overlap only)

Sternoclavicular — Protraction(+)/retraction(-)	Sternoclavicular — Depression(+)/elevation(-)	Sternoclavicular — Backwards(+)/forward(-) rot
single group · K = 9	single group · K = 4	single group · K = 3
not modelled	not modelled	not modelled
single group · K = 11	single group · K = 11	single group · K = 11
single group · K = 18	single group · K = 10	single group · K = 14

blue line = EX VIVO – IN VIVO, band = 95% CI · squares along the bottom = band excludes 0. Note the sign: in vivo is the reference, so this is mirrored against v2.
stars = LIKELIHOOD-RATIO TEST of H0 'no ex-vivo departure' against H1, Bonferroni-adjusted across every compared cell on the plate: *** p<0.001, ** p<0.01, * p<0.05, ns = not significant.

ex/in · studies ex/in | K, this cell's own spline df chosen by BIC | mean and max |Δ| in degrees. Curves span the full abscissa — every x any shoulder reaches is fitted, so the thinnest stretches rest on a single shoulder and the 95% CI is the widest.

ex/in · studies ex/in | K, this cell's own spline df chosen by BIC | mean and max $|\Delta|$ in degrees. Curves span the full abscissa — every x any shoulder reaches is fitted, so the thinnest stretches rest on a single shoulder and the 95%