

internal-external rotation 90 degree-abducted — random spline, per-cell df by BIC, D diagonal

Sternoclavicular — Protraction(+)/retraction(-)

not modelled

Sternoclavicular — Depression(+)/elevation(-)

not modelled

Sternoclavicular — Backwards(+)/forward(-) rot

not modelled

Acromioclavicular — Protraction(+)/retraction(-)

not modelled


Acromioclavicular — Medial(+)/lateral(-) rot

not modelled


Acromioclavicular — Posterior(+)/anterior(-) tilt

not modelled


Scapulothoracic — Protraction(+)/retraction(-)




Scapulothoracic — Medial(+)/lateral(-) rot




Scapulothoracic — Posterior(+)/anterior(-) tilt




Glenohumeral — Plane of elevation



Glenohumeral — Elevation(-)/Depression(+)



Glenohumeral — Int(+)/ext(-) rotation



orange = ex vivo, green = in vivo · line = population curve, band = 95% CI. No knots are drawn.
stars = LIKELIHOOD-RATIO TEST of H0 'no ex-vivo departure' against H1, Bonferroni-adjusted across every compared cell on the plate: *** p<0.001, ** p<0.01, * p<0.05, ns = not significant.
ex/in · studies ex/in | K, this cell's own spline df chosen by BIC | mean and max |Δ| in degrees. Curves span the full abscissa — every x any shoulder reaches is fitted, so the thinnest stretches rest on a single shoulder and the 95%