




.....Olympiad Training for Individual Study

Elementary Geometry (Version W)

EVAN CHEN 《陳誼廷》

7 September 2023
BGW-ELEMGEOM



OTIS, © Evan Chen, internal use only. Artwork contributed by Rishabh Mahale.

§1 Lecture notes

This is some practice with the low-tech end of olympiad geometry, i.e. problems whose *canonical*¹ synthetic solutions essentially only use:

- Angle chasing
- Power of a point
- Homothety

Example 1.1 (USEMO 2021/4, Sayandeep Shee)

Let ABC be a triangle with circumcircle ω , and let X be the reflection of A in B . Line CX meets ω again at D . Lines BD and AC meet at E , and lines AD and BC meet at F . Let M and N denote the midpoints of AB and AC .

Can line EF share a point with the circumcircle of triangle AMN ?

21USEM04

Walkthrough. There are lots of different approaches. We'll give the most classical one presented by the author, but there are plenty of other things that could work too!

- Let P be the midpoint of \overline{AD} . Where else does P lie?
- Show that $FB^2 = FP \cdot FA$.
- Show that $EB^2 = EN \cdot EA$.
- Using radical axis with a circle of radius zero at B , prove that line EF is disjoint from (AMN) .

Example 1.2 (JMO 2018/3, Ray Li)

Let $ABCD$ be a quadrilateral inscribed in circle ω with $\overline{AC} \perp \overline{BD}$. Let E and F be the reflections of D over \overline{BA} and \overline{BC} , respectively, and let P be the intersection of \overline{BD} and \overline{EF} . Suppose that the circumcircles of EPD and FPD meet ω at Q and R different from D . Show that $EQ = FR$.

18JMO3

Walkthrough. Most of this problem is about realizing where the points P , Q , R are.

- Using what you know about the Simson line, figure out where point P is.
- Determine the circumcenters of $\triangle EPD$ and $\triangle FPD$.
- Figure out where the points Q and R are.
- Finish the problem.

¹I emphasize “canonical” since most official solutions can be rewritten in a way that avoids inversion, projective, etc. These problems are the ones for which even students who know this machinery would feasibly find the low-tech solution first.

Example 1.3 (TSTST 2017/5, Ray Li)

Let ABC be a triangle with incenter I . Let D be a point on side BC and let ω_B and ω_C be the incircles of $\triangle ABD$ and $\triangle ACD$, respectively. Suppose that ω_B and ω_C are tangent to segment BC at points E and F , respectively. Let P be the intersection of segment AD with the line joining the centers of ω_B and ω_C . Let X be the intersection point of lines BI and CP and let Y be the intersection point of lines CI and BP . Prove that lines EX and FY meet on the incircle of $\triangle ABC$.

17TSTST5

Walkthrough. Let ω denote the incircle of $\triangle ABC$.

- Identify the point $Z = \overline{EX} \cap \overline{FY}$ in a good diagram. (This was worth a point! Despite this, many contestants were unable to find it.)
- Consider the positive homothety sending ω to ω_C . Determine its center.
- Consider the negative homothety sending ω_C to ω_B . Determine its center.
- The composition of the previous two homotheties in (b) and (c) is a negative homothety sending ω to ω_B . Determine with proof the center of this homothety.
This is not as simple as the previous two parts; you will need to *use* (b) and (c) to do this part, as well as the simple observation that the center should lie on the $\angle B$ bisector.
- Conclude that \overline{FY} passes through the point you claimed in (a).

Experts may notice that this walkthrough gives what is essentially a proof of Monge d'Alembert theorem.

Example 1.4 (TSTST 2016/2, Evan Chen)

Let ABC be a scalene triangle with orthocenter H and circumcenter O and denote by M, N the midpoints of $\overline{AH}, \overline{BC}$. Suppose the circle γ with diameter \overline{AH} meets the circumcircle of ABC at $G \neq A$, and meets line \overline{AN} at $Q \neq A$. The tangent to γ at G meets line OM at P . Show that the circumcircles of $\triangle GNQ$ and $\triangle MBC$ intersect on \overline{PN} .

16TSTST2

Walkthrough. Let E and F be the feet of the altitudes.

- Show that P is really just the intersection of the tangents to γ at A and G (and thus the line \overline{OM} is just a distraction).
- Show that lines $\overline{AG}, \overline{EF}, \overline{BC}$ are concurrent, say at R .
- Prove that $(PAMG), (MBC), (MFDNE)$ are concurrent at a point $T \neq M$.
- Show that $T = \overline{PN} \cap \overline{MR}$.
- Show that $R \in \overline{HQ}$.
- Show that R, G, T, Q, N are concyclic, completing the proof.

§2 Practice problems

Instructions: Solve [36♣]. If you have time, solve [42♣]. Problems with red weights are mandatory.

What kind of idiot would risk his life just to make a fake brother?

Winry in *Full Metal Alchemist: Brotherhood*

♠ REIM

[2♣] **Problem 1** (Reim's theorem). We say two lines ℓ_1, ℓ_2 are *antiparallel* with respect to m_1 and m_2 , if the four points $\ell_i \cap m_j$ are the vertices of a cyclic quadrilateral.

Fix two lines m_1 and m_2 . Prove that if ℓ_1 is antiparallel to ℓ_2 and ℓ_2 is antiparallel to ℓ_3 , then lines ℓ_1 and ℓ_3 are either parallel or coincide.

♠ AOA60

[2♣] **Problem 2** (Added by Karn Chutinan). Let ABC be a triangle with circumcenter O and orthocenter H . Prove that $AO = AH$ if and only if $\angle BAC = 60^\circ$.

♠ O8IMO1

[2♣] **Problem 3** (IMO 2008/1). Let H be the orthocenter of an acute-angled triangle ABC . The circle Γ_A centered at the midpoint of \overline{BC} and passing through H intersects the sideline BC at points A_1 and A_2 . Similarly, define the points B_1, B_2, C_1 , and C_2 . Prove that six points $A_1, A_2, B_1, B_2, C_1, C_2$ are concyclic.

♠ 15APMO1

[2♣] **Problem 4** (APMO 2015/1). Let ABC be a triangle, and let D be a point on side BC . A line through D intersects side AB at X and ray AC at Y . The circumcircle of triangle BXD intersects the circumcircle ω of triangle ABC again at point Z distinct from point B . The lines ZD and ZY intersect ω again at V and W respectively. Prove that $AB = VW$.

♠ 13EGMO1

[2♣] **Problem 5** (EGMO 2013/1). The side BC of the triangle ABC is extended beyond C to D so that $CD = BC$. The side CA is extended beyond A to E so that $AE = 2CA$. Prove that if $AD = BE$ then the triangle ABC is right-angled.

♠ 15UK

[2♣] **Problem 6** (UK 2015). Let $ABCD$ be a cyclic quadrilateral. Let F be the midpoint of the arc AB of its circumcircle which does not contain C or D . Let the lines DF and AC meet at P and the lines CF and BD meet at Q . Prove that the lines PQ and AB are parallel.

♠ 20BAMO4

[3♣] **Problem 7** (BAMO 2020/4). Let ABC be a triangle and let M be any point on segment BC . Denote by O_B and O_C the circumcenters of $\triangle ABM$ and $\triangle ACM$, respectively. Suppose ω is a circle whose center lies on \overline{BC} and passes through A and M . Lines MO_B and MO_C meet ω again at K and L , respectively. Prove that lines BK and CL meet on ω .

♠ 16HMIC2

[3♣] **Problem 8** (HMIC 2016/2). Let ABC be an acute triangle with circumcenter O , orthocenter H , and circumcircle Ω . Let M be the midpoint of BH and N the midpoint of CH . Assume the points M, N, O, H are distinct and lie on a circle ω . Prove that the circles ω and Ω are internally tangent to each other.

♠ 15HMMT4

[3♣] **Problem 9** (HMMT 2015 T4). Convex quadrilateral $ABCD$ with $BC = CD$ is inscribed in circle Ω ; the diagonals of $ABCD$ meet at X . Suppose $AD < AB$, the circumcircle of triangle BCX intersects segment AB at a point $Y \neq B$, and ray \overrightarrow{CY} meets Ω again at a point $Z \neq C$. Prove that ray \overrightarrow{DY} bisects angle ZDB .

🚩 04IM01

[3♣] **Problem 10 (IMO 2004/1)**. Let ABC be an acute-angled triangle with $AB \neq AC$. The circle with diameter BC intersects the sides AB and AC at M and N respectively. Denote by O the midpoint of the side BC . The bisectors of the angles $\angle BAC$ and $\angle MON$ intersect at R . Prove that the circumcircles of the triangles BMR and CNR have a common point lying on the side BC .

🚩 98AM02

[2♣] **Problem 11 (USAMO 1998/2)**. Let C_1 and C_2 be concentric circles, with C_2 in the interior of C_1 . From a point A on C_1 one draws the tangent AB to C_2 ($B \in C_2$). Let C be the second point of intersection of ray AB and C_1 , and let D be the midpoint of \overline{AB} . A line passing through A intersects C_2 at E and F in such a way that the perpendicular bisectors of \overline{DE} and \overline{CF} intersect at a point M on line AB . Find, with proof, the ratio AM/MC .

🚩 12GER3

[3♣] **Problem 12 (Germany 2012/3, added by Joel Gerlach)**. Triangle ABC has $\angle A = 120^\circ$. Let H and O denote the orthocenter and circumcenter of $\triangle ABC$. Prove that

$$|AB| + |AC| = |OH|.$$

🚩 REMOROV

[3♣] **Problem 13**. Let ABC be an acute triangle with circumcircle Γ and intouch triangle DEF . A circle is drawn tangent to \overline{BC} at D and to minor arc BC of Γ at X . Define Y and Z similarly. Prove that lines \overline{DX} , \overline{EY} , \overline{FZ} are concurrent.

🚩 23AM01

[3♣] **Problem 14 (USAMO 2023/1)**. In an acute triangle ABC , let M be the midpoint of \overline{BC} . Let P be the foot of the perpendicular from C to AM . Suppose that the circumcircle of triangle ABP intersects line BC at two distinct points B and Q . Let N be the midpoint of \overline{AQ} . Prove that $NB = NC$.

🚩 06SLG2

[3♣] **Required Problem 15 (Shortlist 2006 G2)**. Let $ABCD$ be a trapezoid with parallel sides $AB > CD$. Points K and L lie on the line segments AB and CD , respectively, so that $AK/KB = DL/LC$. Suppose points P and Q on segment KL obey

$$\angle APB = \angle BCD \quad \text{and} \quad \angle CQD = \angle ABC.$$

Prove that P, Q, B, C are concyclic.

🚩 17EGM01

[3♣] **Problem 16 (EGMO 2017/1)**. Let $ABCD$ be a convex quadrilateral with $\angle DAB = \angle BCD = 90^\circ$ and $\angle ABC > \angle CDA$. Let Q and R be points on segments BC and CD , respectively, such that line QR intersects lines AB and AD at points P and S , respectively. It is given that $PQ = RS$. Let the midpoint of BD be M and the midpoint of QR be N . Prove that the points M, N, A and C lie on a circle.

🚩 20IGOB2

[3♣] **Problem 17 (Iran Geo Olympiad 2020, added by Pedro Rosalba)**. Let ABC be an isosceles triangle ($AB = AC$) with its circumcenter O . Point N is the midpoint of the segment BC and point M is the reflection of the point N with respect to the side AC . Suppose that T is a point so that $ANBT$ is a rectangle. Prove that $\angle OMT = \frac{1}{2}\angle BAC$.

🚩 22USEM04

[9♣] **Required Problem 18 (USEMO 2022/4)**. Let $ABCD$ be a cyclic quadrilateral whose opposite sides are not parallel. Suppose points P, Q, R, S lie in the interiors of segments AB, BC, CD, DA , respectively, such that

$$\angle PDA = \angle PCB, \quad \angle QAB = \angle QDC, \quad \angle RBC = \angle RAD, \quad \text{and} \quad \angle SCD = \angle SBA.$$

Let \overline{AQ} intersect \overline{BS} at X , and \overline{DQ} intersect \overline{CS} at Y . Prove that lines \overline{PR} and \overline{XY} are either parallel or coincide.

20ELMO4

[5♣] **Problem 19** (ELMO 2020/4). Let ABC be an acute scalene triangle ABC with orthocenter H . Let D be the foot of the A -altitude, and let M be the midpoint of \overline{BC} . Let D' be the reflection of D over M . Point P is chosen on line $D'H$ such that $\overline{AP} \parallel \overline{BC}$. Suppose the circumcircles of $\triangle AHP$ and $\triangle BHC$ meet again at a point G other than H . Prove that $\angle MHG = 90^\circ$.

97IMO2

[3♣] **Problem 20** (IMO 1997/2). It is known that $\angle BAC$ is the smallest angle in the triangle ABC . The points B and C divide the circumcircle of the triangle into two arcs. Let U be an interior point of the arc between B and C which does not contain A . The perpendicular bisectors of AB and AC meet the line AU at V and W , respectively. The lines BV and CW meet at T .

Show that $AU = TB + TC$.

15INDTST1

[5♣] **Problem 21** (India TST 2015/1). Diagonals \overline{AC} and \overline{BD} of convex quadrilateral $ABCD$ meet at P . Prove that the incenters of the triangles $\triangle PAB$, $\triangle PBC$, $\triangle PCD$, $\triangle PDA$ are concyclic if and only if their P -excenters are also concyclic.

23TSTST1

[5♣] **Required Problem 22** (TSTST 2023/1). Let ABC be a triangle with centroid G . Points R and S are chosen on rays GB and GC , respectively, such that

$$\angle ABS = \angle ACR = 180^\circ - \angle BGC.$$

Prove that $\angle RAS + \angle BAC = \angle BGC$.

17SLG3

[9♣] **Problem 23** (Shortlist 2017 G3). Let ABC be an acute triangle with orthocenter H and circumcenter O . Let $P = \overline{BH} \cap \overline{AO}$ and $Q = \overline{CH} \cap \overline{AO}$. Prove that the circumcenter of $\triangle HPQ$ lies on the A -median.

16EGMO4

[5♣] **Required Problem 24** (EGMO 2016/4). Two circles ω_1 and ω_2 , of equal radius intersect at different points X_1 and X_2 . Consider a circle ω externally tangent to ω_1 at T_1 and internally tangent to ω_2 at point T_2 . Prove that lines X_1T_1 and X_2T_2 intersect at a point lying on ω .

99AMO6

[3♣] **Problem 25** (USAMO 1999/6). Let $ABCD$ be an isosceles trapezoid with $AB \parallel CD$. The inscribed circle ω of triangle BCD meets CD at E . Let F be a point on the (internal) angle bisector of $\angle DAC$ such that $EF \perp CD$. Let the circumscribed circle of triangle ACF meet line CD at C and G . Prove that the triangle AFG is isosceles.

[1♣] **Mini Survey**. Fill out feedback on the OTIS-WEB portal when submitting this problem set. Any thoughts on problems (e.g. especially nice, instructive, easy, etc.) or overall comments on the unit are welcome.

In addition, if you have any suggestions for problems to add, or want to write hints for one problem you really liked, please do so in the ARCH system!

The maximum number of [♣] for this unit is [89♣], including the mini-survey.


§3 Solutions to the walkthroughs

§3.1 Solution 1.1, USEMO 2021/4, Sayandeep Shee

The answer is no, they never intersect.

¶ **Classical solution, by author** Let P denote the midpoint of \overline{AD} , which

- lies on \overline{BN} , since $\overline{BN} \parallel \overline{CX}$; and
- lies on (AMN) , since it's homothetic to (ABC) through A with factor $\frac{1}{2}$.



Now, note that

$$\angle FBP = \angle CBN = \angle BCD = \angle BAD = \angle BAF \implies FB^2 = FP \cdot FA$$

$$\angle EBN = \angle EDC = \angle BDC = \angle BAC = \angle BAE \implies EB^2 = EN \cdot EA.$$

This means that line EF is the radical axis of the circle centered at B with radius zero, and the circumcircle of triangle AMN . Since B obviously lies outside (AMN) , the disjointness conclusion follows.

¶ **Projective solution, by Ankit Bisain** In this approach we are still going to prove that \overline{EF} is the radical axis of (AMN) and the circle of radius zero at B , but we are not going to use the point P , or even points E and F .

Instead, let $Y = \overline{EF} \cap \overline{AB}$, which by Brokard's theorem on $ABDC$ satisfies $(AB; XY) = -1$. Since $XB = XA$, it follows that $AY : YB = 2$. From here it is straightforward to verify that

$$YB^2 = \frac{1}{9}AB^2 = YM \cdot YA.$$

Thus Y lies on the radical axis.

Finally, by Brokard's theorem again, if O is the center of ω then $\overline{OX} \perp \overline{EF}$. Taking a homothety with scale factor 2 at A , it follows that the line through B and the center of (AMN) is perpendicular to \overline{EF} .

Since \overline{EF} contains Y , it now follows that \overline{EF} is the radical axis, as claimed.

¶ **Solution with inversion, projective, and Cartesian coordinates, by Ankan Bhat-tacharya** In what follows, let O be the center of ω . Note that Brokard's theorem gives that \overline{EF} is the polar of X .

Note that since none of E, F, X are points at infinity, O is different from all three.

We consider inversion in ω to eliminate the polar:

- The circumcircle of $\triangle AMN$, i.e. the circle with diameter \overline{AO} , is sent to the line ℓ tangent to ω at A .
- The line EF , as the polar of X , is sent to the circle with diameter \overline{OX} . (It is indeed a circle, because O does not lie on line EF .)

Thus, if the posed question is true, then we see that ℓ intersects (OX) . We claim this is impossible.

Establish Cartesian coordinates with $A = (0, 0)$ and $O = (2, 0)$, so ℓ is the y -axis. Let T be the center of (OX) : the midpoint of \overline{OX} . Observe:

- B lies on the circle with center $(2, 0)$ and radius 2.
- X lies on the circle with center $(4, 0)$ and radius 4.
- T lies on the circle with center $(3, 0)$ and radius 2.

Thus, let the coordinates of T be (x, y) , with $(x - 3)^2 + y^2 = 4$. The intersection of ℓ and (OX) being nonempty is equivalent to

$$\begin{aligned} d(T, \ell)^2 &\leq OT^2 \\ \iff x^2 &\leq (x - 2)^2 + y^2 \\ \iff x^2 &\leq (x - 2)^2 + [4 - (x - 3)^2] \\ \iff (x - 1)^2 &\leq 0, \end{aligned}$$

or $x = 1$ (which forces $y = 0$); i.e. $T = (1, 0)$. However, this forces


$$B = (0, 0) = A,$$

which is not permitted. Thus, line ℓ cannot share a point with (OX) , and so line EF cannot share a point with (AMN) .

§3.2 Solution 1.2, JMO 2018/3, Ray Li

Most of this problem is about realizing where the points P, Q, R are.

¶ **First solution (Evan Chen)** Let X, Y , be the feet from D to $\overline{BA}, \overline{BC}$, and let $Z = \overline{BD} \cap \overline{AC}$. By Simson theorem, the points X, Y, Z are collinear. Consequently, the point P is the reflection of D over Z , and so we conclude P is the orthocenter of $\triangle ABC$.



Suppose now we extend ray CP to meet ω again at Q' . Then \overline{BA} is the perpendicular bisector of both $\overline{PQ'}$ and \overline{DE} ; consequently, $PQ'ED$ is an isosceles trapezoid. In particular, it is cyclic, and so $Q' = Q$. In the same way R is the second intersection of ray \overline{AP} with ω .


Now, because of the two isosceles trapezoids we have found, we conclude

$$EQ = PD = FR$$

as desired.

Remark. Alternatively, after identifying P , one can note \overline{BQE} and \overline{BRF} are collinear. Since $BE = BD = BF$, upon noticing $BQ = BP = BR$ we are also done.

¶ **Second solution (Danielle Wang)** Here is a solution which does not identify the point P at all. We know that $BE = BD = BF$, by construction.



Claim — The points B, Q, E are collinear. Similarly the points B, R, F are collinear.

Proof. Work directed modulo 180° . Let Q' be the intersection of \overline{BE} with $(ABCD)$. Let $\alpha = \angle DEB = \angle BDE$ and $\beta = \angle BFD = \angle FDB$.

Observe that $BE = BD = BF$, so B is the circumcenter of $\triangle DEF$. Thus, $\angle DEP = \angle DEF = 90^\circ - \beta$. Then


$$\begin{aligned}\angle DPE &= \angle DEP + \angle PDE = (90^\circ - \beta) + \alpha \\ &= \alpha - \beta + 90^\circ \\ \angle DQ'B &= \angle DCB = \angle DCA + \angle ACB \\ &= \angle DBA - (90^\circ - \angle DBC) = -(90^\circ - \alpha) - (90^\circ - (90^\circ - \beta)) \\ &= \alpha - \beta + 90^\circ.\end{aligned}$$

Thus Q' lies on the desired circle, so $Q' = Q$. \square

Now, by power of a point we have $BQ \cdot BE = BP \cdot BD = BR \cdot BF$, so $BQ = BP = BR$. Hence $EQ = PD = FR$.

§3.3 Solution 1.3, TSTST 2017/5, Ray Li

¶ First solution (homothety) Let Z be the diametrically opposite point on the incircle. We claim this is the desired intersection.



Note that:

- P is the insimilicenter of ω_B and ω_C
- C is the exsimilicenter of ω and ω_C .

Thus by Monge theorem, the insimilicenter of ω_B and ω lies on line CP .

This insimilicenter should also lie on the line joining the centers of ω and ω_B , which is \overline{BI} , hence it coincides with the point X . So $X \in \overline{EZ}$ as desired.

¶ Second solution (harmonic) Let $T = \overline{BI_C} \cap \overline{BC}$, and W the foot from I to \overline{BC} . Define $Z = \overline{FY} \cap \overline{IW}$. Because $\angle I_B D I_C = 90^\circ$, we have

$$-1 = (I_B I_C; PT) \stackrel{B}{=} (I I_C; YC) \stackrel{F}{=} (I \infty; ZW)$$

So I is the midpoint of \overline{ZW} as desired.

¶ **Third solution (outline, barycentric, Andrew Gu)** Let $AD = t$, $BD = x$, $CD = y$, so $a = x + y$ and by Stewart's theorem we have

$$(x + y)(xy + t^2) = b^2x + c^2y. \quad (1)$$

We then have $D = (0 : y : x)$ and so

$$\overline{AI_B} \cap \overline{BC} = \left(0 : y + \frac{tx}{c+t} : \frac{cx}{c+t} \right)$$

hence intersection with BI gives

$$I_B = (ax : cy + at : cx).$$

Similarly,

$$I_C = (ay : by : bx + at).$$

Then, we can compute

$$P = (2axy : y(at + bx + cy) : x(at + bx + cy))$$

since $P \in \overline{I_B I_C}$, and clearly $P \in \overline{AD}$. Intersection now gives

$$X = (2ax : at + bx + cy : 2cx)$$

$$Y = (2ay : 2by : at + bx + cy).$$

Finally, we have $BE = \frac{1}{2}(c + x - t)$, and similarly for CF . Now if we reflect $D = (0, \frac{s-c}{a}, \frac{s-b}{a})$ over $I = (\frac{a}{2s}, \frac{b}{2s}, \frac{c}{2s})$, we get the antipode

$$Q := (4a^2 : -a^2 + 2ab - b^2 + c^2 : -a^2 + 2ac - c^2 + b^2).$$

We may then check Q lies on each of lines EX and FY (by checking $\det(Q, E, X) = 0$ using the equation (1)).


§3.4 Solution 1.4, TSTST 2016/2, Evan Chen

We present two solutions, one using essentially only power of a point, and the other more involved.


¶ **First solution (found by contestants)** Denote by $\triangle DEF$ the orthic triangle. Observe \overline{PA} and \overline{PG} are tangents to γ , since \overline{OM} is the perpendicular bisector of \overline{AG} . Also note that \overline{AG} , \overline{EF} , \overline{BC} are concurrent at some point R by radical axis on (ABC) , γ , (BC) .

Now, consider circles $(PAGM)$, $(MFDNE)$, and (MBC) . They intersect at M but have radical center R , so are coaxial; assume they meet again at $T \in \overline{RM}$, say. Then $\angle PTM$ and $\angle MTN$ are both right angles, hence T lies on \overline{PN} .

Finally H is the orthocenter of $\triangle ARN$, and thus the circle with diameter \overline{RN} passes through G , Q , N .



¶ **Alternate solution (by proposer)** Let L be diametrically opposite A on the circumcircle. Denote by $\triangle DEF$ the orthic triangle. Let $X = \overline{AH} \cap \overline{EF}$. Finally, let T be the second intersection of $(MFDNE)$ and (MBC) .



We begin with a few easy observations. First, points H, G, N, L are collinear and $\angle AGL = 90^\circ$. Also, Q is the foot from H to \overline{AN} . Consequently, lines AG, EF, HQ, BC, TM concur at a point R (radical axis). Moreover, we already know $\angle MTN = 90^\circ$. This implies T lies on the circle with diameter \overline{RN} , which is exactly the circumcircle of $\triangle GQN$.

Note by Brokard's Theorem on $AFHE$, the point X is the orthocenter of $\triangle MBC$. But $\angle MTN = 90^\circ$ already, and N is the midpoint of \overline{BC} . Consequently, points T, X ,

N are collinear.

Finally, we claim P, X, N are collinear, which solves the problem. Note $P = \overline{GG} \cap \overline{AA}$. Set $K = \overline{HNL} \cap \overline{AP}$. Then by noting

$$-1 = (D, X; A, H) \stackrel{N}{=} (\infty, \overline{NX} \cap \overline{AK}; A, K)$$

we see that \overline{NX} bisects segment \overline{AK} , as desired. (A more projective finish is to show that \overline{PXN} is the polar of R to γ).

Remark. The original problem proposal reads as follows:

Let ABC be a triangle with orthocenter H and circumcenter O and denote by M, N the midpoints of $\overline{AH}, \overline{BC}$. Suppose ray OM meets the line parallel to \overline{BC} through A at P . Prove that the line through the circumcenter of $\triangle MBC$ and the midpoint of \overline{OH} is parallel to \overline{NP} .

The points G and Q were added to the picture later to prevent the problem from being immediate by coordinates.