Please check the examination details	below before ente	ering your candidate information
Candidate surname		Other names
Centre Number Candidate	Number	
Pearson Edexcel Inte	ernation	al Advanced Level
Time 1 hour 30 minutes	Paper reference	WME01/01
Mathematics International Advanced Mechanics M1	Subsidiar	y/Advanced Level
You must have: Mathematical Formulae and Statist	tical Tables (Ye	Total Marks (Pllow), calculator

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear.
 Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \,\mathrm{m \, s^{-2}}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶



P71403A
©2022 Pearson Education Ltd.



1.	1. Two particles, P and Q , are moving towards each other in opposite directions along the same straight line when they collide directly. Immediately before the collision the speed of Q is $2u$. The mass of Q is $3m$ and the magnitude of the impulse exerted by P on Q in the collision is $4mu$.					
	Find					
	(a) the speed of Q immediately after the collision, (3)					
	(b) the direction of motion of Q immediately after the collision. (1)					
_						

	Leave
Question 1 continued	blank
Question 1 continued	
	Q1
(Total 4 marks)	
(Total 4 illaiks)	



2.	A motorbike is moving with constant acceleration along a straight horizontal road.
	The motorbike passes a point P and 10 seconds later passes a point Q .
	The speed of the motorbike as it passes Q is $28 \mathrm{ms^{-1}}$
	Given that $PQ = 220 \mathrm{m}$,
	(a) find the acceleration of the motorbike,
	(3)
	(b) find the distance travelled by the motorbike during the fifth second after passing P (4)

estion 2 continued	action 2 continued	
	estion 2 continued	



Question 2 continued	

Question 2 continued	blank
	Q2
(Total 7 marks)	



3.	A tractor of mass 6 tonnes is dragging a large block of mass 2 tonnes along rough
	horizontal ground. The cable connecting the tractor to the block is horizontal and parallel
	to the direction of motion.

The cable is modelled as being light and inextensible.

The driving force of the tractor is $7400 \,\mathrm{N}$ and the resistance to the motion of the tractor is $200 \,\mathrm{N}$. The resistance to the motion of the block is R newtons, where R is a constant.

Given that the tension in the cable is 6000 N and the tractor is accelerating,

(a)	find	the	value	of R .
-----	------	-----	-------	----------

(6)

(b)	State how	you have	used the	fact that	t the cable	e is modelled	l as being	inextensible
-----	-----------	----------	----------	-----------	-------------	---------------	------------	--------------

-	4	1
- 1	П	۱ ۱
•	Л	L



Question 3 continued	Leave
	02
	Q3
(Total 7 marks)	



4.




Figure 1

A small block of mass 5 kg lies at rest on a rough horizontal plane.

The coefficient of friction between the block and the plane is $\frac{3}{7}$

A force of magnitude P newtons is applied to the block in a direction which makes an angle of 30° with the plane, as shown in Figure 1.

The block is modelled as a particle.

Given that P = 14

(a) find the magnitude of the frictional force exerted on the block by the plane and describe what happens to the block, justifying your answer.

(6)

The value of *P* is now changed so that the block is on the point of slipping along the plane.

(b) Find the value of P

(6)

	Leave blank
Question 4 continued	Olalik



uestion 4 contin	rued		

Question 4 continued	Leave
Question reminieu	
	Q4
(Total 12 marks)	
(22	-






Figure 2

A uniform rod AB has length 5 m and mass 5 kg. The rod rests in equilibrium in a horizontal position on two supports C and D, where AC = 1 m and DB = 2 m, as shown in Figure 2.

A particle of mass $10 \,\mathrm{kg}$ is placed on the rod at A and a particle of mass $M \,\mathrm{kg}$ is placed on the rod at B. The rod remains horizontal and in equilibrium.

(a) Find, in terms of M, the magnitude of the reaction on the rod at C.

(3)

(b) Find, in terms of M, the magnitude of the reaction on the rod at D.

(3)

(c) Hence, or otherwise, find the range of possible values of M.

(3)

	Leave blank
Question 5 continued	



Question 5 continued	

Question 5 continued	blank
	Q5
(Total 9 marks)	



Leave

6.	A particle P	is moving	with constant	acceleration.

At time t = 1 second, P has velocity $(-\mathbf{i} + 4\mathbf{j}) \,\mathrm{m \, s}^{-1}$

At time t = 4 seconds, P has velocity $(5\mathbf{i} - 8\mathbf{j}) \,\mathrm{m \, s}^{-1}$

Find the speed of P at time t = 3.5 seconds.

(6)		



Question 6 continued	Leave
	Q6
(Total 6 marks)	



7. Two small children, Ajaz and Beth, are running a 100 m race along a straight horizontal track.

They both start from rest, leaving the start line at the same time.

Ajaz accelerates at $0.8\,\mathrm{m\,s^{-2}}$ up to a speed of $4\,\mathrm{m\,s^{-1}}$ and then maintains this speed until he crosses the finish line.

Beth accelerates at $1\,\mathrm{m\,s^{-2}}$ for T seconds and then maintains a constant speed until she crosses the finish line.

Ajaz and Beth cross the finish line at the same time.

(a) Sketch, on the same axes, a speed-time graph for each child, from the instant when they leave the start line to the instant when they cross the finish line.

(3)

(b) Find the time taken by Ajaz to complete the race.

(4)

(c) Find the value of T

(4)

(d) Find the difference in the speeds of the two children as they cross the finish line.

(2)

Question 7 continued	Leave blank



Question 7 continued	

Question 7 continued	Leave blank
	Q7
(Total 13 marks)	



8. [In this question, **i** and **j** are horizontal unit vectors directed due east and due north respectively and position vectors are given relative to a fixed origin O.]

Two boats, P and Q, are moving with constant velocities.

The velocity of P is $15i \,\mathrm{m \, s}^{-1}$ and the velocity of Q is $(20i - 20j) \,\mathrm{m \, s}^{-1}$

(a) Find the direction in which Q is travelling, giving your answer as a bearing.

(2)

The boats are modelled as particles.

At time t = 0, P is at the origin O and Q is at the point with position vector 200j m.

At time t seconds, the position vector of P is \mathbf{p} m and the position vector of Q is \mathbf{q} m.

(b) Show that

$$\overrightarrow{PQ} = [5t\mathbf{i} + (200 - 20t)\mathbf{j}] \,\mathrm{m}$$

(5)

(c) Find the bearing of P from Q when t = 10

(2)

(d) Find the distance between P and Q when Q is north east of P

(5)

(e) Find the times when P and Q are 200 m apart.

(3)





	blank
Questian 9 continued	Diank
Question 8 continued	
	1



Question 8 continued

	Leave
	blank
Question 8 continued	



	(Total 17 marks)