Please check the examination det	ails below	before ente	ring your can	didate information
Candidate surname			Other name	es
Pearson Edexcel GCE	Centre	Number		Candidate Number
Monday 13 M	lay	201	9	
Afternoon (Time: 1 hour 30 minu	utes)	Paper Re	eference 6	667/01
Further Pure M Advanced/Advanced S			tics F	P1
You must have: Mathematical Formulae and Sta	ntistical T	ables (Pir	nk)	Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P55807A
©2019 Pearson Education Ltd.



Leave blank $f(z) = z^3 + az^2 + bz - 26$ 1. where a and b are real constants. Given that z = 3i - 2 is a solution of the equation f(z) = 0(a) Use algebra to solve f(z) = 0 completely. **(5)** (b) Find the value of a and the value of b. **(3)**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave blank

Question 1 continued	blank

DO NOT WRITE IN THIS AREA

Question 1 continued	Lea bla
	Q1 al 8 marks)



Leave blank

2. $\mathbf{M} = \begin{pmatrix} 9+k & -3 \\ 4-k & 2 \end{pmatrix}, \text{ where } k \text{ is a constant}$

The triangle T has vertices at the points (2, 1), (7, 1) and (7, 12).

Triangle T is transformed onto triangle T' by the transformation represented by the matrix M.

(a) Find, in terms of k, the coordinates of the vertices of the triangle T'

(3)

Given that the area of triangle T' is 770 square units,

(b) find the two possible values of k.

(4)

DO NOT WRITE IN THIS AREA

Question 2 continued	bla
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_



uestion 2 continued	blank

DO NOT WRITE IN THIS AREA

uestion 2 continued	
	(Total 7 marks)



Leave blank

DO NOT WRITE IN THIS AREA

3.	The rectangular hyperbola H has equation
	xy = 10
	The point $P(5, 2)$ lies on H .
	(a) Using calculus, find an equation of the normal to H at the point P , giving your answer in the form $ax + by + c = 0$, where a , b and c are integers. (5)
	The normal at P meets H again at the point Q .
	(b) Find the coordinates of Q . (5)
	(c) Find the exact area of triangle <i>OPQ</i> , where <i>O</i> is the origin. (4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

0	blank
Question 3 continued	



Leave

DO NOT WRITE IN THIS AREA

Question 3 continued		blank
	_	

DO NOT WRITE IN THIS AREA

estion 3 continued	
	(Total 14 marks)



Leave

- **4.** (i) $f(x) = 3x^2 \frac{1}{2\sqrt{x}} 5x, \quad x > 0$
 - (a) Show that the equation f(x) = 0 has a root α in the interval [1, 2].
 - (b) Find f'(x). (2)
 - (c) Using $x_0 = 2$ as a first approximation to α , apply the Newton-Raphson procedure once to f(x) to find a second approximation to α , giving your answer to 2 decimal places.

(2)

(2)

(ii)
$$g(\theta) = 3\theta + 6 - \tan\left(\frac{\theta}{3}\right), \quad -\pi < \theta < 0$$

The equation $g(\theta) = 0$ has a root β in the interval [-3, -2]. Use linear interpolation once on the interval [-3, -2] to find an approximation for β , giving your answer to 3 decimal places.

(4)

NOT WRITE IN THIS A

DO NOT WRITE IN THIS AREA

estion 4 continued	



Leave blank

5. (a) Use the standard results for $\sum_{r=1}^{n} r^2$ and $\sum_{r=1}^{n} r^3$ to show that, for all positive integers n,

$$\sum_{r=1}^{n} r^{2}(4+r) = \frac{1}{12}n(n+1)(an^{2}+bn+c)$$

where a, b and c are integers to be found.

(4)

(b) Hence find the sum of the series

$$5^2 \times 10 + 6^2 \times 11 + 7^2 \times 12 + \dots + 20^2 \times 25$$

(4)

DO NOT WRITE IN THIS AREA

	Leave
	blank
Question 5 continued	



Question 5 continued	Leave blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

uestion 5 continued	
	(Total 8 marks)

Leave

Leave blank

6. $z = \frac{a+2i}{a+5i} \quad a \in \mathbb{R}, a > 0$

Given that the real part of z is $\frac{13}{28}$, find

(a) the value of a,

(4)

(b) arg(z), giving your answer in radians to 2 decimal places,

(3)

(c) the value of zz^* in its simplest form.

(2)

DO NOT WRITE IN THIS AREA

Question 6 continued	blank



DO NOT WRITE IN THIS AREA

	Leave
Question 6 continued	blank

	THIS AREA
24KXX	
-X X X	$\times \times \times \times \times$
-XXX	CKARKO
	AA SEE X
	2
-XX	
	XX 3 AXX
\sim \times \times	
₩	O.
	<u> </u>
	Œ
	E
	3
	WR
	WRI
	WRI
	WR
	TWRIT
	TWR
T E	TWR
	OT WRI
	NOT WRIT
	NOT WRI
	NOT WRI
TEIN AD	NOTWR
TEIC AD	NOTWR
	S NOT WRIT
TEIR ADD	O NOT WRI
TEIC ADE	O NOT WRI
TENC ADEL	O NOT WRI
	DO NOT WRIT
N TEIR AREA	DO NOT WRI
THIS AREA	0
THIS ADEA	DO NOT WRI
TEIIC AREA	DO NOT WRI

uestion 6 continued	
	_
	(Total 9 marks)



Leave blank

7. $\mathbf{P} = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$

(a) Describe fully the single geometrical transformation U represented by the matrix P. (2)

The transformation V, represented by the matrix \mathbf{Q} , is a reflection in the line y = -x

(b) Write down the matrix \mathbf{Q} .

(1)

The transformation U followed by the transformation V is the transformation T.

Transformation T is represented by the matrix \mathbf{R} .

(c) Find the matrix **R**.



(d) Find the exact value of the real constant k for which the transformation T maps the point (1, k) onto itself.

-	- 4	1
	∕∎	- N





DO NOT WRITE IN THIS AREA

uestion 7 continued	
destion / continued	



DO NOT WRITE IN THIS AREA

Question 7 continued	blan

DO NOT WRITE IN THIS AREA

estion 7 continued	



Leave blank

8. (i) Prove by induction, that for $n \in \mathbb{Z}^+$

$$\begin{pmatrix} -3 & -2 \\ 8 & 5 \end{pmatrix}^n = \begin{pmatrix} 1 - 4n & -2n \\ 8n & 1 + 4n \end{pmatrix}$$

(5)

(ii) Prove by induction, that for $n \in \mathbb{Z}^+$

$$\sum_{r=1}^{n} \frac{1}{r(r+1)(r+2)} = \frac{n(n+3)}{4(n+1)(n+2)}$$

(5)

DO NOT WRITE IN THIS AREA

	Leave
	blank
Question 8 continued	



DO NOT WRITE IN THIS AREA

Question 8 continued	blan

DO NOT WRITE IN THIS AREA

uestion 8 continued	
uestion o continueu	

