5-Amino-1MQ

NNMT Inhibitor | Longevity & Metabolic Enhancement

Oral | Injectable

Typical Dose	Route	Cycle	Storage
50-100mg	Oral	8-12 weeks	Room temp
1x daily	N/A - Oral administration	Typical duration	Refrigerated

Did you know? You can suggest edits to improve this peptide information.

Edit

Overview

What is 5-Amino-1MQ?

5-Amino-1MQ is a small molecule NNMT (nicotinamide N-methyltransferase) inhibitor that enhances cellular NAD+ levels and mitochondrial function. Originally developed for metabolic research, it's gaining attention for its potential anti-aging properties and ability to improve energy metabolism at the cellular level.

Key Benefits

Enhanced NAD+ levels, improved mitochondrial function, longevity benefits, better energy metabolism, enhanced cellular repair, improved insulin sensitivity, convenient daily dosing

Mechanism of Action

Orally administered 5-Amino-1MQ inhibits NNMT enzyme systematically, preventing NAD+ degradation and maintaining higher cellular NAD+ levels throughout the body, supporting mitochondrial function and metabolic processes

Quick Start Guide

- Typical Dose: 50-75mg
- How Often: Once daily with food
- How to Take: N/A - Oral administration
- Best Timing: With breakfast or lunch for better absorption
- Effects Timeline: Energy: 1-2 weeks, Metabolic: 4-8 weeks, Longevity: 8-12 weeks
- Storage: Cool, dry place at room temperature
- Cycle Length: 8-12 weeks
- Break Between: 2-4 weeks off between cycles

Pharmacokinetics

Peak: **2 hrs** Half-life: **8 hrs** Cleared: **~1.7 days**

24h 7d 14d 30d



Peak Half-life

Preclinical data only

Community Poll

Question 1 of 10

What is your experience with this compound?

- Currently using
- Used in the past
- Planning to start
- Just researching
- Other (please specify)

Submit Answer

Poll Results

1065 responses

Experience

Currently using	42%
Planning to start	37%
Just researching	17%
Used in the past	4%
Other: What other peppers can be stacked with it	0%

+1 more response

Experience Duration Age Gender Goal Effectiveness Side effects Frequency Recommend Discovery

Research Indications

- **Longevity** — MOST EFFECTIVE

NAD+ Enhancement
Significantly increases cellular NAD+ levels by inhibiting NNMT enzyme, supporting cellular energy production and DNA repair mechanisms essential for healthy aging

Mitochondrial Function
Improves mitochondrial biogenesis and efficiency, enhancing cellular energy production and reducing oxidative stress markers associated with aging

Cellular Repair
Enhanced NAD+ availability supports sirtuins and other longevity pathways involved in DNA repair, cellular maintenance, and stress resistance

- **Metabolism** — EFFECTIVE
- **Weight Loss** — MODERATE

Research Protocols

DISCLAIMER
These are commonly discussed research protocols and not medical advice. Consult a healthcare provider before use.

Goal	Dose	Frequency	Route
Conservative approach	25mg	1x daily	Oral (with food)
Standard starting dose	50mg	1x daily	Oral (with food)
Typical maintenance	75mg	1x daily	Oral (with food)
Maximum dose	100mg	1x daily	Oral (with food)
Split dosing option	50mg	2x daily	Oral (with meals)

TIMING
Take with food to improve absorption and reduce potential GI effects. Consistent daily timing helps maintain steady NAD+ levels. Morning or early afternoon dosing is preferred to avoid potential sleep interference.

Community Insights

1,065 responses

- Effectiveness: 74% report positive results
- Would Recommend: 75% of 74 users
- Common Side Effects: Injection site reaction, Headache, Nausea
- Best Results At: 3-6 months (50% effective)

From 457 users tracking

Weight Change: ↓ -2.8% avg, 67% saw decrease

Sleep Duration: ↑ +0.9 avg, 57% saw increase

Resting Heart Rate: 31% decrease, 51% increase

Peptide Interactions

- NAD+ Precursors (NMN, NR) — SYNERGISTIC
- Resveratrol — SYNERGISTIC
- Metformin — MONITOR COMBINATION
- BPC-157 — COMPATIBLE
- Blood Thinners — UNKNOWN
- Berberine — MONITOR COMBINATION

Was this helpful?

Yes No

Your feedback helps us improve PepPedia

How to Reconstitute

Calculator

IMPORTANT
Always use bacteriostatic water (BAC). Sterile technique is essential.

1. Use pre-made capsules for convenience and accuracy (recommended)
2. If using powder: measure precise dose with milligram scale
3. Take with food to improve absorption and reduce GI effects
4. Maintain consistent daily timing for optimal results
5. Store in cool, dry place away from light and moisture
6. Consider cycling: 8-12 weeks on, 2-4 weeks off

Quality Indicators

- ✓ **White to off-white capsules** — High-quality capsules should be uniform in color and size with clear labeling and batch information
- ✓ **Third-party testing certificates** — Reputable suppliers provide COA with purity analysis (should be >98% pure) and heavy metal testing
- ✓ **Proper pharmaceutical packaging** — Sealed bottles with moisture protection, clear expiration dates, and professional labeling
- ✗ **Moisture damage or clumping** — Capsules should not be sticky, discolored, or show signs of moisture exposure or heat damage
- ✗ **Unusual odor or appearance** — Should be relatively odorless; strong chemical odors may indicate contamination or degradation

What to Expect

- Week 1-2: Gradual increase in energy levels and mental clarity
- Week 2-4: Enhanced exercise performance and recovery
- Week 4-8: Improved metabolic markers and body composition
- Week 8-12: Sustained longevity benefits and cellular health
- Side effects: Generally minimal, occasional mild GI adjustment
- Most effective for: Metabolic health, longevity, energy enhancement

Side Effects & Safety

Side Effects 7 | When to Stop 7

- No human clinical trials published - all dosing extrapolated from animal studies
- 38.4% oral bioavailability demonstrated in rats (PubMed 34304099)
- Start with lowest effective dose (25-50mg) to assess tolerance
- Take with food to reduce potential GI effects
- Monitor for any unusual symptoms and discontinue if adverse effects occur
- Not recommended for pregnant or breastfeeding women
- Consult healthcare provider before use, especially with existing conditions

References

Research Studies 3

Muscle Function in Aged Mice (2024)
Animal study | 10mg/kg/day SubQ | 6 weeks
40% grip strength improvement in sedentary mice, 60% with exercise. No adverse effects observed.
View Study →

Rat Pharmacokinetics Study (2021)
Animal PK study | Various doses | Bioavailability
Demonstrated 38.4% oral bioavailability, half-life 6.9hr (oral) and 3.8hr (IV). Supports oral administration viability.
View Study →

Mouse Obesity Study - Neelakantan et al. (2018)
Animal study | 10mg/kg TID 300mg/kg/day | 11 days | n=10 mice/group
Diet-induced obese mice showed 5.1% weight loss, 30% adipocyte size reduction. Human equivalent ~5mg/kg/day using FDA allometric scaling.
View Study →

Related Peptides

EXTENSIVELY STUDIED
MOTS-c
Metabolism Anti-Aging Exercise
Learn more →

FDA APPROVED
SS-31 (Elamipretide)
Energy Production Heart Health Anti-Aging
Learn more →

EXTENSIVELY STUDIED
NAD+
Energy & Metabolism Anti-Aging Neurological Support
Learn more →

EMERGING RESEARCH
SLU-PP-332
Metabolic Health Heart Health Kidney Health
Learn more →

Curated from research and clinical sources, reviewed by the community. Educational content only. Use responsibly and with caution.

Download on the App Store GET IT ON Google Play

Amazing Supporters

Supporters help fund this project through donations. Listing does not imply endorsement or vetting of any vendors.

- bodyandscience.com
- aussiepeptides.au
- ignitepeptides.com
- STResearch.net
- Roxanne
- JPM
- dr.tanlav
- Hero_labz
- nullifepeptides.com
- JHBiosciences.com
- lg.peptide.warehouse
- Xyr
- nextgenpeppys
- Kevin

Become a supporter →

PepPedia