

Energy-Energy Correlators at Strong Coupling

Lecheng Ren

Queen Mary University of London

September 18, 2026

Based on 2601.05312 with Bo Wang and Congkao Wen
2607.21139 with Max Jackson, Bo Wang, and Congkao Wen

Why energy correlators

- Measured in experiments: e^+e^- colliders, ep colliders, pp colliders.
- Infrared safe; bridge formal theory and experiments.

[Moult, Zhu, 2025 review]

Why $\mathcal{N} = 4$ super Yang–Mills

- AdS/CFT allows analytic computation at strong coupling.

Question

- How does an AdS string amplitude determine the angular energy correlator?

Main Takeaway

- We express EEC as a worldsheet integral.
- This circumvents the complexity of the amplitude itself and helps obtain the EEC in a more efficient way.
- The EEC transformation and the strong coupling expansion *might* not commute.

[Wang, Wen, LR; Jackson, Wang, Wen, LR]

Energy flux operator

$$\mathcal{E}(\hat{n})|X\rangle = \sum_{k \in X} k^0 \delta^2(\Omega_n - \Omega_k) |X\rangle.$$

[Sterman]

An operator definition

$$\mathcal{E}(\hat{n}) = \frac{1}{4} \int_{-\infty}^{\infty} du \lim_{r \rightarrow \infty} r^2 T_{\mu\nu}(u+r, r\hat{n}) \bar{n}^\mu \bar{n}^\nu, \quad \bar{n} = (1, -\hat{n}).$$

where $n = (1, \vec{n})$ is null.

[Sveshnikov, Tkachov; Korchemsky, Sterman; Hofman, Maldacena]

[Kologlu, Kravchuk, Simmons-Duffin, Zhiboedov]

The energy-energy correlator

$$\text{EEC}(\xi) = \frac{8\pi^2}{q^2 \sigma_{\text{tot}}(q)} \int d^4x e^{iq \cdot x} \langle \mathcal{O}(x) \mathcal{E}(n_1) \mathcal{E}(n_2) \mathcal{O}(0) \rangle.$$

$$\sigma_{\text{tot}}(q) = \int d^4x e^{iq \cdot x} \langle \mathcal{O}(x) \mathcal{O}(0) \rangle.$$

- A local source creates the state with total momentum q .
- Two detectors measure the angular correlation of energy flux.
- In the rest frame,

$$\xi = \frac{1 - \cos \theta}{2} = \sin^2 \frac{\theta}{2}, \quad 0 \leq \xi \leq 1.$$

[Hofman, Maldacena; Belitsky, Hohenegger, Korchemsky, Sokatchev, Zhiboedov]

Energy and momentum conservation

$$\int_0^1 d\xi \text{EEC}(\xi) = 1, \quad \int_0^1 d\xi (1 - 2\xi) \text{EEC}(\xi) = 0.$$

- The angular dependence is naturally expanded in Legendre multipoles.

$$\text{EEC}(\xi) = \sum_{s=0}^{\infty} c_s P_s(1 - 2\xi), \quad c_0 = 1, \quad c_1 = 0.$$

- The first unconstrained angular moment is c_2 .

[Dempsey, Karlsson, Pufu, Zahraee, Zhiboedov]

- We choose the source to be a half-BPS scalar operator $\mathcal{O}_p$.
- For $p = 2$, $\mathcal{O}_2$ belongs to the stress-tensor multiplet. The two $\mathcal{O}_2$ insertions are related to energy detectors by supersymmetry and light-ray transforms.
- The local input is the four-point correlator

$$\langle \mathcal{O}_p \mathcal{O}_2 \mathcal{O}_2 \mathcal{O}_p \rangle.$$

[Belitsky, Hohenegger, Korchemsky, Sokatchev, Zhiboedov; Jackson, Wang, Wen, LR]

Examples of EEC

- Weak coupling:

$$\text{EEC}_2(\xi) = \tfrac{1}{2}[\delta(\xi) + \delta(1 - \xi)] + \mathcal{O}(\lambda).$$

- Strong coupling:

$$\text{EEC}(\xi) = 1 + \mathcal{O}(\lambda^{-1}).$$

[Hofman, Maldacena]

- Weak coupling: fruitful knowledge at high multiplicity and high loop order.

[Dixon, Luo, Shtabovenko, Yang, Zhu; Henn, Sokatchev, Yan, Zhiboedov;
He, Jiang, Yang, Zhang; Volovich, Wu, Yan]

- Numerical bootstrap: non-perturbative bounds and finite coupling information from CFT data.

[Dempsey, Karlsson, Pufu, Zahraee, Zhiboedov]

- Strong coupling: the four-point AdS amplitude is known up to first curvature for general p and second curvature correction for $p = 2$.

[Alday, Hansen, Silva; Alday, Hansen; Fardelli, Hansen, Silva]

After analytic continuation and the light-ray transform,

$$\text{EEC}_p(\xi) = \int \frac{ds dt}{(2\pi i)^2} \underbrace{\mathcal{M}_p(s, t)}_{\substack{\text{dynamics} \\ \text{(amplitude)}}} \underbrace{\mathcal{K}_p(t; \xi)}_{\substack{\text{detector/angular} \\ \text{dependence}}} .$$

[Belitsky, Hohenegger, Korchemsky, Sokatchev, Zhiboedov]

- $\mathcal{M}_p(s, t)$ is the Mellin amplitude of the four-point correlator.
- The detector kernel contains the angular dependence. For $p = 2$,

$$\mathcal{K}_2(t; \xi) = \frac{-\pi t^2 (t-2)^2}{128 \xi^3 \sin(\pi t/2)} \left(\frac{\xi}{1-\xi} \right)^{1+t/2} .$$

- The kernel is independent of s . The s -integral is the essential dynamical step.

Note that

$$\text{EEC}_p(\xi) = \int_{i\mathbb{R}} \frac{ds dt}{(2\pi i)^2} \underbrace{\mathcal{M}_p(s, t)}_{\substack{\text{dynamics} \\ \text{(amplitude)}}} \underbrace{\mathcal{K}_p(t; \xi)}_{\substack{\text{detector/angular} \\ \text{dependence}}}.$$

Define

$$Q_{n,p}(\xi) = \int \frac{dt}{2\pi i} t^n \mathcal{K}_p(t; \xi).$$

with Q being a linear combination of Legendre polynomials:

$$Q_{0,2} = P_2(1 - 2\xi), \quad Q_{1,2} = P_2(1 - 2\xi) + 3P_3(1 - 2\xi), \dots$$

Then we have the expansion

$$\text{EEC}_p(\xi) = 1 + \sum_{k=0}^{\infty} \lambda^{-1-k/2} \sum_{n=0}^k c_{k,n}(p) Q_{k-n,p}(\xi).$$

For $p = 2$, the low-energy expansion begins with

$$\mathcal{M}_2(s, t) = \underbrace{\frac{8}{(s-2)(t-2)(\tilde{u}-2)}}_{\text{supergravity}} + \underbrace{\frac{120\zeta(3)}{\lambda^{3/2}} + \dots}_{\text{stringy corrections}}, \quad s + t + \tilde{u} = 4.$$

[Rastelli, Zhou; Alday, Hansen, Silva]

- SUGRA gives $\text{EEC}(\xi) = 1$.
- The stringy terms in the low-energy expansion are polynomials in s, t .
- Their s -integrals are not well-defined term by term on an infinite vertical contour.

We must reorganize the summation and integration.

[Gonçalves]

Relate the AdS amplitude to the Mellin amplitude through a Borel representation:

[Penedones]

$$\mathcal{M}_p(s, t) = \frac{1}{2\lambda^{3/2}\Gamma(p-1)\Gamma(p)} \int_0^\infty d\beta \beta^{p+3} e^{-\beta} A(\beta\hat{s}, \beta\hat{t}),$$
$$\hat{s} = \frac{s - 2p/3}{2\sqrt{\lambda}}, \quad \hat{t} = \frac{t - 2p/3}{2\sqrt{\lambda}}, \quad S = \beta\hat{s}, \quad T = \beta\hat{t}.$$

- In Mellin space, fixed orders in the strong-coupling expansion are polynomial low-energy data.
- In AdS amplitude space, each fixed-curvature object keeps the resummed string-energy dependence:

$$A^{(k)}(S, T) \lesssim \frac{1}{|S|^2}, \quad S \rightarrow i\infty$$

[Maldacena, Shenker, Stanford; Costa, Goncalves, Penedones; Alday, Hansen]

AdS curvature expansion

$$A(S, T) = \sum_{k=0}^{\infty} \lambda^{-k/2} A^{(k)}(S, T), \quad \frac{1}{\sqrt{\lambda}} = \frac{\alpha'}{R^2}, \quad S + T + U = 0.$$

[Alday, Hansen, Silva; Alday, Hansen]

The leading term is the full flat-space Virasoro-Shapiro amplitude:

$$A^{(0)}(S, T) = -\frac{\Gamma(-S)\Gamma(-T)\Gamma(S+T)}{\Gamma(1+S)\Gamma(1+T)\Gamma(1-S-T)}.$$

[Virasoro; Shapiro]

The essential object is

$$I^{(k)}(T) = \int \frac{dS}{2\pi i} A^{(k)}(S, T).$$

Coefficients of $I^{(k)}(T)$ feed the data of EEC expansion in Q .

From $I(T)$ to EEC expansion data

$$I^{(r)}(T) = \sum_{i \geq 0} e_i^{(r)} T^i.$$

$$[Q_{j,p}]_{\lambda^{-k/2-1}} = \frac{1}{\Gamma(p-1)\Gamma(p)} \sum_{i=j}^k \frac{\Gamma(p+3+i)}{2^i} \binom{i}{j} \left(-\frac{2p}{3}\right)^{i-j} e_i^{(k-i)}.$$

	$Q_{0,p}$	$Q_{1,p}$	$Q_{2,p}$
λ^{-1}	$\{e_0^{(0)}\}$	$\{\}$	$\{\}$
$\lambda^{-3/2}$	$\{e_0^{(1)}, e_1^{(0)}\}$	$\{e_1^{(0)}\}$	$\{\}$
λ^{-2}	$\{e_0^{(2)}, e_1^{(1)}, e_2^{(0)}\}$	$\{e_1^{(1)}, e_2^{(0)}\}$	$\{e_2^{(0)}\}$

Closing the S contour

The basic object is a vertical contour integral,

$$I^{(k)}(T) = \int_{\varepsilon - i\infty}^{\varepsilon + i\infty} \frac{dS}{2\pi i} A^{(k)}(S, T).$$

- With the Regge falloff, the contour can be closed in the appropriate half-plane.
- The EEC data can then be read from a sum over pole residues.
- This turns the problem into summing over the massive-pole data of the AdS amplitudes.

$$\text{EEC}^{(k)} \sim \sum_{\text{massive poles}} \text{Res } A^{(k)}.$$

Closing the S contour

The basic object is a vertical contour integral,

$$I^{(k)}(T) = \int_{\varepsilon - i\infty}^{\varepsilon + i\infty} \frac{dS}{2\pi i} A^{(k)}(S, T).$$

- With the Regge falloff, the contour can be closed in the appropriate half-plane.
- The EEC data can then be read from a sum over pole residues.
- This turns the problem into summing over the massive-pole data of the AdS amplitudes.

$$\text{EEC}^{(k)} \stackrel{?}{\sim} \sum_{\text{massive poles}} \text{Res } A^{(k)}.$$

The leading strong-coupling moments are finite:

$$c_3 = \frac{360\zeta(3)}{\lambda^{3/2}} + \cdots, \quad c_4 = \frac{684\pi^4}{7\lambda^2} + \cdots.$$

- The strong-coupling correction is reconstructed from stringy operators.
- The problem appears when the large- λ expansion is taken before resumming the higher-order data.
- The higher order of the residue summations diverge!

[Dempsey, Karlsson, Pufu, Zahraee, Zhiboedov]

Higher orders of the residue sum**Example: the T^1 coefficient of $I^{(1)}(T)$**

$$\text{raw residue sum} = -2\zeta(2)H_\Delta + C_1 + o(1), \quad \Delta \rightarrow \infty,$$

$$C_1 = 2\zeta(2) - 4\zeta(3) + \frac{53}{2}\zeta(4), \quad H_\Delta = \sum_{\delta=1}^{\Delta} \frac{1}{\delta}.$$

The raw residue sum diverges!

- So the formal step “close contour then sum residues” cannot be used naively after the large- λ expansion.

Keep the imaginary-axis integral

$$I^{(k)}(T) = \int_{\varepsilon-i\infty}^{\varepsilon+i\infty} \frac{dS}{2\pi i} A^{(k)}(S, T)$$

- At fixed curvature order, $A^{(k)} \sim 1/|S|^2$ on the imaginary S axis.
- The strong-coupling expansion of the AdS amplitude is well-defined.
- Therefore the S integral itself should give a finite answer for the EEC data.

[large λ expansion, residue sum] $\neq 0$

[large λ expansion, residue sum] $\neq 0$

Instead of the divergence, the non-commutativity also affect the finite terms:

Example: the T^0 coefficient of $I^{(1)}(T)$

The OPE expansion includes:

$$A^{(1)}(S, 0) = (\text{safe terms}) - \sum_{\delta=1}^{\infty} Z_2(\delta-1) \left[\frac{1}{(S-\delta)^2} + \frac{1}{(S+\delta)^2} \right].$$

$$\sum_{\delta} \int \frac{dS}{2\pi i} (\cdots) = 0, \quad \int \frac{dS}{2\pi i} \sum_{\delta} (\cdots) = -\zeta(2).$$

The direct imaginary-axis integral is finite, but reconstructing it from all massive-pole data is cumbersome.

At fixed curvature order, we use two simpler representations:

- ① Worldsheet integrals over a disk and its boundary.
- ② Resummation of the low-energy Wilson coefficients.

[Wang, Wen, LR; Jackson, Wang, Wen, LR]

- Both match with integral along the imaginary axis.
- A guaranteed finite answer.

Worldsheet formulation for AdS Virasoro-Shapiro amplitudes:

$$A^{(k)}(S, T) = \int d^2z |z|^{-2S-2} |1-z|^{-2T-2} G_{\text{tot}}^{(k)}(z, \bar{z}; S, T).$$

[Alday, Hansen, Silva; Alday, Hansen; Fardelli, Hansen, Silva]

- $G_{\text{tot}}^{(k)}$ contains single-valued multiple polylogarithms of weight $3k$.
- First curvature: weight three, including general Kaluza-Klein external states.
- Second curvature: weight six, known for $p = 2$.

Compute $I^{(k)}(T)$ directly by performing the S -integral inside the worldsheet representation.

The S -integral in $I^{(k)}(T)$ acts on terms of the form

$$J_j(T) = \int_{\varepsilon-i\infty}^{\varepsilon+i\infty} \frac{dS}{2\pi i} \int d^2z S^j |z|^{-2S-2} |1-z|^{-2T-2} J(z, \bar{z}; T),$$

with $\varepsilon > 0$, the S -integral localizes the worldsheet integral:

$$j < 0 : \quad \text{contour deformation} \quad \implies \quad |z| < 1,$$

$$j \geq 0 : \quad \delta^{(j)}(\log |z|^2) \quad \implies \quad |z| = 1.$$

- ① Expand the moduli-space integrand at the desired curvature order.
- ② Perform the S -integral first.
- ③ Evaluate disk and boundary-circle integrals and expand over T .

[Wang, Wen, LR; Jackson, Wang, Wen, LR]

Virasoro-Shapiro example revisited

$$\begin{aligned} I^{(0)}(T) &= \int_{|z|<1} d^2z |1-z|^{-2T-2} (-\log |z|^2) \\ &= \frac{1}{T^2} \left[\frac{\Gamma(1-2T)}{\Gamma(1-T)^2} - 1 \right]. \end{aligned}$$

$$I^{(0)}(T) = \zeta(2) + 2\zeta(3)T + \frac{19}{4}\zeta(4)T^2 + \cdots.$$

$$c_{0,0} = 24\zeta(2), \quad c_{1,0} = 120\zeta(3), \quad c_{2,0} = 855\zeta(4).$$

[Dempsey, Karlsson, Pufu, Zahraee, Zhiboedov; Wang, Wen, LR]

Examples from the first curvature correction

These are the two kinds of worldsheet terms needed for $I^{(1)}(T)$.

- Disk integral

$$\begin{aligned} I_1^{(1)}(T) &= \frac{p^2 T}{36} \int_{|z| < 1} d^2 z |z|^{2T-2} |1 - z|^{-2T-2} \log^3 \frac{|z|^2}{|1 - z|^2} \\ &= -\frac{p^2 T}{36} \frac{\partial^3}{\partial T^3} \frac{\Gamma(-2T)}{\Gamma(1 - T)^2}. \end{aligned}$$

- Circle integral, $z = e^{i\theta}$

$$I_3^{(1)}(T) = \oint_{|z|=1} \frac{d\theta}{2\pi} |1 - z|^{-2T-2} (2\zeta(3) - \text{Li}_3(z) - \text{Li}_3(\bar{z})).$$

[Alday, Hansen; Fardelli, Hansen, Silva; Jackson, Wang, Wen, LR]

First curvature correction

- Disk integrals: logarithms and derivatives of Gamma-function integrals.
- Boundary-circle integrals: weight-three polylogarithms.

[Alday, Hansen, Silva; Fardelli, Hansen, Silva]

For $p = 2$, the regular expansion is

$$\begin{aligned} I^{(1)}(T) &= \zeta(2) - \frac{10}{3}\zeta(3) \\ &\quad + \left[-2\zeta(2) + 4\zeta(3) - \frac{53}{2}\zeta(4) \right] T + \mathcal{O}(T^2). \end{aligned}$$

This gives

$$\begin{aligned} c_{1,1} &= 24\zeta(2) - 240\zeta(3), \\ c_{2,1} &= -120\zeta(2) + 240\zeta(3) - 3870\zeta(4). \end{aligned}$$

[Wang, Wen, LR; Jackson, Wang, Wen, LR]

Second curvature correction, $p = 2$

- The worldsheet correlator contains weight-six single-valued multiple polylogarithms.
- To determine the EEC through λ^{-2} , only $I^{(2)}(0)$ is needed.
- This constant receives a contribution only from the unit-circle integral. The regular disk contribution starts at order T .

[Alday, Hansen; Jackson, Wang, Wen, LR]

$$I^{(2)}(0) = 6\zeta(2) - \frac{13}{2}\zeta(3) + \frac{263}{6}\zeta(4) \\ + \zeta(2)\zeta(3) - \frac{5}{3}\zeta(5).$$

EECs from Wilson coefficients

$$\begin{aligned}
\mathcal{M}_2(s, t) &= \mathcal{M}_{\text{SUGRA}}(s, t) \\
&+ \sum_{a, b \geq 0} \frac{\Gamma(2a + 3b + 6)}{8^{a+b} \lambda^{3/2 + a + 3b/2}} (s_1^2 + s_2^2 + s_3^2)^a (s_1 s_2 s_3)^b \\
&\times [\alpha_{a,b}^{(0)} + \lambda^{-1/2} \alpha_{a,b}^{(1)} + \dots], \\
s_1 &= s - \frac{4}{3}, \quad s_2 = t - \frac{4}{3}, \quad s_1 + s_2 + s_3 = 0.
\end{aligned}$$

[Alday, Hansen, Silva; Alday, Hansen]

- Extract the required power of t .
- Resum the Wilson coefficients before performing the unbounded s -integral.

The leading example

$$c_{0,0} = \int \frac{ds_1}{2\pi i} \sum_{a=0}^{\infty} \frac{\Gamma(2a+6)}{4^a} \zeta(2a+3) s_1^{2a}.$$

[Gonçaves]

A useful modified Borel representation is

$$\zeta(n) = \frac{2^{n-1}}{\Gamma(n+1)} \int_0^{\infty} d\omega \frac{\omega^n}{\sinh^2 \omega}, \quad \text{Re } n > 1.$$

[Arutyunov, Dorigoni, Savin; Dorigoni, Green, Wen]

- Sum the infinite low-energy series using this representation.
- The contour integral gives $c_{0,0} = 24\zeta(2)$, in agreement with the worldsheet calculation.

Analytic continuation and localization

$$\begin{aligned}\mathcal{I} &= \int \frac{ds_1}{2\pi i} \sum_{a \geq 0} \frac{f(a)\Gamma(2a+p)}{4^a} \zeta(2a+q, \vec{w}) s_1^{2a+m} \\ &= 2^m f\left(-\frac{m+1}{2}\right) \Gamma(p-m-1) \zeta(q-m-1, \vec{w}).\end{aligned}$$

- The resummed integral localizes the summation variable to $a = -(m+1)/2$.

[Jackson, Wang, Wen, LR]

A term that survives the limit

The first-curvature Wilson coefficient $\alpha_{a,0}^{(1)}$ contains

$$-(2a+1)\zeta(2a+2, 2).$$

At the localized value,

$$\lim_{\varepsilon \rightarrow 0} -\varepsilon \zeta(1+\varepsilon, 2) = -\zeta(2), \quad a = -\frac{1}{2} + \frac{\varepsilon}{2}.$$

In the S -integral on summation over massive poles:

$$\int \frac{dS}{2\pi i} \sum_{\delta=1}^{\infty} Z_2(\delta-1) \left[\frac{1}{(S-\delta)^2} + \frac{1}{(S+\delta)^2} \right] = -\zeta(2)$$

The result also agrees with the worldsheet calculation.

[Alday, Hansen, Silva; Wang, Wen, LR; Jackson, Wang, Wen, LR]

General half-BPS sources

$$\begin{aligned} \text{EEC}_p(\xi) = 1 + \frac{c_{0,0}(p)}{\lambda} Q_{0,p}(\xi) \\ + \frac{1}{\lambda^{3/2}} [c_{1,0}(p) Q_{1,p}(\xi) + c_{1,1}(p) Q_{0,p}(\xi)] + \cdots, \end{aligned}$$

$$c_{0,0}(p) = \frac{\Gamma(p+3)}{\Gamma(p-1)\Gamma(p)} \zeta(2), \quad c_{1,0}(p) = \frac{\Gamma(p+4)}{\Gamma(p-1)\Gamma(p)} \zeta(3),$$

$$c_{1,1}(p) = \frac{\Gamma(p+3)}{\Gamma(p-1)\Gamma(p)} [\zeta(2) - (p^2 + 2p + 2)\zeta(3)].$$

[Jackson, Wang, Wen, LR]

For $p = 2$, through order λ^{-2}

$$\begin{aligned} \text{EEC}(\xi) = & 1 + \frac{24\zeta(2)}{\lambda} Q_{0,2}(\xi) \\ & + \frac{1}{\lambda^{3/2}} [120\zeta(3)Q_{1,2}(\xi) + (24\zeta(2) - 240\zeta(3))Q_{0,2}(\xi)] \\ & + \frac{1}{\lambda^2} [c_{2,0}Q_{2,2}(\xi) + c_{2,1}Q_{1,2}(\xi) + c_{2,2}Q_{0,2}(\xi)] + \cdots, \end{aligned}$$


$$c_{2,0} = 855\zeta(4),$$

$$c_{2,1} = -120\zeta(2) + 240\zeta(3) - 3870\zeta(4),$$


$$c_{2,2} = 304\zeta(2) - 476\zeta(3) + 24\zeta(2)\zeta(3) + 4692\zeta(4) - 40\zeta(5).$$

Worldsheet integrals and the regulated low-energy calculation agree, as well as the S -integral along the imaginary axis numerically.


[Wang, Wen, LR; Jackson, Wang, Wen, LR]



[Dempsey, Karlsson, Pufu, Zahraee, Zhiboedov; Jackson, Wang, Wen, LR]



[Dempsey, Karlsson, Pufu, Zahraee, Zhiboedov; Jackson, Wang, Wen, LR]



[Dempsey, Karlsson, Pufu, Zahraee, Zhiboedov; Jackson, Wang, Wen, LR]

Summary

- EECs in $\mathcal{N} = 4$ SYM are closely related to AdS Virasoro-Shapiro amplitudes.
- Worldsheet disk and boundary integrals provide a direct calculation.
- Resummation of Wilson coefficients gives an independent analytic cross-check.
- Both approach guarantee a finite result, therefore should work at any higher orders.

Open questions

- Can we bootstrapp moduli space integrand for EEC directly without AdS amplitudes?
- Numerical bootstrapp to more precise value.
- EECs with heavy states and large charge.
- Less supersymmetric theories?

Thank you!

The remaining EEC integrals

$$\text{EEC}_p(\xi) = \frac{1}{\lambda \Gamma(p-1) \Gamma(p)} \int \frac{dt}{2\pi i} \int_0^\infty d\beta e^{-\beta} \beta^{p+2} I(\beta \hat{t}) \mathcal{K}_p(t; \xi).$$

$$\hat{t} = \frac{t - 2p/3}{2\sqrt{\lambda}}, \quad \int_0^\infty d\beta e^{-\beta} \beta^{p+2+i} = \Gamma(p+3+i).$$

- Expand $I(T)$ to the required order.
- Include the shift in $\hat{t}$ before using $Q_{n,p}$.
- Treat the massless or SUGRA singular contribution separately from the regular string expansion.

[Wang, Wen, LR; Jackson, Wang, Wen, LR]

First curvature: cutoff dependence

Raw simple-pole residue sums, as $\Delta \rightarrow \infty$:

$$T^1 : -2\zeta(2)H_\Delta + C_1 + o(1), \quad T^2 : -2\zeta(2)H_\Delta^2 + C_2 + o(1).$$

$$C_1 = 2\zeta(2) - 4\zeta(3) + \frac{53}{2}\zeta(4),$$

$$C_2 = -4\zeta(2) - 6\zeta(4) + 32\zeta(2)\zeta(3) + 8\zeta(3) + 74\zeta(5).$$

After finite-axis integration and mass summation, the T^1 contributions are

$$\text{simple:} \quad 2\zeta(2)(\log Y + \gamma_E) - C_1 + o(1),$$

$$\text{double:} \quad -2\zeta(2)(\log Y + \gamma_E) + o(1).$$

- Their sum tends to $-C_1$; the raw residue sum still diverges.
- At T^2 , squared logarithms cancel between simple and double poles.
- These are ordered asymptotics, not exact finite- (Y, Δ) identities.

Second curvature: cancellation between pole orders

At $T = 0$, keep common cutoffs. The divergent coefficients are

Pole order	simple	double	triple	fourth
$\zeta(4)YH_{\Delta}/\pi$	-7	2	5	0
$\zeta(2)(\log Y + \gamma_E)$	$-29/6$	$43/6$	$-4/3$	-1

$$-7 + 2 + 5 = 0, \quad -\frac{29}{6} + \frac{43}{6} - \frac{4}{3} - 1 = 0.$$

- The first row concerns mass-cutoff asymptotics at fixed Y ; the second concerns the subsequent large- Y expansion.
- Fifth, sixth and seventh pole orders give $o(1)$ in the ordered limit, not zero at finite cutoffs.
- The combined principal parts still leave $-19\zeta(4)Y/(2\pi)$.

Second curvature: the Regge condition

The massive-pole principal parts approach $-19\zeta(4)/2$ on the large imaginary axis. Regge decay fixes the compensating amplitude constant:

$$\int_{-iY}^{iY} \frac{dS}{2\pi i} \frac{19}{2} \zeta(4) = \frac{19\zeta(4)Y}{2\pi}.$$

This is an amplitude constant, not a missing simple-pole residue.

$$\begin{aligned} I^{(2)}(0) &= \frac{263}{6} \zeta(4) + 6\zeta(2) + \left(\zeta(2) - \frac{13}{2} \right) \zeta(3) - \frac{5}{3} \zeta(5) \\ &= 49.7471610332818994 \dots \end{aligned}$$

- This is the ordered limit with the Regge completion included.
- Even at $Y = 10^6$, the completed contour gives $49.7282427441 \dots$
- Numerical checks of subtracted residue remainders do not establish the convergence rate of the finite-axis integral.