

Due before lecture on Monday, November 25, 2019

- Find all eigenvectors and eigenvalues of the matrix $B = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 4 & 0 \\ 12 & 2 & 2 \end{bmatrix}$. Write 1-3 sentences that interpret these geometrically—in other words, what do the eigenvectors and eigenvalues tell you about the transformation geometrically (in terms of stretch factors and stretch directions)?
- Construct an example for each of the following, or explain why such an example does not exist:
 - a 2×2 matrix that is invertible but not diagonalizable
 - a 2×2 non-diagonal matrix that is diagonalizable but not invertible
- (Strang §5.2 #11) If all eigenvalues of A are 1, 1, and 2, which of the following are certain to be true? Give a reason if true or a counterexample if false.
 - A is invertible.
 - A is diagonalizable.
 - A is not diagonalizable.
- (Strang §5.2 #12) Suppose the only eigenvectors of A are multiples of $\vec{x} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, which of the following are certain to be true? Give a reason if true or a counterexample if false.
 - A is not invertible.
 - A has a repeated eigenvalue.
 - A is not diagonalizable.
- (Strang §5.1 #18) Suppose a 3×3 matrix A has eigenvalues 0, 3, and 5 with associated eigenvectors \vec{u} , \vec{v} , and \vec{w} respectively.
 - Since the eigenvalues of A are all distinct, the set $\{\vec{u}, \vec{v}, \vec{w}\}$ is _____.
 - Write down a basis for the nullspace $N(A)$ and the column space $C(A)$.
 - Find one particular solution to $A\vec{x} = \vec{v} + \vec{w}$. Find all solutions to $A\vec{x} = \vec{v} + \vec{w}$.
 - Explain why $A\vec{x} = \vec{u}$ does not have a solution. (Hint: If there is a solution, then _____ is in $C(A)$. Explain why that is impossible.)
 - Is A invertible? Why or why not?
- Let A be an n -by- n matrix. Suppose $A\vec{u} = 2\vec{u}$ and $A\vec{v} = 5\vec{v}$ for nonzero vectors \vec{u} and \vec{v} . Complete the following proof that $\{\vec{u}, \vec{v}\}$ is linearly independent.
 Proof: In order for $\{\vec{u}, \vec{v}\}$ to be linearly independent, we need to show that

the only solution to the vector equation $x\vec{u} + y\vec{v} = \vec{0}$ is the trivial one.

Note that in this equation, \vec{u} and \vec{v} are known vectors, while x and y are unknown scalars. Now assume that $x = a$ and $y = b$ is some solution to the above equation. That is

$$a\vec{u} + b\vec{v} = \vec{0}. \tag{1}$$

Multiply both sides of equation (1) by matrix A from the left. Show your calculation details to explain why the following has to be true as well

$$2a\vec{u} + 5b\vec{v} = \vec{0}. \quad (2)$$

Explain in detail how combining vectors equations (1) and (2) leads to the conclusion that $a = 0 = b$. Therefore, the only solution to $x\vec{u} + y\vec{v} = \vec{0}$ is the trivial solution.

7. In one of the reflection questions, you saw that if A is 2×2 matrix then the product of its eigenvalues is equal to $\det(A)$ and the sum is equal to $\text{trace}(A)$. The following shows that both claims still hold true for $n \times n$ matrices.

Suppose that $\lambda_1, \lambda_2, \dots, \lambda_n$ are the n eigenvalues of an $n \times n$ matrix A . λ_i 's are the roots of the polynomial $\det(A - \lambda I)$, which means that we have a factorization

$$\det(A - \lambda I) = (\lambda_1 - \lambda)(\lambda_2 - \lambda) \cdots (\lambda_n - \lambda) \quad (3)$$

- (a) (*Strang §5.1 #8*) By making a clever choice of the value for λ in equation (3), show that $\det(A)$ is equal to the product of eigenvalues.
- (b) (*Strang §5.1 #9*) Show that $\text{trace}(A)$ is equal to the sum of eigenvalues in three steps. First, find the coefficient of $(-\lambda)^{n-1}$ on the righthand side of equation (3). Next, find all terms on the righthand side of the following that involves $(-\lambda)^{n-1}$

$$\det(A - \lambda I) = \det \begin{bmatrix} a_{11} - \lambda & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} - \lambda & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} - \lambda \end{bmatrix},$$

where a_{ij} are the entries of A . Add up all those terms to find the coefficient of $(-\lambda)^{n-1}$. Lastly, compare the coefficient of $(-\lambda)^{n-1}$ found in these two different ways.