Elevated N-glycosylation of immunoglobulin G variable regions in myasthenia gravis highlights a commonality across autoantibody-associated diseases

What is the Manuscript Microscope Sentence Audit?

The Manuscript Microscope Sentence Audit is a research paper introspection system that parses the text of your manuscript into minimal sentence components for faster, more accurate, enhanced proofreading.

Why use a Sentence Audit to proofread your manuscript?

- Accelerated Proofreading: Examine long technical texts in a fraction of the usual time.
- Superior Proofreading: Detect subtle errors that are invisible to traditional methods.
- Focused Proofreading: Inspect each individual sentence component in isolation.
- Reliable Proofreading: Ensure every single word of your manuscript is correct.
- Easier Proofreading: Take the hardship out of crafting academic papers.

Bonus 1: Improved Productivity: Rapidly refine rough drafts to polished papers.
Bonus 2: Improved Authorship: Cultivate a clear, concise, consistent, writing style.
Bonus 3: Improved Reputation: Become known for rigorously precise publications.

Manuscript Source: https://www.biorxiv.org/content/10.1101/2021.03.06.434199v1

Manuscript Authors: Caleigh Mandel-Brehm, Miriam L. Fichtner, Ruoyi Jiang, Valerie J. Winton,

Sara E. Vazquez, Minh C. Pham, Kenneth B. Hoehn, Neil L. Kelleher,

Richard J. Nowak, Steven H. Kleinstein, Michael R. Wilson, Joseph L. DeRisi

& Kevin C. O'Connor

Audit Date: 29/03/21 Audit Identifier: B851094K5GY4PH3 Code Version: 3.6

Features of the Sentence Audit:

The Sentence Audit combines two complementary proofreading approaches:

- 1. Each sentence of your text is parsed and displayed in isolation for focused inspection.
- 2. Each individual sentence is further parsed into Minimal Sentence Components for a deeper review of the clarity, composition and consistency of the language you used.

The Minimal Sentence Components shown are the smallest coherent elements of each sentence of your text as derived from it's conjunctions, prepositions and selected punctuation symbols (i.e. commas, semicolons, round and square brackets).

The combined approaches ensure easier, faster, more effective proofreading.

Comments and Caveats:

- The sentence parsing is achieved using a prototype natural language processing pipeline written in Python and may include occasional errors in sentence segmentation.
- Depending on the source of the input text, the Sentence Audit may contain occasional html artefacts that are parsed as sentences (E.g. "Download figure. Open in new tab").
- Always consult the original research paper as the true reference source for the text.

Contact Information:

To get a Manuscript Microscope Sentence Audit of any other research paper, simply forward any copy of the text to John.James@OxfordResearchServices.com.

All queries, feedback or suggestions are also very welcome.

Research Paper Sections:

The sections of the research paper input text parsed in this audit.

Section No.	Headings	Sentences
Section: 1	Abstract	10
Section: 2	Introduction	17
N/A		0

Title regions in myasthenia gravis highlights a commonality across autoantibody-associated diseases

S1 [001] Abstract

S1 [002] Elevated N-linked glycosylation of immunoglobulin G variable regions (lgG-VN-Glyc) is an emerging molecular phenotype associated with autoimmune disorders.

Elevated N-linked glycosylation ...
... of immunoglobulin G variable regions ...
... (IgG-VN-Glyc) ...
... is an emerging molecular phenotype associated ...
... with autoimmune disorders.

S1 [003] To test the broader specificity of elevated IgG-VN-Glyc, we studied patients with distinct subtypes of myasthenia gravis (MG), a B cell-mediated autoimmune disease.

To test the broader specificity ...
... of elevated IgG-VN-Glyc, ...
... we studied patients ...
... with distinct subtypes ...
... of myasthenia gravis ...
... (MG), ...
... a B cell-mediated autoimmune disease.

... and antigen binding assays ...
... to confirm occupation ...
... of glycosylation sites ...
... and determine ...

... whether they alter binding.

S1 [004] Our experimental design included adaptive immune receptor repertoire sequencing to quantify and characterize N-glycosylation sites in the global B cell receptor repertoire, proteomics to examine glycosylation patterns of the circulating IgG, and production of human-derived recombinant autoantibodies, which were studied with mass spectrometry and antigen binding assays to confirm occupation of glycosylation sites and determine whether they alter binding.

Our experimental design included adaptive immune receptor repertoire sequencing ...
... to quantify ...
... and characterize N-glycosylation sites ...
... in the global B cell receptor repertoire, ...
... proteomics ...
... to examine glycosylation patterns ...
... of the circulating IgG, ...
... and production ...
... of human-derived recombinant autoantibodies, ...
... which were studied ...
... with mass spectrometry ...

S1 [005] We found that the frequency of IgG-VN-Glyc motifs was increased in the B cell repertoire of MG patients when compared to healthy donors. We found that the frequency of IgG-VN-Glyc motifs was increased in the B cell repertoire of MG patients when compared to healthy donors. S1 [006] Motifs were introduced by both biased V gene segment usage and somatic hypermutation. Motifs were introduced by both biased V gene segment usage and somatic hypermutation. S1 [007] IgG-VN-Glyc could be observed in the circulating IgG in a subset of MG patients. IgG-VN-Glyc could be observed in the circulating IgG in a subset of MG patients. S1 [008] Autoantigen binding, by patient-derived MG autoantigen-specific monoclonal antibodies with experimentally confirmed presence of IgG-VN-Glyc, was not altered by the glycosylation. Autoantigen binding, by patient-derived MG autoantigen-specific monoclonal antibodies with experimentally confirmed presence of IgG-VN-Glyc, was not altered by the glycosylation. S1 [009] Our findings extend prior work on patterns of variable region N-linked glycosylation in autoimmunity to MG subtypes. Our findings extend prior work on patterns of variable region N-linked glycosylation in autoimmunity to MG subtypes. S1 [010] Although occupied IgG-VN-Glyc motifs are found on MG autoantigen-specific monoclonal antibodies, they are not required for binding to the autoantigen in this disease.

Although occupied IgG-VN-Glyc motifs are found on MG autoantigen-specific monoclonal antibodies, ...

... they are not required ...

... for binding ...
... to the autoantigen ...

S2 [011] Introduction

S2 [012] The vast diversity of immunoglobulin G variable regions (IgG-V) is critical for host immunity.

```
The vast diversity ...
... of immunoglobulin G variable regions ...
... (IgG-V) ...
... is critical ...
... for host immunity.
```

S2 [013] This diversity arises through VDJ recombination and somatic hypermutation (SHM).

```
This diversity arises ...
... through VDJ recombination ...
... and somatic hypermutation ...
... (SHM).
```

S2 [014] Historically, IgG-V diversity has been represented by amino acid sequence alone with little focus on post-translational modifications.

```
Historically, ...
... IgG-V diversity has been represented ...
... by amino acid sequence alone ...
... with little focus ...
... on post-translational modifications.
```

S2 [015] Recently, the presence of N-linked glycosylation in IgG-V (IgG-VN-Glyc) has been shown to contribute to diversity (1, 2).

```
Recently, ...
... the presence ...
... of N-linked glycosylation ...
... in IgG-V ...
... (IgG-VN-Glyc) ...
... has been shown ...
... to contribute ...
... to diversity ...
... (1, 2)...
```

S2 [016] IgG-VN-Glyc is contingent upon the presence of the predictive N-Glyc amino acid motif N-X-S/T, where X can be any amino acid except for proline.

```
IgG-VN-Glyc is contingent ...
... upon the presence ...
... of the predictive N-Glyc amino acid motif N-X-S/T, ...
... where X can be any amino acid except ...
... for proline.
```

S2 [017] This motif is most often introduced as a consequence of SHM (3).

```
This motif is most often introduced ...
... as a consequence ...
... of SHM ...
... (3).
```

S2 [018] Less often it can be provided by the few germline gene segments (IGHV1-8, IGHV4-34, IGHV5-10-1, IGLV3-12, and IGLV5-37) in which it is encoded (4).

```
Less often it can be provided ...
... by the few germline gene segments ...
... (IGHV1-8, ...
... IGHV4-34, ...
... IGHV5-10-1, ...
... IGLV3-12, ...
... and IGLV5-37) ...
... in which it is encoded ...
... (4).
```

S2 [019] The percentage of IgG in healthy individuals that includes V-region glycosylation is approximately 15-25%; the range reflects different approaches of measurement (1).

```
The percentage ...
... of IgG ...
... in healthy individuals ...
... that includes V-region glycosylation is approximately 15-25%; ...
... the range reflects different approaches ...
... of measurement ...
... (1).
```

S2 [020] The occurrence of IgG-VN-Glyc also varies among the IgG subclasses, with skewing toward higher frequencies in antibodies of the IgG4 subclass (5).

```
The occurrence ...
... of IgG-VN-Glyc also varies ...
... among the IgG subclasses, ...
... with skewing toward higher frequencies ...
... in antibodies ...
... of the IgG4 subclass ...
... (5).
```

S2 [021] Higher frequencies of IgG-VN-Glyc than that which is found in healthy individuals have been observed in B cell malignancies (6–10) and in autoimmune diseases (11).

```
Higher frequencies ...
... of IgG-VN-Glyc ...
... than that ...
... which is found ...
... in healthy individuals have been observed ...
... in B cell malignancies ...
... (6–10) ...
... and in autoimmune diseases ...
```

End of Sample Audit

This is a truncated Manuscript Microscope Sample Audit.

To get the full audit of this text (or any other research paper), forward a copy of the research paper to John James at John.James@OxfordResearchServices.com