



Report: Transient Grating Project

Manuel Fernando Sánchez Alarcón

May 9, 2026

Contents

1	Introduction	2
2	Fitting Models	3
2.1	Model 1	3
2.2	Model 2	3
2.3	Model 3	4
3	Results	5
3.1	Phase Space	5
3.2	Model 1	6
3.2.1	Constant Intensity	6
3.2.2	Constant Energy	13
3.3	Model 2	19
3.3.1	Constant Intensity	19
3.3.2	Constant Energy	29
3.4	Model 3	37
3.4.1	Constant Intensity	37
3.4.2	Constant Energy	45
3.5	Comparison of Models	52
3.5.1	Constant Intensity	52
3.5.2	Constant Energy	59

Chapter 1

Introduction

This report summarizes the current status of transient grating signal analysis. It is intended as the companion document for the code and notebook workflow in this project.

Chapter 2

Fitting Models

For this report, three different models have been implemented and fitted to the data. The models are described in the following sections.

2.1 Model 1

The first model is described by the following equation:

$$\begin{aligned} f(t) &= [A_1 e^{-t/\tau} + A_{offset}] \otimes \frac{1}{\sqrt{2\pi}\sigma} e^{-(t-t_0)^2/(2\sigma^2)} \\ &= A_1 e^{-(t-t_0)/\tau} e^{\sigma^2/(2\tau^2)} \left(1 + \operatorname{erf} \left(\frac{t-t_0-\sigma^2/\tau}{\sqrt{2}\sigma} \right) \right) \\ &\quad + A_{offset} \left(1 + \operatorname{erf} \left(\frac{t-t_0}{\sqrt{2}\sigma} \right) \right) \\ &= f_{1,1}(t) + f_{1,2}(t). \end{aligned} \tag{2.1}$$

2.2 Model 2

The second model is described by the following equation:

$$\begin{aligned} f(t) &= [A_1 e^{-t/\tau} + A_2 e^{-(t-t_{0,2})/\tau_2} + A_{offset}] \otimes \frac{1}{\sqrt{2\pi}\sigma} e^{-(t-t_0)^2/(2\sigma^2)} \\ &= A_1 e^{-(t-t_0)/\tau} e^{\sigma^2/(2\tau^2)} \left(1 + \operatorname{erf} \left(\frac{t-t_0-\sigma^2/\tau}{\sqrt{2}\sigma} \right) \right) \\ &\quad + A_2 e^{-(t-t'_{0,2})/\tau_2} e^{\sigma^2/(2\tau_2^2)} \left(1 + \operatorname{erf} \left(\frac{t-t'_{0,2}-\sigma^2/\tau_2}{\sqrt{2}\sigma} \right) \right) \\ &\quad + A_{offset} \left(1 + \operatorname{erf} \left(\frac{t-t_0}{\sqrt{2}\sigma} \right) \right) \\ &= f_{2,1}(t) + f_{2,2}(t) + f_{2,3}(t). \end{aligned} \tag{2.2}$$

2.3 Model 3

The third model is described by the following equation:

$$\begin{aligned}
f(t) &= [A_1 e^{-t/\tau} + A_2 e^{-(t-t_{0,2})/\tau_2} + A_3 e^{-(t-t_{0,3})/\tau_3}] \circledast \frac{1}{\sqrt{2\pi}\sigma} e^{-(t-t_0)^2/(2\sigma^2)} \\
&= A_1 e^{-(t-t_0)/\tau} e^{\sigma^2/(2\tau^2)} \left(1 + \operatorname{erf} \left(\frac{t-t_0-\sigma^2/\tau}{\sqrt{2}\sigma} \right) \right) \\
&\quad + A_2 e^{-(t-t'_{0,2})/\tau_2} e^{\sigma^2/(2\tau_2^2)} \left(1 + \operatorname{erf} \left(\frac{t-t'_{0,2}-\sigma^2/\tau_2}{\sqrt{2}\sigma} \right) \right) \\
&\quad + A_3 e^{-(t-t'_{0,3})/\tau_3} e^{\sigma^2/(2\tau_3^2)} \left(1 + \operatorname{erf} \left(\frac{t-t'_{0,3}-\sigma^2/\tau_3}{\sqrt{2}\sigma} \right) \right) \\
&= f_{3,1}(t) + f_{3,2}(t) + f_{3,3}(t).
\end{aligned} \tag{2.3}$$

Chapter 3

Results

3.1 Phase Space

The available scans for the dataset CoOx12 are represented by each point in the phase space plot shown in Figure 3.1. The phase space plot is a scatter plot of the energy and intensity of the scans.




Figure 3.1: Phase space plot of the available scans for the dataset CoOx12.

3.2 Model 1

3.2.1 Constant Intensity

In this case, only scans with a constant intensity of 2 μJ are considered. Each individual signal was fitted with the Model 1 and the results are shown in Figure 3.2, together with the corresponding residuals (in green). All signals (experimental and fitted) are shown one on top of the other in Figure 3.4. The single fit components are plotted in Figure 3.3.




Figure 3.2: Fitted signals for the Model 1 with constant intensity of 2 μJ . The fitted curve is shown in red, the experimental data is shown in blue, and the residuals are shown in green. Values for the χ^2/dof goodness of fit are shown in the title of each plot, as well as the R^2 value.




Figure 3.3: Single fit components for the Model 1 with constant intensity of 2 μJ . The components are shown in green ($f_{1,1}(t)$) and purple ($f_{1,2}(t)$). The fit is shown in red.




Figure 3.4: Stacked signals for the Model 1 with constant intensity of $2 \mu\text{J}$. The experimental data is shown in blue, and the fitted curve is shown in red. All signals are shown one on top of the other.

The values of A_1 for the Model 1 are shown in Figure 3.5.




Figure 3.5: Values of A_1 for the Model 1 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of τ for the Model 1 are shown in Figure 3.6.




Figure 3.6: Values of τ for the Model 1 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of σ for the Model 1 are shown in Figure 3.7.




Figure 3.7: Values of σ for the Model 1 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of A_{offset} for the Model 1 are shown in Figure 3.8.




Figure 3.8: Values of A_{offset} for the Model 1 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of t_0 for the Model 1 are shown in Figure 3.9.




Figure 3.9: Values of t_0 for the Model 1 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of R^2 for the Model 1 are shown in Figure 3.10.




Figure 3.10: Values of R^2 for the Model 1 with constant intensity of 2 μJ . The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

3.2.2 Constant Energy

In this case, only scans with a constant energy of 63.6 eV are considered. Each individual signal was fitted with the Model 1 and the results are shown in Figure 3.11, together with the corresponding residuals (in green). All signals (experimental and fitted) are shown one on top of the other in Figure 3.13. The single fit components are plotted in Figure 3.12.




Figure 3.11: Fitted signals for the Model 1 with constant energy of 63.6 eV. The fitted curve is shown in red, the experimental data is shown in blue, and the residuals are shown in green. Values for the χ^2/dof goodness of fit are shown in the title of each plot, as well as the R^2 value.




Figure 3.12: Single fit components for the Model 1 with constant energy of 63.6 eV. The components are shown in green ($f_{1,1}(t)$) and purple ($f_{1,2}(t)$). The fit is shown in red.




Figure 3.13: Stacked signals for the Model 1 with constant energy of 63.6 eV. The experimental data is shown in blue, and the fitted curve is shown in red. All signals are shown one on top of the other.

The values of A_1 for the Model 1 are shown in Figure 3.14.




Figure 3.14: Values of A_1 for the Model 1 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of τ for the Model 1 are shown in Figure 3.15.




Figure 3.15: Values of τ for the Model 1 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of σ for the Model 1 are shown in Figure 3.16.




Figure 3.16: Values of σ for the Model 1 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of A_{offset} for the Model 1 are shown in Figure 3.17.




Figure 3.17: Values of A_{offset} for the Model 1 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of t_0 for the Model 1 are shown in Figure 3.18.




Figure 3.18: Values of t_0 for the Model 1 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of R^2 for the Model 1 are shown in Figure 3.19.




Figure 3.19: Values of R^2 for the Model 1 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

3.3 Model 2

3.3.1 Constant Intensity

In this case, only scans with a constant intensity of 2 μJ are considered. Each individual signal was fitted with the Model 2 and the results are shown in Figure 3.20, together with the corresponding residuals (in green). All signals (experimental and fitted) are shown one on top of the other in Figure 3.22. The single fit components are plotted in Figure 3.21.




Figure 3.20: Fitted signals for the Model 2 with constant intensity of $2 \mu\text{J}$. The fitted curve is shown in red, the experimental data is shown in blue, and the residuals are shown in green. Values for the χ^2/dof goodness of fit are shown in the title of each plot, as well as the R^2 value.




Figure 3.21: Single fit components for the Model 2 with constant intensity of 2 μ J. The components are shown in green ($f_{2,1}(t)$), orange ($f_{2,2}(t)$), and purple ($f_{2,3}(t)$). The fit is shown in red.




Figure 3.22: Stacked signals for the Model 2 with constant intensity of $2 \mu\text{J}$. The experimental data is shown in blue, and the fitted curve is shown in red. All signals are shown one on top of the other.

The values of A_1 for the Model 2 are shown in Figure 3.23.




Figure 3.23: Values of A_1 for the Model 2 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of τ for the Model 2 are shown in Figure 3.24.




Figure 3.24: Values of τ for the Model 2 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of σ for the Model 2 are shown in Figure 3.25.




Figure 3.25: Values of σ for the Model 2 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of A_{offset} for the Model 2 are shown in Figure 3.26.




Figure 3.26: Values of A_{offset} for the Model 2 with constant intensity of 2 μJ . The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of t_0 for the Model 2 are shown in Figure 3.27.




Figure 3.27: Values of t_0 for the Model 2 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of A_2 for the Model 2 are shown in Figure 3.28.




Figure 3.28: Values of A_2 for the Model 2 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of τ_2 for the Model 2 are shown in Figure 3.29.




Figure 3.29: Values of τ_2 for the Model 2 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of R^2 for the Model 2 are shown in Figure 3.30.




Figure 3.30: Values of R^2 for the Model 2 with constant intensity of 2 μJ . The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

3.3.2 Constant Energy

In this case, only scans with a constant energy of 63.6 eV are considered. Each individual signal was fitted with the Model 2 and the results are shown in Figure 3.31, together with the corresponding residuals (in green). All signals (experimental and fitted) are shown one on top of the other in Figure 3.33. The single fit components are plotted in Figure 3.32.




Figure 3.31: Fitted signals for the Model 2 with constant energy of 63.6 eV. The fitted curve is shown in red, the experimental data is shown in blue, and the residuals are shown in green. Values for the χ^2/dof goodness of fit are shown in the title of each plot, as well as the R^2 value.




Figure 3.32: Single fit components for the Model 2 with constant energy of 63.6 eV. The components are shown in green ($f_{2,1}(t)$), orange ($f_{2,2}(t)$), and purple ($f_{2,3}(t)$). The fit is shown in red.




Figure 3.33: Stacked signals for the Model 2 with constant energy of 63.6 eV. The experimental data is shown in blue, and the fitted curve is shown in red. All signals are shown one on top of the other.

The values of A_1 for the Model 2 are shown in Figure 3.34.




Figure 3.34: Values of A_1 for the Model 2 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of τ for the Model 2 are shown in Figure 3.35.




Figure 3.35: Values of τ for the Model 2 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of σ for the Model 2 are shown in Figure 3.36.




Figure 3.36: Values of σ for the Model 2 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of A_{offset} for the Model 2 are shown in Figure 3.37.




Figure 3.37: Values of A_{offset} for the Model 2 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of t_0 for the Model 2 are shown in Figure 3.38.




Figure 3.38: Values of t_0 for the Model 2 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of A_2 for the Model 2 are shown in Figure 3.39.




Figure 3.39: Values of A_2 for the Model 2 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of τ_2 for the Model 2 are shown in Figure 3.40.




Figure 3.40: Values of τ_2 for the Model 2 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of R^2 for the Model 2 are shown in Figure 3.41.




Figure 3.41: Values of R^2 for the Model 2 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

3.4 Model 3

3.4.1 Constant Intensity

In this case, only scans with a constant intensity of 2 μJ are considered. Each individual signal was fitted with the Model 3 and the results are shown in Figure 3.42, together with the corresponding residuals (in green). All signals (experimental and fitted) are shown one on top of the other in Figure 3.44. The single fit components are plotted in Figure 3.43.




Figure 3.42: Fitted signals for the Model 3 with constant intensity of 2 μJ . The fitted curve is shown in red, the experimental data is shown in blue, and the residuals are shown in green. Values for the χ^2/dof goodness of fit are shown in the title of each plot, as well as the R^2 value.




Figure 3.43: Single fit components for the Model 3 with constant intensity of 2 μJ . The components are shown in green ($f_{3,1}(t)$), orange ($f_{3,2}(t)$), and purple ($f_{3,3}(t)$). The fit is shown in red.




Figure 3.44: Stacked signals for the Model 3 with constant intensity of 2 μJ . The experimental data is shown in blue, and the fitted curve is shown in red. All signals are shown one on top of the other.

The values of A_1 for the Model 3 are shown in Figure 3.45.




Figure 3.45: Values of A_1 for the Model 3 with constant intensity of 2 μJ . The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of A_2 for the Model 3 are shown in Figure 3.46.




Figure 3.46: Values of A_2 for the Model 3 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of t_0 for the Model 3 are shown in Figure 3.47.




Figure 3.47: Values of t_0 for the Model 3 with constant intensity of 2 μJ . The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of τ for the Model 3 are shown in Figure 3.48.




Figure 3.48: Values of τ for the Model 3 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{2+} , Co^{3+} and Co_3O_4 are shown as area plots.

The values of τ_2 for the Model 3 are shown in Figure 3.49.




Figure 3.49: Values of τ_2 for the Model 3 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of σ for the Model 3 are shown in Figure 3.50.




Figure 3.50: Values of σ for the Model 3 with constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

The values of R^2 for the Model 3 are shown in Figure 3.51.




Figure 3.51: Values of R^2 for the Model 3 with constant intensity of 2 μJ . The values are shown as a function of the energy of the scan. As a visual reference, values of absorbance for Co^{+2} , Co^{+3} and Co_3O_4 are shown as area plots.

3.4.2 Constant Energy

In this case, only scans with a constant energy of 63.6 eV are considered. Each individual signal was fitted with the Model 3 and the results are shown in Figure 3.52, together with the corresponding residuals (in green). All signals (experimental and fitted) are shown one on top of the other in Figure 3.54. The single fit components are plotted in Figure 3.53.




Figure 3.52: Fitted signals for the Model 3 with constant energy of 63.6 eV. The fitted curve is shown in red, the experimental data is shown in blue, and the residuals are shown in green. Values for the χ^2/dof goodness of fit are shown in the title of each plot, as well as the R^2 value.




Figure 3.53: Single fit components for the Model 3 with constant energy of 63.6 eV. The components are shown in green, orange, purple and black. The fit is shown in red.




Figure 3.54: Stacked signals for the Model 3 with constant energy of 63.6 eV. The experimental data is shown in blue, and the fitted curve is shown in red. All signals are shown one on top of the other.

The values of A_1 for the Model 3 are shown in Figure 3.55.




Figure 3.55: Values of A_1 for the Model 3 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of A_2 for the Model 3 are shown in Figure 3.56.




Figure 3.56: Values of A_2 for the Model 3 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of t_0 for the Model 3 are shown in Figure 3.57.




Figure 3.57: Values of t_0 for the Model 3 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of τ for the Model 3 are shown in Figure 3.58.




Figure 3.58: Values of τ for the Model 3 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of τ_2 for the Model 3 are shown in Figure 3.59.




Figure 3.59: Values of τ_2 for the Model 3 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of σ for the Model 3 are shown in Figure 3.60.




Figure 3.60: Values of σ for the Model 3 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The values of R^2 for the Model 3 are shown in Figure 3.61.




Figure 3.61: Values of R^2 for the Model 3 with constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

3.5 Comparison of Models

3.5.1 Constant Intensity

The value of A_1 for the Model 1, Model 2 and Model 3 considering a constant intensity of 2 μJ are shown in Figure 3.62.




Figure 3.62: Values of A_1 for the Model 1, Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan.

The value of τ for the Model 1, Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$ are shown in Figure 3.63.




Figure 3.63: Values of τ for the Model 1, Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan.

The value of σ for the Model 1, Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$ are shown in Figure 3.64.




Figure 3.64: Values of σ for the Model 1, Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan.

The value of t_0 for the Model 1, Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$ are shown in Figure 3.65.




Figure 3.65: Values of t_0 for the Model 1, Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan.

The value of R^2 for the Model 1, Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$ are shown in Figure 3.66.




Figure 3.66: Values of R^2 for the Model 1, Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan.

The value of A_2 for the Model 2 and Model 3 considering a constant energy of 63.6 eV are shown in Figure 3.67.




Figure 3.67: Values of A_2 for the Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan.

The value of τ_2 for the Model 2 and Model 3 considering a constant energy of 63.6 eV are shown in Figure 3.68.




Figure 3.68: Values of τ_2 for the Model 2 and Model 3 considering a constant intensity of $2 \mu\text{J}$. The values are shown as a function of the energy of the scan.

3.5.2 Constant Energy

The value of A_1 for the Model 1, Model 2 and Model 3 considering a constant energy of 63.6 eV are shown in Figure 3.69.




Figure 3.69: Values of A_1 for the Model 1, Model 2 and Model 3 considering a constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The value of τ for the Model 1, Model 2 and Model 3 considering a constant energy of 63.6 eV are shown in Figure 3.70.




Figure 3.70: Values of τ for the Model 1, Model 2 and Model 3 considering a constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The value of σ for the Model 1, Model 2 and Model 3 considering a constant energy of 63.6 eV are shown in Figure 3.71.




Figure 3.71: Values of σ for the Model 1, Model 2 and Model 3 considering a constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The value of t_0 for the Model 1, Model 2 and Model 3 considering a constant energy of 63.6 eV are shown in Figure 3.72.




Figure 3.72: Values of t_0 for the Model 1, Model 2 and Model 3 considering a constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The value of R^2 for the Model 1, Model 2 and Model 3 considering a constant energy of 63.6 eV are shown in Figure 3.73.




Figure 3.73: Values of R^2 for the Model 1, Model 2 and Model 3 considering a constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The value of A_2 for the Model 2 and Model 3 considering a constant energy of 63.6 eV are shown in Figure 3.74.




Figure 3.74: Values of A_2 for the Model 2 and Model 3 considering a constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.

The value of τ_2 for the Model 2 and Model 3 considering a constant energy of 63.6 eV are shown in Figure 3.75.




Figure 3.75: Values of τ_2 for the Model 2 and Model 3 considering a constant energy of 63.6 eV. The values are shown as a function of the intensity of the scan.