

Question **2** ▼ Flag question Mark 0.00 out of 1.00 Incorrect If $y_h(x)$ is a solution of a homogeneous problem, it can always be added to that of a particular solution, therefore there are infinitely many solutions. Select one or more: a. False Careful! ☐ b. True Your answer is incorrect. The correct answer is: True Question **3** Mark 1.00 out of 1.00 Correct Methods of integrating factor and variation of parameters lead to the same solution. Select one or more: ☑ a. Yes ✓ OK, if the solution exists. ☐ b. No Your answer is correct. The correct answer is: Yes Finish review



■ Lecture 8 (Activation)

Quiz)

Grades

Sections

» Welcome

>> Materials

>> STACK

» Assignments

>> Home Exam

Dashboard

Site home

Calendar

Learner Metrics

L Teacher Metrics

» Homework submission

Tuki / Support

Opiskelijoille / Students

- MyCourses instructions for <u>students</u>
- email: mycourses(at)aalto.fi

Opettajille / Teachers

- MyCourses help
- MyTeaching Support form

Palvelusta

- MyCourses rekisteriseloste
- <u>Tietosuojailmoitus</u>
- <u>Palvelukuvaus</u>

Lecture 10 (Activation

Quiz) ►

• <u>Saavutettavuusseloste</u>

About service

- MyCourses protection of <u>privacy</u>
- Privacy notice
- Service description
- Accessibility summary

Service

- MyCourses
- registerbeskrivining • <u>Dataskyddsmeddelande</u>
- <u>Beskrivining av tjänsten</u> • <u>Sammanfattning av</u>
- <u>tillgängligheten</u>

Binh Nguyen (Log out)