


Decision making in case of uncertainties



Bayesian Analysis

- ▶ Based on Bayesian probability theory
 - ▶ uncertainty is presented with probabilities
 - ▶ probabilities are updated based on new information

Bayesian Analysis

- ▶ Based on Bayesian probability theory
 - ▶ uncertainty is presented with probabilities
 - ▶ probabilities are updated based on new information
- ▶ Thomas Bayes (170?–1761)
 - ▶ English nonconformist, Presbyterian minister, mathematician
 - ▶ considered the problem of *inverse probability*
 - ▶ significant part of the Bayesian theory

Bayesian Analysis

- ▶ Based on Bayesian probability theory
 - ▶ uncertainty is presented with probabilities
 - ▶ probabilities are updated based on new information
- ▶ Thomas Bayes (170?–1761)
 - ▶ English nonconformist, Presbyterian minister, mathematician
 - ▶ considered the problem of *inverse probability*
 - ▶ significant part of the Bayesian theory
- ▶ Bayes did not invent all, but was first to solve problem of inverse probability in special case
- ▶ Modern Bayesian theory with rigorous proofs developed in 20th century

Bayesian Analysis

- ▶ Based on Bayesian probability theory
 - ▶ uncertainty is presented with probabilities
 - ▶ probabilities are updated based on new information
- ▶ Thomas Bayes (170?–1761)
 - ▶ English nonconformist, Presbyterian minister, mathematician
 - ▶ considered the problem of *inverse probability*
 - ▶ significant part of the Bayesian theory
- ▶ Bayes did not invent all, but was first to solve problem of inverse probability in special case
- ▶ Modern Bayesian theory with rigorous proofs developed in 20th century
- ▶ A nice book about history: Sharon Bertsch McGrayne, *The Theory That Would Not Die*, 2012.

Term Bayesian used first time in mid 20th century

- ▶ Earlier there was just "probability theory"
 - ▶ concept of the probability was not strictly defined, although it was close to modern Bayesian interpretation
 - ▶ in the end of 19th century there were increasing demand for more strict definition of probability (mathematical and philosophical problem)

Term Bayesian used first time in mid 20th century

- ▶ Earlier there was just "probability theory"
 - ▶ concept of the probability was not strictly defined, although it was close to modern Bayesian interpretation
 - ▶ in the end of 19th century there were increasing demand for more strict definition of probability (mathematical and philosophical problem)
- ▶ In the beginning of 20th century frequentist view gained popularity
 - ▶ accepts definition of probabilities only through frequencies
 - ▶ does not accept inverse probability or use of prior
 - ▶ gained popularity due to apparent objectivity and "cook book" like reference books

Term Bayesian used first time in mid 20th century

- ▶ Earlier there was just "probability theory"
 - ▶ concept of the probability was not strictly defined, although it was close to modern Bayesian interpretation
 - ▶ in the end of 19th century there were increasing demand for more strict definition of probability (mathematical and philosophical problem)
- ▶ In the beginning of 20th century frequentist view gained popularity
 - ▶ accepts definition of probabilities only through frequencies
 - ▶ does not accept inverse probability or use of prior
 - ▶ gained popularity due to apparent objectivity and "cook book" like reference books
- ▶ R. A. Fisher used in 1950 first time term "Bayesian" to emphasize the difference to general term "probability theory"
 - ▶ term became quickly popular, because alternative descriptions were longer

Uncertainty and probabilistic modeling

- ▶ Two types of uncertainty: aleatoric and epistemic
- ▶ Representing uncertainty with probabilities
- ▶ Updating uncertainty

Two types of uncertainty

- ▶ Aleatoric uncertainty due to randomness
- ▶ Epistemic uncertainty due to lack of knowledge

Two types of uncertainty

- ▶ Aleatoric uncertainty due to randomness
 - ▶ we are not able to obtain observations which could reduce this uncertainty
- ▶ Epistemic uncertainty due to lack of knowledge

Two types of uncertainty

- ▶ Aleatoric uncertainty due to randomness
 - ▶ we are not able to obtain observations which could reduce this uncertainty
- ▶ Epistemic uncertainty due to lack of knowledge
 - ▶ we are able to obtain observations which can reduce this uncertainty
 - ▶ two observers may have different epistemic uncertainty

Updating uncertainty

Updating uncertainty

- Probability of red $\frac{\text{\#red}}{\text{\#red} + \text{\#yellow}} = \theta$

Updating uncertainty

- ▶ Probability of red $\frac{\text{\#red}}{\text{\#red} + \text{\#yellow}} = \theta$
- ▶ $p(y = \text{\#red} | \theta) = \theta$ aleatoric uncertainty

Updating uncertainty

- ▶ Probability of red $\frac{\text{\#red}}{\text{\#red} + \text{\#yellow}} = \theta$
- ▶ $p(y = \text{\#red} | \theta) = \theta$ aleatoric uncertainty
- ▶ $p(\theta)$ epistemic uncertainty

Updating uncertainty

- ▶ Probability of red $\frac{\text{\#red}}{\text{\#red} + \text{\#yellow}} = \theta$
- ▶ $p(y = \text{\#red} | \theta) = \theta$ aleatoric uncertainty
- ▶ $p(\theta)$ epistemic uncertainty
- ▶ Picking many chips updates our uncertainty about the proportion
- ▶ $p(\theta | y = \text{\#red}, \text{\#yellow}, \text{\#red}, \text{\#red}, \dots) = ?$

Updating uncertainty

- ▶ Probability of red $\frac{\text{\#red}}{\text{\#red} + \text{\#yellow}} = \theta$
- ▶ $p(y = \text{\#red} | \theta) = \theta$ aleatoric uncertainty
- ▶ $p(\theta)$ epistemic uncertainty
- ▶ Picking many chips updates our uncertainty about the proportion
- ▶ $p(\theta | y = \text{\#red}, \text{\#yellow}, \text{\#red}, \text{\#red}, \dots) = ?$
- ▶ Bayes rule $p(\theta | y) = \frac{p(y|\theta)p(\theta)}{\int p(y|\theta)p(\theta)d\theta}$

Model vs. likelihood


- ▶ Bayes rule $p(\theta|y) \propto p(y|\theta)p(\theta)$
- ▶ Model: $p(y|\theta)$ as a function of y given fixed θ describes the aleatoric uncertainty
- ▶ Likelihood: $p(y|\theta)$ as a function of θ given fixed y provides information about epistemic uncertainty, but is not a probability distribution

Model vs. likelihood

- ▶ Bayes rule $p(\theta|y) \propto p(y|\theta)p(\theta)$
- ▶ Model: $p(y|\theta)$ as a function of y given fixed θ describes the aleatoric uncertainty
- ▶ Likelihood: $p(y|\theta)$ as a function of θ given fixed y provides information about epistemic uncertainty, but is not a probability distribution
- ▶ Bayes rule combines the likelihood with prior uncertainty $p(\theta)$ and transforms them to updated posterior uncertainty


Example of uncertainty in modeling

Data




Example of uncertainty in modeling

Posterior mean




Example of uncertainty in modeling

Predictive distribution given posterior mean




Example of uncertainty in modeling

Posterior draws




Example of uncertainty in modeling

Posterior draws and predictive distribution




Example of uncertainty in modeling

Posterior draws and predictive distribution



Example application: Drug dosage for liver transplant¹


- ▶ Everolimus is immunosuppressant to prevent rejection of organ transplants
- ▶ Pharmacokinetic model of drug and body, optimal dosage depends on weight



¹with E. Siivola, Aalto and S. Weber, Novartis Pharma

Example application: Drug dosage for liver transplant¹

- ▶ Everolimus is immunosuppressant to prevent rejection of organ transplants
- ▶ Pharmacokinetic model of drug and body, optimal dosage depends on weight




- ▶ Model fitted with 500 adults, extrapolation to children?

¹with E. Siivola, Aalto and S. Weber, Novartis Pharma

Example application: Drug dosage for liver transplant¹

- ▶ Everolimus is immunosuppressant to prevent rejection of organ transplants
- ▶ Pharmacokinetic model of drug and body, optimal dosage depends on weight



- ▶ Model fitted with 500 adults, extrapolation to children?
- ▶ Maturation effect, 17 observations from children

¹with E. Siivola, Aalto and S. Weber, Novartis Pharma

The art of probabilistic modeling

- ▶ The art of probabilistic modeling is to describe in a mathematical form (model and prior distributions) what we already know and what we don't know

The art of probabilistic modeling

- ▶ The art of probabilistic modeling is to describe in a mathematical form (model and prior distributions) what we already know and what we don't know
- ▶ “Easy” part is to use Bayes rule to update the uncertainties
 - ▶ computational challenges

The art of probabilistic modeling

- ▶ The art of probabilistic modeling is to describe in a mathematical form (model and prior distributions) what we already know and what we don't know
- ▶ “Easy” part is to use Bayes rule to update the uncertainties
 - ▶ computational challenges
- ▶ Other parts of the art of probabilistic modeling are, for example,
 - ▶ model checking: is data in conflict with our prior knowledge?
 - ▶ presentation: presenting the model and the results to the application experts

- ▶ Galaxy clusters for cosmology
- ▶ Coagulation of blood
- ▶ Gene regulation
- ▶ Pharmacokinetics and -dynamics
- ▶ Decision support
- ▶ Effects of nutrition for diabetes
- ▶ Evolutionary anthropology
- ▶ Clinical trial designs
- ▶ Daily demand for gas
- ▶ Brain structure trees
- ▶ School enrollment
- ▶ Sports
- ▶ Product demand
- ▶ Cocoa bean fermentation
- ▶ Marine propulsion power
- ▶ Alcohol consumption trends
- ▶ Flood probability
- ▶ Instantaneous heart rate distributions
- ▶ Drug dosing regimens in pediatrics
- ▶ Human T stem cell memory cells
- ▶ Fairness in university admission policies
- ▶ Destruction of bacteria and bacterial spores under heat

Bayesian data analysis

Example analyses

- ▶ Treatment/control
 - ▶ randomize patients to treatment or control
 - ▶ is the treatment effective?

Bayesian data analysis

Example analyses

- ▶ Treatment/control
 - ▶ randomize patients to treatment or control
 - ▶ is the treatment effective?
- ▶ Continuous valued treatment
 - ▶ randomize patients with different dosages
 - ▶ which dosage is sufficient without too many side effects?

Bayesian data analysis

Example analyses

- ▶ Treatment/control
 - ▶ randomize patients to treatment or control
 - ▶ is the treatment effective?
- ▶ Continuous valued treatment
 - ▶ randomize patients with different dosages
 - ▶ which dosage is sufficient without too many side effects?
- ▶ Different effects for different patients?
 - ▶ Is the treatment effect different for male/female, child/adult, light/heavy, ...

Bayesian approach

- ▶ Benefits of Bayesian approach
 - ▶ integrate over uncertainties to focus to interesting parts
 - ▶ use relevant prior information
 - ▶ hierarchical models
 - ▶ model checking and evaluation

Computation

We need to be able to compute expectations with respect to posterior distribution $p(\theta|y)$

$$E_{\theta|y} [g(\theta)] = \int p(\theta|y) g(\theta) d\theta$$

- ▶ Analytic
 - ▶ only for very simple models
- ▶ Monte Carlo, Markov chain Monte Carlo
 - ▶ generic
- ▶ Distributional approximations
 - ▶ e.g. Laplace, variational, expectation propagation
 - ▶ less generic, but can be much faster with sufficient accuracy

Probabilistic programming



Enables agile workflow for developing probabilistic models

language – automated inference – diagnostics



mc-stan.org

Binomial model for treatment/control comparison

- ▶ Two groups of patients: treatment and control
 - ▶ Binary outcome
 - ▶ Is the treatment useful?

Binomial model for treatment/control comparison

```
parameters {  
  real<lower=0,upper=1> theta1;  
  real<lower=0,upper=1> theta2;  
}  
model {  
  theta1 ~ beta(1, 1);  
  theta2 ~ beta(1, 1);  
  y1 ~ binomial(N1, theta1);  
  y2 ~ binomial(N2, theta2);  
}
```

Binomial model for treatment/control comparison

RStanARM

```
fit_bin2 <- stan_glm(y/N ~ grp2, family = binomial(),  
                    data = d_bin2, weights = N)
```

Modeling nature

- ▶ Drop a ball from different heights and measure time

Modeling nature

- ▶ Drop a ball from different heights and measure time
 - ▶ Newton
 - ▶ air resistance, air pressure, shape and surface structure of the ball
 - ▶ relativity

Modeling nature

- ▶ Drop a ball from different heights and measure time
 - ▶ Newton
 - ▶ air resistance, air pressure, shape and surface structure of the ball
 - ▶ relativity
- ▶ Taking into account the accuracy of the measurements, how accurate model is needed?

Modeling nature

- ▶ Drop a ball from different heights and measure time
 - ▶ Newton
 - ▶ air resistance, air pressure, shape and surface structure of the ball
 - ▶ relativity
- ▶ Taking into account the accuracy of the measurements, how accurate model is needed?
 - ▶ often simple models are adequate and useful
 - ▶ *All models are wrong, but some of them are useful*, George P. Box

Reminder: Uncertainty and probabilistic modeling

- ▶ Two types of uncertainty: aleatoric and epistemic
- ▶ Representing uncertainty with probabilities
- ▶ Updating uncertainty

Questions

- ▶ Pick a number between 1–5

Questions

- ▶ Pick a number between 1–5
 - ▶ raise as many fingers

Questions

- ▶ Pick a number between 1–5
 - ▶ raise as many fingers
 - ▶ is the number of fingers raised random (by you or by others)?

Questions

- ▶ Pick a number between 1–5
 - ▶ raise as many fingers
 - ▶ is the number of fingers raised random (by you or by others)?
- ▶ If we build a robot with very fast vision which can observe the rotating coin accurately, is the throw random for the robot?

Questions

- ▶ Pick a number between 1–5
 - ▶ raise as many fingers
 - ▶ is the number of fingers raised random (by you or by others)?
- ▶ If we build a robot with very fast vision which can observe the rotating coin accurately, is the throw random for the robot?
- ▶ Is the quantum uncertainty aleatoric or epistemic?

Questions

- ▶ Pick a number between 1–5
 - ▶ raise as many fingers
 - ▶ is the number of fingers raised random (by you or by others)?
- ▶ If we build a robot with very fast vision which can observe the rotating coin accurately, is the throw random for the robot?
- ▶ Is the quantum uncertainty aleatoric or epistemic?
- ▶ What is your own example with both aleatoric and epistemic uncertainty?

Rest of the course

- ▶ Basic models which can be used as building blocks
- ▶ Basic computation
- ▶ Typical simple scientific data analysis cases
 - ▶ e.g. comparison of treatments
- ▶ Presentation of the results

Some important terms

- probability
- probability density
- probability mass
- probability density function (pdf)
- probability mass function (pmf)
- probability distribution
- discrete probability distribution
- continuous probability distribution
- cumulative distribution function (cdf)
- likelihood

Ambiguous notation in statistics

$$\ln p(y|\theta)$$

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value

we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value
we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$
- θ can be variable or value
we could clarify by using $p(y|\Theta)$ or $p(y|\theta)$

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value
we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$
- θ can be variable or value
we could clarify by using $p(y|\Theta)$ or $p(y|\theta)$
- p can be a discrete or continuous function of y or θ
we could clarify by using P_Y , P_Θ , p_Y or p_Θ

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value
we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$
- θ can be variable or value
we could clarify by using $p(y|\Theta)$ or $p(y|\theta)$
- p can be a discrete or continuous function of y or θ
we could clarify by using P_Y , P_Θ , p_Y or p_Θ
- $P_Y(Y|\Theta = \theta)$ is a probability mass function, sampling distribution, observation model

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value
we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$
- θ can be variable or value
we could clarify by using $p(y|\Theta)$ or $p(y|\theta)$
- p can be a discrete or continuous function of y or θ
we could clarify by using P_Y , P_Θ , p_Y or p_Θ
- $P_Y(Y|\Theta = \theta)$ is a probability mass function, sampling distribution, observation model
- $P(Y = y|\Theta = \theta)$ is a probability

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value
we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$
- θ can be variable or value
we could clarify by using $p(y|\Theta)$ or $p(y|\theta)$
- p can be a discrete or continuous function of y or θ
we could clarify by using P_Y , P_Θ , p_Y or p_Θ
- $P_Y(Y|\Theta = \theta)$ is a probability mass function, sampling distribution, observation model
- $P(Y = y|\Theta = \theta)$ is a probability
- $P_\Theta(Y = y|\Theta)$ is a likelihood function (can be discrete or continuous)

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value
we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$
- θ can be variable or value
we could clarify by using $p(y|\Theta)$ or $p(y|\theta)$
- p can be a discrete or continuous function of y or θ
we could clarify by using P_Y , P_Θ , p_Y or p_Θ
- $P_Y(Y|\Theta = \theta)$ is a probability mass function, sampling distribution, observation model
- $P(Y = y|\Theta = \theta)$ is a probability
- $P_\Theta(Y = y|\Theta)$ is a likelihood function (can be discrete or continuous)
- $p_Y(Y|\Theta = \theta)$ is a probability density function, sampling distribution, observation model

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value
we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$
- θ can be variable or value
we could clarify by using $p(y|\Theta)$ or $p(y|\theta)$
- p can be a discrete or continuous function of y or θ
we could clarify by using P_Y , P_Θ , p_Y or p_Θ
- $P_Y(Y|\Theta = \theta)$ is a probability mass function, sampling distribution, observation model
- $P(Y = y|\Theta = \theta)$ is a probability
- $P_\Theta(Y = y|\Theta)$ is a likelihood function (can be discrete or continuous)
- $p_Y(Y|\Theta = \theta)$ is a probability density function, sampling distribution, observation model
- $p(Y = y|\Theta = \theta)$ is a density

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value
we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$
- θ can be variable or value
we could clarify by using $p(y|\Theta)$ or $p(y|\theta)$
- p can be a discrete or continuous function of y or θ
we could clarify by using P_Y , P_Θ , p_Y or p_Θ
- $P_Y(Y|\Theta = \theta)$ is a probability mass function, sampling distribution, observation model
- $P(Y = y|\Theta = \theta)$ is a probability
- $P_\Theta(Y = y|\Theta)$ is a likelihood function (can be discrete or continuous)
- $p_Y(Y|\Theta = \theta)$ is a probability density function, sampling distribution, observation model
- $p(Y = y|\Theta = \theta)$ is a density
- $p_\Theta(Y = y|\Theta)$ is a likelihood function (can be discrete or continuous)

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value
we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$
- θ can be variable or value
we could clarify by using $p(y|\Theta)$ or $p(y|\theta)$
- p can be a discrete or continuous function of y or θ
we could clarify by using P_Y , P_Θ , p_Y or p_Θ
- $P_Y(Y|\Theta = \theta)$ is a probability mass function, sampling distribution, observation model
- $P(Y = y|\Theta = \theta)$ is a probability
- $P_\Theta(Y = y|\Theta)$ is a likelihood function (can be discrete or continuous)
- $p_Y(Y|\Theta = \theta)$ is a probability density function, sampling distribution, observation model
- $p(Y = y|\Theta = \theta)$ is a density
- $p_\Theta(Y = y|\Theta)$ is a likelihood function (can be discrete or continuous)
- y and θ can also be mix of continuous and discrete

Ambiguous notation in statistics

In $p(y|\theta)$

- y can be variable or value
we could clarify by using $p(Y|\theta)$ or $p(y|\theta)$
- θ can be variable or value
we could clarify by using $p(y|\Theta)$ or $p(y|\theta)$
- p can be a discrete or continuous function of y or θ
we could clarify by using P_Y , P_Θ , p_Y or p_Θ
- $P_Y(Y|\Theta = \theta)$ is a probability mass function, sampling distribution, observation model
- $P(Y = y|\Theta = \theta)$ is a probability
- $P_\Theta(Y = y|\Theta)$ is a likelihood function (can be discrete or continuous)
- $p_Y(Y|\Theta = \theta)$ is a probability density function, sampling distribution, observation model
- $p(Y = y|\Theta = \theta)$ is a density
- $p_\Theta(Y = y|\Theta)$ is a likelihood function (can be discrete or continuous)
- y and θ can also be mix of continuous and discrete
- Due to the sloppiness sometimes likelihood is used to refer $P_{Y,\theta}(Y|\Theta)$, $p_{Y,\theta}(Y|\Theta)$

Chapter 1

Reading instructions

- ▶ 1.1-1.3 important terms
- ▶ 1.4 a useful example
- ▶ 1.5 foundations
- ▶ 1.6 & 1.7 examples (can be skipped, but may be useful to read)
- ▶ 1.8 & 1.9 background material, good to read before doing the exercises
- ▶ 1.10 a point of view for using Bayesian inference