

MS-A0001 - Matrix Algebra,
26.10.2020-08.12.2020

Grades

Sections

» Welcome

» Materials

» Assignments

» STACK

» Homework Submission

» ONLINE EXAM

» END EXAM 8.12

Dashboard

Site home

Calendar

Learner Metrics

Teacher Metrics

MS-A0001 - Matrix Algebra, 26.10.2020-08.12.2020

Dashboard / Courses / School of Science / department of... / ms-a0001 - ma... / Sections / STACK / Lecture 5

Started on	Friday, 13 November 2020, 3:27 PM
State	Finished
Completed on	Friday, 13 November 2020, 3:48 PM
Time taken	21 mins 33 secs
Grade	2.00 out of 2.00 (100%)

Question 1

Flag question Mark 1.00 out of 1.00 Correct

Compute the inverse of

$$A = \begin{bmatrix} 3 & -1 \\ -2 & -2 \end{bmatrix}$$

using Gauss-Jordan elimination.

Input the augmented matrix $[A|I_2]$:

3	-1	1	0
-2	-2	0	1

Your last answer was interpreted as follows:

$$\begin{bmatrix} 3 & -1 & 1 & 0 \\ -2 & -2 & 0 & 1 \end{bmatrix}$$

Next compute the reduced row echelon (rref) of the matrix. Input the matrix rref $[A|I_2]$:

1	0	1/4	-1/8
0	1	-1/4	-3/8

Your last answer was interpreted as follows:

$$\begin{bmatrix} 1 & 0 & \frac{1}{4} & -\frac{1}{8} \\ 0 & 1 & -\frac{1}{4} & -\frac{3}{8} \end{bmatrix}$$

From the previous matrix we may extract the inverse of A. Input the sought A^{-1} .

1/4	-1/8
-1/4	-3/8

Your last answer was interpreted as follows:

$$\begin{bmatrix} \frac{1}{4} & -\frac{1}{8} \\ -\frac{1}{4} & -\frac{3}{8} \end{bmatrix}$$

Your answer is correct!

Your answer is correct!

The augmented matrix is correct!

Marks for this submission: 0.33/0.33.

Your answer is correct!

The reduced row echelon form of the augmented matrix is correct!

Marks for this submission: 0.33/0.33.

Your answer is correct!

The inverse A^{-1} is correct!

Marks for this submission: 0.33/0.33.

Worked solution:

First form a matrix where the left half is the matrix whose inverse we wish to calculate and the right half is an identity matrix of the same dimensions.

$$[A|I_2] = \begin{bmatrix} 3 & -1 & 1 & 0 \\ -2 & -2 & 0 & 1 \end{bmatrix}$$

Next we apply row operations of the matrix until the left half is an identity matrix.

$$\begin{aligned} \begin{bmatrix} 3 & -1 & 1 & 0 \\ -2 & -2 & 0 & 1 \end{bmatrix} &\xrightarrow{\frac{1}{3}R_1} \begin{bmatrix} 1 & -\frac{1}{3} & \frac{1}{3} & 0 \\ -2 & -2 & 0 & 1 \end{bmatrix} \\ &\xrightarrow{R_2+2R_1} \begin{bmatrix} 1 & -\frac{1}{3} & \frac{1}{3} & 0 \\ 0 & -\frac{8}{3} & \frac{2}{3} & 1 \end{bmatrix} \\ &\xrightarrow{\left(-\frac{3}{8}\right)R_2} \begin{bmatrix} 1 & -\frac{1}{3} & \frac{1}{3} & 0 \\ 0 & 1 & -\frac{1}{4} & -\frac{3}{8} \end{bmatrix} \\ &\xrightarrow{R_1+\frac{1}{3}R_2} \begin{bmatrix} 1 & 0 & \frac{1}{4} & -\frac{1}{8} \\ 0 & 1 & -\frac{1}{4} & -\frac{3}{8} \end{bmatrix} \end{aligned}$$

Now the matrix is in reduced row echelon form (rref) and its left block is I_2 . The right block must now be the sought inverse:

$$A^{-1} = \begin{bmatrix} \frac{1}{4} & -\frac{1}{8} \\ -\frac{1}{4} & -\frac{3}{8} \end{bmatrix}.$$

A correct answer is $\begin{bmatrix} 3 & -1 & 1 & 0 \\ -2 & -2 & 0 & 1 \end{bmatrix}$.

A correct answer is $\begin{bmatrix} 1 & 0 & \frac{1}{4} & -\frac{1}{8} \\ 0 & 1 & -\frac{1}{4} & -\frac{3}{8} \end{bmatrix}$.

A correct answer is $\begin{bmatrix} \frac{1}{4} & -\frac{1}{8} \\ -\frac{1}{4} & -\frac{3}{8} \end{bmatrix}$.

Question 2

Flag question Mark 1.00 out of 1.00 Correct

Let

$$A = \begin{bmatrix} 0 & 3 & 7 \\ 5 & 6 & 0 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 6 & 4 \\ 7 & 4 \end{bmatrix}.$$

Calculate $A^T B$.

35	20
60	36
42	28

Your last answer was interpreted as follows:

$$\begin{bmatrix} 35 & 20 \\ 60 & 36 \\ 42 & 28 \end{bmatrix}$$

Your answer is correct!

Marks for this submission: 1.00/1.00.

Worked solution:

The transpose of A is

$$A^T = \begin{bmatrix} 0 & 5 \\ 3 & 6 \\ 7 & 0 \end{bmatrix}.$$

Let's compute the elements of the product next. The element on the ith row and jth column of $A^T B$ can be calculated by taking the dot product of the ith row of A^T and the jth column of B. The element in the first row and first column is then

$$(A^T B)_{11} = (0, 5) \cdot (6, 7) = 0 \cdot 6 + 5 \cdot 7 = 35$$

and the element in the first row and second column is

$$(A^T B)_{12} = (0, 5) \cdot (4, 4) = 0 \cdot 4 + 5 \cdot 4 = 20.$$

The rest of the elements can be similarly calculated. The answer is

$$A^T B = \begin{bmatrix} 35 & 20 \\ 60 & 36 \\ 42 & 28 \end{bmatrix}.$$

A correct answer is $\begin{bmatrix} 35 & 20 \\ 60 & 36 \\ 42 & 28 \end{bmatrix}$.

Quiz navigation

1	2
✓	✓

Finish review

◀ Lecture 4

Lecture 6 ▶



Tuki / Support

Opiskelijoille / Students

- MyCourses instructions for students
- email: mycourses(at)aalto.fi

Opettajille / Teachers

- MyCourses help
- MyTeaching Support form

Palvelusta

- MyCourses rekisteriseloste
- Tietosuojailmoitus
- Palvelukuvaus
- Saavutettavuusseloste

About service

- MyCourses protection of privacy
- Privacy notice
- Service description
- Accessibility summary

Service

- MyCourses registerbeskrivning
- Data skyddsmeddelande
- Beskrivning av tjänsten
- Sammanfattning av tillgängligheten

