

# CS-C3240 – Machine Learning D

## Round 3: From features to classification


Stephan Sigg

Department of Communications and Networking  
Aalto University, School of Electrical Engineering  
[stephan.sigg@aalto.fi](mailto:stephan.sigg@aalto.fi)

Version 2.3, February 12, 2022

# Issues related to high dimensional input data

Exponential growth Volume of the space grows exponentially with dimension




# Issues related to high dimensional input data

Exponential growth Volume of the space grows exponentially with dimension

## Curse of dimensionality

Too sparse samples across regions to estimate a distribution in that space  
(Problematic for methods that require statistical significance)



# Issues related to high dimensional input data


**Exponential growth** Volume of the space grows exponentially with dimension

## Curse of dimensionality

Too sparse samples across regions to estimate a distribution in that space  
(Problematic for methods that require statistical significance)

## Hughes (peaking) phenomenon

Predictive power of classifier first increases with dimension, then decreases




# Issues related to high dimensional input data



Exponential growth Volume of the space grows exponentially with dimension

Counter-intuitive properties in higher dimensional spaces



# Issues related to high dimensional input data

Exponential growth Volume of the space grows exponentially with dimension

Counter-intuitive properties in higher dimensional spaces

## Example – Volume of a sphere

Consider a sphere of radius  $r = 1$  in a  $D$ -dimensional space

# Issues related to high dimensional input data

Exponential growth Volume of the space grows exponentially with dimension

Counter-intuitive properties in higher dimensional spaces

## Example – Volume of a sphere

Consider a sphere of radius  $r = 1$  in a  $D$ -dimensional space

Fraction of the volume between radius  $r = 1$  and  $r' = 1 - \epsilon$ ?

# Issues related to high dimensional input data

Exponential growth Volume of the space grows exponentially with dimension

Counter-intuitive properties in higher dimensional spaces

## Example – Volume of a sphere

Consider a sphere of radius  $r = 1$  in a  $D$ -dimensional space

Fraction of the volume between radius  $r = 1$  and  $r' = 1 - \epsilon$ ?

Volume of sphere with radius  $r$ :

$$V_D(r) = \delta_D r^D \quad \text{for appropriate } \delta_D$$



# Issues related to high dimensional input data

Exponential growth Volume of the space grows exponentially with dimension

Counter-intuitive properties in higher dimensional spaces

## Example – Volume of a sphere

Consider a sphere of radius  $r = 1$  in a  $D$ -dimensional space

Fraction of the volume between radius  $r = 1$  and  $r' = 1 - \varepsilon$ ?

Volume of sphere with radius  $r$ :

$$V_D(r) = \delta_D r^D \quad \text{for appropriate } \delta_D$$

Given by

$$\frac{V_D(1) - V_D(1 - \varepsilon)}{V_D(1)} = 1 - (1 - \varepsilon)^D$$

# Issues related to high dimensional input data

Exponential growth Volume of the space grows exponentially with dimension

Counter-intuitive properties in higher dimensional spaces

## Example – Volume of a sphere

Consider a sphere of radius  $r = 1$  in a  $D$ -dimensional space

Fraction of the volume between radius  $r = 1$  and  $r' = 1 - \varepsilon$ ?

Volume of sphere with radius  $r$ :

$$V_D(r) = \delta_D r^D \quad \text{for appropriate } \delta_D$$

Given by

$$\frac{V_D(1) - V_D(1 - \varepsilon)}{V_D(1)} = 1 - (1 - \varepsilon)^D$$

For large  $D$ , this fraction tends to 1

In high dimensions, most of the volume of a sphere concentrates near the surface

# Issues related to high dimensional input data

**Exponential growth** Volume of the space grows exponentially with dimension

**Counter-intuitive properties** in higher dimensional spaces

## Example – Volume of a sphere

Consider a sphere of radius  $r = 1$  in a  $D$ -dimensional space


**Fraction of the volume between radius  $r = 1$  and  $r' = 1 - \epsilon$ ?**

Volume of sphere with radius  $r$ :

$$V_D(r) = \delta_D r^D \quad \text{for appropriate } \delta_D$$

Given by

$$\frac{V_D(1) - V_D(1 - \epsilon)}{V_D(1)} = 1 - (1 - \epsilon)^D$$



For large  $D$ , this fraction tends to 1

In high dimensions, most of the volume of a sphere concentrates near the surface

# Issues related to high dimensional input data



**Exponential growth** Volume of the space grows exponentially with dimension

**Counter-intuitive properties** in higher dimensional spaces

Example – Gaussian distribution

Probability mass concentrated in a thin shell

(here plotted as distance from the origin in a polar coordinate system)



# Issues related to high dimensional input data


**Exponential growth** Volume of the space grows exponentially with dimension

**Counter-intuitive properties** in higher dimensional spaces

Example – Gaussian distribution

Probability mass concentrated in a thin shell

(here plotted as distance from the origin in a polar coordinate system)



## Curse of Dimensionality

Mechanisms to efficiently reduce dimensions or classifiers that respect properties of high-dimensional spaces required.

# Questions?

Stephan Sigg

`stephan.sigg@aalto.fi`

Si Zuo

`si.zuo@aalto.fi`

# Literature

- C.M. Bishop: Pattern recognition and machine learning, Springer, 2007.
- R.O. Duda, P.E. Hart, D.G. Stork: Pattern Classification, Wiley, 2001.

