

CS-C3240 - Machine Learning D

Feature Engineering

Stephan Sigg

Department of Communications and Networking Aalto University, School of Electrical Engineering stephan.sigg@aalto.fi

Version 1.0, February 14, 2022

For two features \mathcal{X}_1 , \mathcal{X}_2 , and the arithmetic mean $E[\cdot]$, consider sets of measurements with zero mean:

$$\overrightarrow{x_1} = \{x_1^{(1)}, \dots, x_1^{(n)}\}\ \overrightarrow{x_2} = \{x_2^{(1)}, \dots, x_2^{(n)}\}\$$



For two features \mathcal{X}_1 , \mathcal{X}_2 , and the arithmetic mean $E[\cdot]$, consider sets of measurements with zero mean:

$$\overrightarrow{x_1} = \{x_1^{(1)}, \dots, x_1^{(n)}\}\$$
 $\overrightarrow{x_2} = \{x_2^{(1)}, \dots, x_2^{(n)}\}\$

Variance:
$$E[(\overrightarrow{x_i} - E[\overrightarrow{x_i}])^2] = E[(\overrightarrow{x_i} - \mu)^2]$$



For two features $\mathcal{X}_1, \mathcal{X}_2$, and the arithmetic mean $E[\cdot]$, consider sets of measurements with zero mean:

$$\overrightarrow{x_1} = \{x_1^{(1)}, \dots, x_1^{(n)}\}\ \overrightarrow{x_2} = \{x_2^{(1)}, \dots, x_2^{(n)}\}\$$

Variance:
$$E[(\overrightarrow{x_j} - E[\overrightarrow{x_j}])^2] = E[(\overrightarrow{x_j} - \mu)^2]$$
 $\xrightarrow{\text{zero mean}} E[\overrightarrow{x_j} \cdot \overrightarrow{x_j}]$



For two features \mathcal{X}_1 , \mathcal{X}_2 , and the arithmetic mean $E[\cdot]$, consider sets of measurements with zero mean:

$$\overrightarrow{x_1} = \{x_1^{(1)}, \dots, x_1^{(n)}\}\ \overrightarrow{x_2} = \{x_2^{(1)}, \dots, x_2^{(n)}\}\$$

Variance:
$$E[(\overrightarrow{x_j} - E[\overrightarrow{x_j}])^2] = E[(\overrightarrow{x_j} - \mu)^2]$$

$$\xrightarrow{\text{zero mean}} E[\overrightarrow{x_j} \cdot \overrightarrow{x_j}]$$

$$\Rightarrow \frac{1}{n} \sum_{i=1}^{n} (\overrightarrow{x_i} \cdot \overrightarrow{x_i})$$



For two features \mathcal{X}_1 , \mathcal{X}_2 , and the arithmetic mean $E[\cdot]$, consider sets of measurements with zero mean:

$$\overrightarrow{x_1} = \{x_1^{(1)}, \dots, x_1^{(n)}\}\$$
 $\overrightarrow{x_2} = \{x_2^{(1)}, \dots, x_2^{(n)}\}\$

Variance:
$$E[(\overrightarrow{x_j} - E[\overrightarrow{x_j}])^2] = E[(\overrightarrow{x_j} - \mu)^2]$$

$$\xrightarrow{\text{zero mean}} E[\overrightarrow{x_j} \cdot \overrightarrow{x_j}]$$


$$\Rightarrow \frac{1}{n} \sum_{i=1}^{n} (\overrightarrow{x_i} \cdot \overrightarrow{x_i})$$

Covariance: $E[(\overrightarrow{x_i} - E[\overrightarrow{x_i}])(\overrightarrow{x_i} - E[\overrightarrow{x_i}])]$





Covariance matrix







Questions?

Stephan Sigg stephan.sigg@aalto.fi

Si Zuo si.zuo@aalto.fi





Literature

- C.M. Bishop: Pattern recognition and machine learning, Springer, 2007.
- R.O. Duda, P.E. Hart, D.G. Stork: Pattern Classification, Wiley, 2001.

