



Aalto University

# MS-C1350 Partial differential equations

## Chapter 2.3 Fourier series

Riikka Korte

Department of Mathematics and Systems Analysis  
Aalto University  
riikka.korte@aalto.fi

September 3, 2024

# Fourier series

- ▶ Let  $e_j = e^{ij t} \in L^1([-\pi, \pi])$ .

# Fourier series

- ▶ Let  $e_j = e^{ij t} \in L^1([-\pi, \pi])$ .
- ▶ Recall that  $\{e_j\}_{j=-\infty}^{\infty}$  form an orthonormal set with the inner product given in previous slides.

# Fourier series

- ▶ Let  $e_j = e^{ij t} \in L^1([-\pi, \pi])$ .
- ▶ Recall that  $\{e_j\}_{j=-\infty}^{\infty}$  form an orthonormal set with the inner product given in previous slides.
- ▶ Let  $f \in L^1([-\pi, \pi])$ . The  $n$ th partial sum of a Fourier series is

$$S_n f(t) = \sum_{j=-n}^n \widehat{f}(j) e^{ij t}, \quad n = 0, 1, 2, \dots,$$

where

$$\widehat{f}(j) = \langle f, e_j \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-ij t} dt, \quad j \in \mathbb{Z},$$

is the  $j$ th Fourier coefficient of  $f$ .

# Fourier series

- ▶ Let  $e_j = e^{ijt} \in L^1([-\pi, \pi])$ .
- ▶ Recall that  $\{e_j\}_{j=-\infty}^{\infty}$  form an orthonormal set with the inner product given in previous slides.
- ▶ Let  $f \in L^1([-\pi, \pi])$ . The  $n$ th partial sum of a Fourier series is

$$S_n f(t) = \sum_{j=-n}^n \hat{f}(j) e^{ijt}, \quad n = 0, 1, 2, \dots,$$

where

$$\hat{f}(j) = \langle f, e_j \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-ijt} dt, \quad j \in \mathbb{Z},$$

is the  $j$ th Fourier coefficient of  $f$ .

- ▶ The Fourier series of  $f$  is the limit of the partial sums  $S_n f$  as  $n \rightarrow \infty$ , provided the limit exists in some reasonable sense. In this case we may write

$$f(t) = \lim_{n \rightarrow \infty} S_n f(t) = \lim_{n \rightarrow \infty} \sum_{j=-n}^n \hat{f}(j) e^{ijt} = \sum_{j=-\infty}^{\infty} \hat{f}(j) e^{ijt}.$$

# Remarks about Fourier series

- ▶ Definition makes sense if  $f \in L^1([-\pi, \pi])$ , but we will see that  $L^2([-\pi, \pi])$  is needed to understand the Fourier coefficients and the convergence of the series.

# Remarks about Fourier series

- ▶ Definition makes sense if  $f \in L^1([-\pi, \pi])$ , but we will see that  $L^2([-\pi, \pi])$  is needed to understand the Fourier coefficients and the convergence of the series.
- ▶ We always consider symmetric Fourier series (sum from  $-n$  to  $n$ ).

# Remarks about Fourier series

- ▶ Definition makes sense if  $f \in L^1([-\pi, \pi])$ , but we will see that  $L^2([-\pi, \pi])$  is needed to understand the Fourier coefficients and the convergence of the series.
- ▶ We always consider symmetric Fourier series (sum from  $-n$  to  $n$ ).
- ▶ It corresponds to having trigonometric functions  $\sin(t), \cos(t), \sin(2t), \cos(2t), \dots, \sin(nt), \cos(nt)$ .



# Remarks about Fourier series

- ▶ Definition makes sense if  $f \in L^1([-\pi, \pi])$ , but we will see that  $L^2([-\pi, \pi])$  is needed to understand the Fourier coefficients and the convergence of the series.
- ▶ We always consider symmetric Fourier series (sum from  $-n$  to  $n$ ).
- ▶ It corresponds to having trigonometric functions  $\sin(t), \cos(t), \sin(2t), \cos(2t), \dots, \sin(nt), \cos(nt)$ .
- ▶  $e_j, j \in \mathbb{Z}$  is always  $2\pi$ -periodic. Therefore  $S_n f$  is always  $2\pi$ -periodic. Therefore we can only approximate  $2\pi$ -periodic functions. (And by change of variables, other periodic functions.)