



Aalto University

MS-C1350 Partial differential equations

Chapters 2.1–2.2 – Periodic functions, L^p -functions and inner product

Riikka Korte

Department of Mathematics and Systems Analysis
Aalto University
riikka.korte@aalto.fi

September 3, 2024

2 Fourier series (intro)

- ▶ Fourier series is a series representation of a function as trigonometric polynomials.
- ▶ <https://www.geogebra.org/m/fdaYrVCH>

2 Fourier series (intro)

- ▶ Fourier series is a series representation of a function as trigonometric polynomials.
- ▶ <https://www.geogebra.org/m/fdaYrVCH>
- ▶ Euler's formula:

$$e^{ikt} = \cos(kt) + i \sin kt, \quad e^{-ikt} = \cos(kt) - i \sin kt$$

⇒ We can use exponential function instead of trigonometric functions.

2 Fourier series (intro)

- ▶ Fourier series is a series representation of a function as trigonometric polynomials.
- ▶ <https://www.geogebra.org/m/fdaYrVCH>
- ▶ Euler's formula:

$$e^{ikt} = \cos(kt) + i \sin kt, \quad e^{-ikt} = \cos(kt) - i \sin kt$$

⇒ We can use exponential function instead of trigonometric functions.

- ▶ Compare to Taylor series – both express a function as infinite sum of elementary functions

2 Fourier series (intro)

- ▶ Fourier series is a series representation of a function as trigonometric polynomials.
- ▶ <https://www.geogebra.org/m/fdaYrVCH>
- ▶ Euler's formula:

$$e^{ikt} = \cos(kt) + i \sin kt, \quad e^{-ikt} = \cos(kt) - i \sin kt$$

⇒ We can use exponential function instead of trigonometric functions.

- ▶ Compare to Taylor series – both express a function as infinite sum of elementary functions
- ▶ Fourier series gives the best square approximation of the function.

2 Fourier series (intro)

- ▶ Fourier series is a series representation of a function as trigonometric polynomials.
- ▶ <https://www.geogebra.org/m/fdaYrVCH>
- ▶ Euler's formula:

$$e^{ikt} = \cos(kt) + i \sin kt, \quad e^{-ikt} = \cos(kt) - i \sin kt$$

⇒ We can use exponential function instead of trigonometric functions.

- ▶ Compare to Taylor series – both express a function as infinite sum of elementary functions
- ▶ Fourier series gives the best square approximation of the function.
- ▶ Fourier series converges pointwise if the function is smooth enough.

2 Fourier series (intro)

- ▶ Fourier series is a series representation of a function as trigonometric polynomials.
- ▶ <https://www.geogebra.org/m/fdaYrVCH>
- ▶ Euler's formula:

$$e^{ikt} = \cos(kt) + i \sin kt, \quad e^{-ikt} = \cos(kt) - i \sin kt$$

⇒ We can use exponential function instead of trigonometric functions.

- ▶ Compare to Taylor series – both express a function as infinite sum of elementary functions
- ▶ Fourier series gives the best square approximation of the function.
- ▶ Fourier series converges pointwise if the function is smooth enough.
- ▶ Many PDE problems can be solved using using Fourier series and convolutions.

2 Fourier series (intro)

- ▶ Fourier series is a series representation of a function as trigonometric polynomials.
- ▶ <https://www.geogebra.org/m/fdaYrVCH>
- ▶ Euler's formula:

$$e^{ikt} = \cos(kt) + i \sin kt, \quad e^{-ikt} = \cos(kt) - i \sin kt$$

⇒ We can use exponential function instead of trigonometric functions.

- ▶ Compare to Taylor series – both express a function as infinite sum of elementary functions
- ▶ Fourier series gives the best square approximation of the function.
- ▶ Fourier series converges pointwise if the function is smooth enough.
- ▶ Many PDE problems can be solved using using Fourier series and convolutions.
- ▶ We need following concepts to be able to do Fourier theory:
 - ▶ periodic functions
 - ▶ L^p -functions, in particular L^2 -functions
 - ▶ Innerproduct spaces

Periodic functions (Chapter 2.1)

- We say that a function $f : \mathbb{R} \rightarrow \mathbb{C}$ is 2π -periodic if for every $t \in \mathbb{R}$ we have

$$(1) \qquad f(t + 2\pi) = f(t).$$

Periodic functions (Chapter 2.1)

- ▶ We say that a function $f : \mathbb{R} \rightarrow \mathbb{C}$ is 2π -periodic if for every $t \in \mathbb{R}$ we have

$$(1) \qquad f(t + 2\pi) = f(t).$$

- ▶ More generally, a function $f : \mathbb{R} \rightarrow \mathbb{C}$ is called **T -periodic**, $T \in \mathbb{R}$, $T \neq 0$, if

$$(2) \qquad f(t + T) = f(t)$$

for every $t \in \mathbb{R}$.

Periodic functions (Chapter 2.1)

- ▶ We say that a function $f : \mathbb{R} \rightarrow \mathbb{C}$ is 2π -periodic if for every $t \in \mathbb{R}$ we have

$$(1) \qquad f(t + 2\pi) = f(t).$$

- ▶ More generally, a function $f : \mathbb{R} \rightarrow \mathbb{C}$ is called T -periodic , $T \in \mathbb{R}, T \neq 0$, if

$$(2) \qquad f(t + T) = f(t)$$

for every $t \in \mathbb{R}$.

- ▶ Every function $f : [a, b] \rightarrow \mathbb{C}, a, b \in \mathbb{R}$, can be extended to a periodic function on the whole \mathbb{R} .

Periodic functions (Chapter 2.1)

- ▶ We say that a function $f : \mathbb{R} \rightarrow \mathbb{C}$ is 2π -periodic if for every $t \in \mathbb{R}$ we have

$$(1) \qquad f(t + 2\pi) = f(t).$$

- ▶ More generally, a function $f : \mathbb{R} \rightarrow \mathbb{C}$ is called T -periodic , $T \in \mathbb{R}, T \neq 0$, if

$$(2) \qquad f(t + T) = f(t)$$

for every $t \in \mathbb{R}$.

- ▶ Every function $f : [a, b] \rightarrow \mathbb{C}, a, b \in \mathbb{R}$, can be extended to a periodic function on the whole \mathbb{R} .
- ▶ Fourier series are defined only for periodic functions, but this is not a serious restriction, because we can extend the functions periodically.

L^p space (Chapter 2.2)

Solutions to PDEs are not always smooth. Therefore we use L^p spaces ($p \geq 1$). Especially L^1 and L^2 are important for Fourier series.

L^p space (Chapter 2.2)

Solutions to PDEs are not always smooth. Therefore we use L^p spaces ($p \geq 1$). Especially L^1 and L^2 are important for Fourier series.

Definition

Let $1 \leq p < \infty$. A function $f : [-\pi, \pi] \rightarrow \mathbb{C}$ belongs to $L^p([-\pi, \pi])$, if

$$\|f\|_{L^p([-\pi, \pi])} = \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |f(t)|^p dt \right)^{\frac{1}{p}} < \infty.$$

The number $\|f\|_{L^p([-\pi, \pi])}$ is called the L^p -norm of f .

L^p space (Chapter 2.2)

Solutions to PDEs are not always smooth. Therefore we use L^p spaces ($p \geq 1$). Especially L^1 and L^2 are important for Fourier series.

Definition

Let $1 \leq p < \infty$. A function $f : [-\pi, \pi] \rightarrow \mathbb{C}$ belongs to $L^p([-\pi, \pi])$, if

$$\|f\|_{L^p([-\pi, \pi])} = \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |f(t)|^p dt \right)^{\frac{1}{p}} < \infty.$$

The number $\|f\|_{L^p([-\pi, \pi])}$ is called the L^p -norm of f .

- L^p functions need not be smooth or even continuous.

L^p space (Chapter 2.2)

Solutions to PDEs are not always smooth. Therefore we use L^p spaces ($p \geq 1$). Especially L^1 and L^2 are important for Fourier series.

Definition

Let $1 \leq p < \infty$. A function $f : [-\pi, \pi] \rightarrow \mathbb{C}$ belongs to $L^p([-\pi, \pi])$, if

$$\|f\|_{L^p([-\pi, \pi])} = \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |f(t)|^p dt \right)^{\frac{1}{p}} < \infty.$$

The number $\|f\|_{L^p([-\pi, \pi])}$ is called the L^p -norm of f .

- ▶ L^p functions need not be smooth or even continuous.
- ▶ $C([-\pi, \pi]) \subset L^2([-\pi, \pi])$ **BUT** $C((-\pi, \pi)) \not\subset L^2([-\pi, \pi])$

L^p space (Chapter 2.2)

Solutions to PDEs are not always smooth. Therefore we use L^p spaces ($p \geq 1$). Especially L^1 and L^2 are important for Fourier series.

Definition

Let $1 \leq p < \infty$. A function $f : [-\pi, \pi] \rightarrow \mathbb{C}$ belongs to $L^p([-\pi, \pi])$, if

$$\|f\|_{L^p([-\pi, \pi])} = \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |f(t)|^p dt \right)^{\frac{1}{p}} < \infty.$$

The number $\|f\|_{L^p([-\pi, \pi])}$ is called the L^p -norm of f .

- ▶ L^p functions need not be smooth or even continuous.
- ▶ $C([-\pi, \pi]) \subset L^2([-\pi, \pi])$ BUT $C((-\pi, \pi)) \not\subset L^2([-\pi, \pi])$
- ▶ L^p functions can have singularities.

(Complex) Inner product

- ▶ Inner product space is a vector space V with a map:
 $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{C}$ that satisfies ($\forall x, y, z \in V, a \in \mathbb{C}$)

- ▶ Linearity

$$\langle ax, y \rangle = a \langle x, y \rangle, \quad \langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle,$$

- ▶ conjugate symmetry

$$\langle x, y \rangle = \overline{\langle y, x \rangle},$$

- ▶ positivedefinitess

$$\langle x, x \rangle > 0 \text{ if } x \neq 0.$$

(Complex) Inner product

- ▶ Inner product space is a vector space V with a map:
 $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{C}$ that satisfies ($\forall x, y, z \in V, a \in \mathbb{C}$)

- ▶ Linearity

$$\langle ax, y \rangle = a \langle x, y \rangle, \quad \langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle,$$

- ▶ conjugate symmetry

$$\langle x, y \rangle = \overline{\langle y, x \rangle},$$

- ▶ positivedefinitess

$$\langle x, x \rangle > 0 \text{ if } x \neq 0.$$

- ▶ There are both finite dimensional vector spaces (e.g $\mathbb{R}^n, \mathbb{C}^n$) and infinite dimensional vector spaces (e.g function spaces).

(Complex) Inner product

- ▶ Inner product space is a vector space V with a map:
 $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{C}$ that satisfies ($\forall x, y, z \in V, a \in \mathbb{C}$)

- ▶ Linearity

$$\langle ax, y \rangle = a \langle x, y \rangle, \quad \langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle,$$

- ▶ conjugate symmetry

$$\langle x, y \rangle = \overline{\langle y, x \rangle},$$

- ▶ positivedefinitess

$$\langle x, x \rangle > 0 \text{ if } x \neq 0.$$

- ▶ There are both finite dimensional vector spaces (e.g $\mathbb{R}^n, \mathbb{C}^n$) and infinite dimensional vector spaces (e.g function spaces).
- ▶ If $\{e_i\}_{i=-\infty}^{\infty}$ is an orthonormalisation basis for V then

$$f = \sum_{i=-\infty}^{\infty} \langle f, e_i \rangle e_i \quad \text{for every } f \in V.$$

Complex vector space $L^2([-\pi, \pi])$

► Addition and multiplication

$$(f + g)(t) = f(t) + g(t), \quad (\alpha f)(t) = \alpha f(t), \quad \alpha \in \mathbb{C}.$$

Complex vector space $L^2([-\pi, \pi])$

► Addition and multiplication

$$(f + g)(t) = f(t) + g(t), \quad (\alpha f)(t) = \alpha f(t), \quad \alpha \in \mathbb{C}.$$

► Inner product:

$$\langle f, g \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) \overline{g(t)} dt$$

Here $\bar{z} = x - iy \in \mathbb{C}$ is the complex conjugate of $z = x + iy$, where $x, y \in \mathbb{R}$ and i is the imaginary unit.

Complex vector space $L^2([-\pi, \pi])$

- Addition and multiplication

$$(f + g)(t) = f(t) + g(t), \quad (\alpha f)(t) = \alpha f(t), \quad \alpha \in \mathbb{C}.$$

- Inner product:

$$\langle f, g \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) \overline{g(t)} dt$$

Here $\bar{z} = x - iy \in \mathbb{C}$ is the complex conjugate of $z = x + iy$, where $x, y \in \mathbb{R}$ and i is the imaginary unit.

- Norm given by the inner product:

$$\|f\|_{L^2([-\pi, \pi])} = \langle f, f \rangle^{1/2}$$

Important example of an inner product

► Let $e_j : [-\pi, \pi] \rightarrow \mathbb{C}$,

$$e_j(t) = e^{ij t} = \cos(jt) + i \sin(jt), \quad j \in \mathbb{Z} \quad (\text{Euler's formula}).$$

Important example of an inner product

- ▶ Let $e_j : [-\pi, \pi] \rightarrow \mathbb{C}$,

$$e_j(t) = e^{ij t} = \cos(jt) + i \sin(jt), \quad j \in \mathbb{Z} \quad (\text{Euler's formula}).$$

- ▶ Then $e_j \in C([-\pi, \pi]) \subset L^2([-\pi, \pi])$ with

$$\|e_j\|_{L^2([-\pi, \pi])} = \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} \underbrace{|e^{ij t}|^2}_{=1} dt \right)^{\frac{1}{2}} = 1, \quad j = 1, 2, \dots$$

Important example of an inner product

- ▶ Let $e_j : [-\pi, \pi] \rightarrow \mathbb{C}$,

$$e_j(t) = e^{ijt} = \cos(jt) + i \sin(jt), \quad j \in \mathbb{Z} \quad (\text{Euler's formula}).$$

- ▶ Then $e_j \in C([-\pi, \pi]) \subset L^2([-\pi, \pi])$ with

$$\|e_j\|_{L^2([-\pi, \pi])} = \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} \underbrace{|e^{ijt}|^2}_{=1} dt \right)^{\frac{1}{2}} = 1, \quad j = 1, 2, \dots$$

- ▶ The inner product of two such functions is

$$\langle e_j, e_k \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ijt} \overline{e^{ikt}} dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(j-k)t} dt = 0, \quad \text{if } j \neq k$$

Important example of an inner product

- ▶ Let $e_j : [-\pi, \pi] \rightarrow \mathbb{C}$,

$$e_j(t) = e^{ijt} = \cos(jt) + i \sin(jt), \quad j \in \mathbb{Z} \quad (\text{Euler's formula}).$$

- ▶ Then $e_j \in C([-\pi, \pi]) \subset L^2([-\pi, \pi])$ with

$$\|e_j\|_{L^2([-\pi, \pi])} = \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} \underbrace{|e^{ijt}|^2}_{=1} dt \right)^{\frac{1}{2}} = 1, \quad j = 1, 2, \dots$$

- ▶ The inner product of two such functions is

$$\langle e_j, e_k \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ijt} \overline{e^{ikt}} dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(j-k)t} dt = 0, \quad \text{if } j \neq k$$

- ▶ Thus the set $\{e_j\}_{j \in \mathbb{Z}}$ is an orthonormal set in $L^2([-\pi, \pi])$

$$\langle e_j, e_k \rangle = \delta_{jk} = \begin{cases} 0, & j \neq k, \\ 1, & j = k. \end{cases}$$

Important example of an inner product

- ▶ Let $e_j : [-\pi, \pi] \rightarrow \mathbb{C}$,

$$e_j(t) = e^{ijt} = \cos(jt) + i \sin(jt), \quad j \in \mathbb{Z} \quad (\text{Euler's formula}).$$

- ▶ Then $e_j \in C([-\pi, \pi]) \subset L^2([-\pi, \pi])$ with

$$\|e_j\|_{L^2([-\pi, \pi])} = \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} \underbrace{|e^{ijt}|^2}_{=1} dt \right)^{\frac{1}{2}} = 1, \quad j = 1, 2, \dots$$

- ▶ The inner product of two such functions is

$$\langle e_j, e_k \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ijt} \overline{e^{ikt}} dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(j-k)t} dt = 0, \quad \text{if } j \neq k$$

- ▶ Thus the set $\{e_j\}_{j \in \mathbb{Z}}$ is an orthonormal set in $L^2([-\pi, \pi])$

$$\langle e_j, e_k \rangle = \delta_{jk} = \begin{cases} 0, & j \neq k, \\ 1, & j = k. \end{cases}$$

- ▶ It is actually also an orthonormal basis (which is not so easy to prove).