



Aalto University

# MS-C1350 Partial differential equations

## Ch 2.5-2.10 Fourier series

Riikka Korte

Department of Mathematics and Systems Analysis  
Aalto University  
[riikka.korte@aalto.fi](mailto:riikka.korte@aalto.fi)

September 10, 2024

## 2.5 Other intervals

The  $n$ th partial sum of a Fourier series of  $f$  on  $[a, b]$  is

$$S_n f(t) = \sum_{j=-n}^n \langle f, e_j \rangle e_j = \sum_{j=-n}^n \hat{f}(j) e^{\frac{2\pi i j t}{b-a}}, \quad n = 0, 1, 2, \dots,$$

where the Fourier coefficients are

$$\hat{f}(j) = \frac{1}{b-a} \int_a^b f(t) e^{\frac{-2\pi i j t}{b-a}} dt, \quad j \in \mathbb{Z}.$$

This follows from a change of variables.

Thus everything works in the same way as with the interval  $[-\pi, \pi]$ , but the formulas get more messy when the length of the interval is not the same as the period of trigonometric functions.

## 2.6 Real form of Fourier series

Let  $f \in L^1([-\pi, \pi])$ . The  $n$ th partial sum of a Fourier series can be written as

$$S_n f(t) = \frac{a_0}{2} + \sum_{j=1}^n (a_j \cos(jt) + b_j \sin(jt)),$$

where

$$a_j = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) \cos(jt) dt, \quad j = 0, 1, 2, \dots$$

and

$$b_j = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) \sin(jt) dt, \quad j = 1, 2, \dots$$

This is called the real form of the Fourier series of  $f$ . The coefficients  $a_j$  are called the Fourier cosine coefficients of  $f$  and  $b_j$  are called the Fourier sine coefficients of  $f$ . The corresponding series are called the Fourier cosine and sine series of  $f$  correspondingly.

## 2.6 Real form of Fourier series

- Recall that

$$e^{ijt} = \cos(jt) + i \sin(jt), \quad e^{-ijt} = \cos(jt) - i \sin(jt).$$

- If  $f$  is real, then also  $a_j, b_j$  are real whereas  $\hat{f}(j)$ 's are not usually all real.

## 2.6 Real form of Fourier series

- Recall that

$$e^{ijt} = \cos(jt) + i \sin(jt), \quad e^{-ijt} = \cos(jt) - i \sin(jt).$$

- If  $f$  is real, then also  $a_j, b_j$  are real whereas  $\hat{f}(j)$ 's are not usually all real.
- Even function:  $f(-t) = f(t)$ . Odd function:  $f(-t) = -f(t)$ .

## 2.6 Real form of Fourier series

- Recall that

$$e^{ijt} = \cos(jt) + i \sin(jt), \quad e^{-ijt} = \cos(jt) - i \sin(jt).$$

- If  $f$  is real, then also  $a_j, b_j$  are real whereas  $\hat{f}(j)$ 's are not usually all real.
- Even function:  $f(-t) = f(t)$ . Odd function:  $f(-t) = -f(t)$ .
- If  $f$  is even, then  $b_j = 0$  for all  $j$ .
- If  $f$  is odd, then  $a_j = 0$  for all  $j$ .

## 2.6 Real form of Fourier series

- Recall that

$$e^{ijt} = \cos(jt) + i \sin(jt), \quad e^{-ijt} = \cos(jt) - i \sin(jt).$$

- If  $f$  is real, then also  $a_j, b_j$  are real whereas  $\hat{f}(j)$ 's are not usually all real.
- Even function:  $f(-t) = f(t)$ . Odd function:  $f(-t) = -f(t)$ .
- If  $f$  is even, then  $b_j = 0$  for all  $j$ .
- If  $f$  is odd, then  $a_j = 0$  for all  $j$ .
- Let  $f : [0, L] \rightarrow \mathbb{R}$ .

## 2.6 Real form of Fourier series

- Recall that

$$e^{ijt} = \cos(jt) + i \sin(jt), \quad e^{-ijt} = \cos(jt) - i \sin(jt).$$

- If  $f$  is real, then also  $a_j, b_j$  are real whereas  $\hat{f}(j)$ 's are not usually all real.
- Even function:  $f(-t) = f(t)$ . Odd function:  $f(-t) = -f(t)$ .
- If  $f$  is even, then  $b_j = 0$  for all  $j$ .
- If  $f$  is odd, then  $a_j = 0$  for all  $j$ .
- Let  $f : [0, L] \rightarrow \mathbb{R}$ .
- Even  $2L$ -periodic extension:  $f(-t) = f(t)$ . Then the series contains only cos-terms.



## 2.6 Real form of Fourier series

- Recall that

$$e^{ijt} = \cos(jt) + i \sin(jt), \quad e^{-ijt} = \cos(jt) - i \sin(jt).$$

- If  $f$  is real, then also  $a_j, b_j$  are real whereas  $\hat{f}(j)$ 's are not usually all real.
- Even function:  $f(-t) = f(t)$ . Odd function:  $f(-t) = -f(t)$ .
- If  $f$  is even, then  $b_j = 0$  for all  $j$ .
- If  $f$  is odd, then  $a_j = 0$  for all  $j$ .
- Let  $f : [0, L] \rightarrow \mathbb{R}$ .
- Even  $2L$ -periodic extension:  $f(-t) = f(t)$ . Then the series contains only  $\cos$ -terms.
- Odd  $2L$ -periodic extension:  $f(-t) = -f(t)$ . Then the series contains only  $\sin$ -terms.

## 2.6 Real form of Fourier series

- Recall that

$$e^{ijt} = \cos(jt) + i \sin(jt), \quad e^{-ijt} = \cos(jt) - i \sin(jt).$$

- If  $f$  is real, then also  $a_j, b_j$  are real whereas  $\hat{f}(j)$ 's are not usually all real.
- Even function:  $f(-t) = f(t)$ . Odd function:  $f(-t) = -f(t)$ .
- If  $f$  is even, then  $b_j = 0$  for all  $j$ .
- If  $f$  is odd, then  $a_j = 0$  for all  $j$ .
- Let  $f : [0, L] \rightarrow \mathbb{R}$ .
- Even  $2L$ -periodic extension:  $f(-t) = f(t)$ . Then the series contains only  $\cos$ -terms.
- Odd  $2L$ -periodic extension:  $f(-t) = -f(t)$ . Then the series contains only  $\sin$ -terms.
- This does not violate the uniqueness of Fourier series as the functions are different on  $[-L, 0]$ .

## 2.7 Differentiation and Fourier series

- ▶ The smoother the function, the faster the Fourier coefficients decay.

## 2.7 Differentiation and Fourier series

- ▶ The smoother the function, the faster the Fourier coefficients decay.
- ▶ Fourier transform of a derivative: Let  $f \in C^1(\mathbb{R})$  be  $2\pi$ -periodic. Then

$$\widehat{f'}(j) = ij\widehat{f}(j), \quad j \in \mathbb{Z}.$$

## 2.7 Differentiation and Fourier series

- ▶ The smoother the function, the faster the Fourier coefficients decay.
- ▶ Fourier transform of a derivative: Let  $f \in C^1(\mathbb{R})$  be  $2\pi$ -periodic. Then

$$\widehat{f'}(j) = ij\widehat{f}(j), \quad j \in \mathbb{Z}.$$

- ▶ Reason: Fourier series can be differentiated termwise:

$$f(t) = \sum_{j=-\infty}^{\infty} \widehat{f}(j)e^{ijt} \implies f'(t) = \sum_{j=-\infty}^{\infty} ij\widehat{f}(j)e^{ijt}.$$

## 2.8 Dirichlet kernel

- ▶ Dirichlet kernel is a useful way of expressing  $S_n f$ .

$$S_n f(t) = \sum_{j=-n}^n \hat{f}(j) e^{ijt}$$

## 2.8 Dirichlet kernel

- ▶ Dirichlet kernel is a useful way of expressing  $S_n f$ .

$$S_n f(t) = \sum_{j=-n}^n \hat{f}(j) e^{ijt} = \sum_{j=-n}^n \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) e^{-ijs} ds \right) e^{ijt}$$

## 2.8 Dirichlet kernel

- ▶ Dirichlet kernel is a useful way of expressing  $S_n f$ .

$$\begin{aligned} S_n f(t) &= \sum_{j=-n}^n \widehat{f}(j) e^{ijt} = \sum_{j=-n}^n \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) e^{-ijs} ds \right) e^{ijt} \\ &= \frac{1}{2\pi} \sum_{j=-n}^n \int_{-\pi}^{\pi} f(s) e^{ij(t-s)} ds \end{aligned}$$



## 2.8 Dirichlet kernel

- ▶ Dirichlet kernel is a useful way of expressing  $S_n f$ .

$$\begin{aligned} S_n f(t) &= \sum_{j=-n}^n \widehat{f}(j) e^{ijt} = \sum_{j=-n}^n \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) e^{-ijs} ds \right) e^{ijt} \\ &= \frac{1}{2\pi} \sum_{j=-n}^n \int_{-\pi}^{\pi} f(s) e^{ij(t-s)} ds = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) \left( \sum_{j=-n}^n e^{ij(t-s)} \right) \end{aligned}$$

## 2.8 Dirichlet kernel

- ▶ Dirichlet kernel is a useful way of expressing  $S_n f$ .

$$\begin{aligned} S_n f(t) &= \sum_{j=-n}^n \widehat{f}(j) e^{ijt} = \sum_{j=-n}^n \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) e^{-ijs} ds \right) e^{ijt} \\ &= \frac{1}{2\pi} \sum_{j=-n}^n \int_{-\pi}^{\pi} f(s) e^{ij(t-s)} ds = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) \left( \sum_{j=-n}^n e^{ij(t-s)} \right) \end{aligned}$$

- ▶ Now we define the Dirichlet kernel as

$$D_n(t) = \sum_{j=-n}^n e^{ijt}, \quad n = 0, 1, 2, \dots$$

## 2.8 Dirichlet kernel

- Dirichlet kernel is a useful way of expressing  $S_n f$ .

$$\begin{aligned} S_n f(t) &= \sum_{j=-n}^n \widehat{f}(j) e^{ijt} = \sum_{j=-n}^n \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) e^{-ijs} ds \right) e^{ijt} \\ &= \frac{1}{2\pi} \sum_{j=-n}^n \int_{-\pi}^{\pi} f(s) e^{ij(t-s)} ds = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) \left( \sum_{j=-n}^n e^{ij(t-s)} \right) \end{aligned}$$

- Now we define the Dirichlet kernel as

$$D_n(t) = \sum_{j=-n}^n e^{ijt}, \quad n = 0, 1, 2, \dots$$

- With this definition we have the formula

$$S_n f(t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) D_n(t-s) ds, \quad t \in [-\pi, \pi], \quad n = 0, 1, 2, \dots$$

# Properties of the Dirichlet kernel



$$D_n f(t) = \sum_{j=-n}^n e^{ijt}$$

# Properties of the Dirichlet kernel



$$D_n f(t) = \sum_{j=-n}^n e^{ijt} = \frac{\sin\left(\left(n + \frac{1}{2}\right)t\right)}{\sin\left(\frac{1}{2}t\right)}, \quad t \neq 0, \quad n = 0, 1, 2, \dots$$

- ▶ The Dirichlet kernel is a  $2\pi$ -periodic function.

# Properties of the Dirichlet kernel



$$D_n f(t) = \sum_{j=-n}^n e^{ijt} = \frac{\sin\left(\left(n + \frac{1}{2}\right)t\right)}{\sin\left(\frac{1}{2}t\right)}, \quad t \neq 0, \quad n = 0, 1, 2, \dots$$

- ▶ The Dirichlet kernel is a  $2\pi$ -periodic function.
- ▶  $D_n(0) = 2n + 1$  and  $D_n(\pi) = (-1)^n$ ,  $n = 0, 1, 2, \dots$

# Properties of the Dirichlet kernel



$$D_n f(t) = \sum_{j=-n}^n e^{ijt} = \frac{\sin\left(\left(n + \frac{1}{2}\right)t\right)}{\sin\left(\frac{1}{2}t\right)}, \quad t \neq 0, \quad n = 0, 1, 2, \dots$$

- ▶ The Dirichlet kernel is a  $2\pi$ -periodic function.
- ▶  $D_n(0) = 2n + 1$  and  $D_n(\pi) = (-1)^n$ ,  $n = 0, 1, 2, \dots$
- ▶ The Dirichlet formula can be written as a convolution

$$S_n f(t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) D_n(t-s) ds = (D_n * f)(t), \quad t \in [-\pi, \pi].$$

# Convolution

- Convolution is an important notation in Fourier analysis and PDEs.



# Convolution

- ▶ Convolution is an important notation in Fourier analysis and PDEs.
- ▶ Let  $f, g \in C(\mathbb{R})$  be  $2\pi$ -periodic.  
The convolution of  $f$  and  $g$  on  $[-\pi, \pi]$  is

$$(f * g)(t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t-s)g(s) ds = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s)g(t-s) ds.$$

# Convolution

- ▶ Convolution is an important notation in Fourier analysis and PDEs.
- ▶ Let  $f, g \in C(\mathbb{R})$  be  $2\pi$ -periodic.  
The convolution of  $f$  and  $g$  on  $[-\pi, \pi]$  is

$$(f * g)(t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t-s)g(s) ds = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s)g(t-s) ds.$$

- ▶ If  $g = 1$  then  $f * g$  is constant.  
Moral: Convolutions can be considered as weighted averages of the functions.

# Convolution

- ▶ Convolution is an important notation in Fourier analysis and PDEs.
- ▶ Let  $f, g \in C(\mathbb{R})$  be  $2\pi$ -periodic.  
The convolution of  $f$  and  $g$  on  $[-\pi, \pi]$  is

$$(f * g)(t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t-s)g(s) ds = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s)g(t-s) ds.$$

- ▶ If  $g = 1$  then  $f * g$  is constant.  
Moral: Convolutions can be considered as weighted averages of the functions.
- ▶  $f * g$  is also  $2\pi$ -periodic.

# Convolution

- ▶ Convolution is an important notation in Fourier analysis and PDEs.
- ▶ Let  $f, g \in C(\mathbb{R})$  be  $2\pi$ -periodic.  
The convolution of  $f$  and  $g$  on  $[-\pi, \pi]$  is

$$(f * g)(t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t-s)g(s) ds = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s)g(t-s) ds.$$

- ▶ If  $g = 1$  then  $f * g$  is constant.  
Moral: Convolutions can be considered as weighted averages of the functions.
- ▶  $f * g$  is also  $2\pi$ -periodic.
- ▶  $f * g = g * f$  and  $f * (g * h) = (f * g) * h$ .

# Convolution

- ▶ Convolution is an important notation in Fourier analysis and PDEs.
- ▶ Let  $f, g \in C(\mathbb{R})$  be  $2\pi$ -periodic.  
The convolution of  $f$  and  $g$  on  $[-\pi, \pi]$  is

$$(f * g)(t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t-s)g(s) ds = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s)g(t-s) ds.$$

- ▶ If  $g = 1$  then  $f * g$  is constant.  
Moral: Convolutions can be considered as weighted averages of the functions.
- ▶  $f * g$  is also  $2\pi$ -periodic.
- ▶  $f * g = g * f$  and  $f * (g * h) = (f * g) * h$ .
- ▶  $\widehat{(f * g)}(j) = \widehat{f}(j)\widehat{g}(j), j \in \mathbb{Z}$ .

# Pointwise convergence

## Theorem

*Let  $f \in C([-\pi, \pi])$  be a  $2\pi$ -periodic function which is differentiable at some point  $t_0 \in [-\pi, \pi]$ . Then*

$$\lim_{n \rightarrow \infty} S_n f(t_0) = f(t_0).$$

# Pointwise convergence

## Theorem

*Let  $f \in C([-\pi, \pi])$  be a  $2\pi$ -periodic function which is differentiable at some point  $t_0 \in [-\pi, \pi]$ . Then*

$$\lim_{n \rightarrow \infty} S_n f(t_0) = f(t_0).$$

- ▶ This shows that the convergence of the Fourier series depends only on the behaviour of the function in an arbitrarily small neighbourhood of the point.

# Pointwise convergence

## Theorem

*Let  $f \in C([-\pi, \pi])$  be a  $2\pi$ -periodic function which is differentiable at some point  $t_0 \in [-\pi, \pi]$ . Then*

$$\lim_{n \rightarrow \infty} S_n f(t_0) = f(t_0).$$

- ▶ This shows that the convergence of the Fourier series depends only on the behaviour of the function in an arbitrarily small neighbourhood of the point.
- ▶ There exists a function  $f \in L^1([-\pi, \pi])$  whose Fourier series diverges at every point.



# Pointwise convergence

## Theorem

*Let  $f \in C([-\pi, \pi])$  be a  $2\pi$ -periodic function which is differentiable at some point  $t_0 \in [-\pi, \pi]$ . Then*

$$\lim_{n \rightarrow \infty} S_n f(t_0) = f(t_0).$$

- ▶ This shows that the convergence of the Fourier series depends only on the behaviour of the function in an arbitrarily small neighbourhood of the point.
- ▶ There exists a function  $f \in L^1([-\pi, \pi])$  whose Fourier series diverges at every point.
- ▶ There are also continuous functions whose Fourier series diverge in a dense set.