



Aalto University

# **MS-C1350 Partial differential equations**

## **Chapter 2.11**

### **Laplace equation in the unit disc**

Riikka Korte

Department of Mathematics and Systems Analysis  
Aalto University  
[riikka.korte@aalto.fi](mailto:riikka.korte@aalto.fi)

September 17, 2024

# Lecture 3

Today we see how to solve Dirichlet problem in the unit disc (Chapter 2.11). Needed techniques include:

- ▶ Moving to polar coordinates
- ▶ Separation of variables
- ▶ Solving equations we obtain by separation of variables.
- ▶ Using Fourier series to find the correct Fourier series solution.

In Chapter 2.12, the heat equation in 1D is solved in a similar way. That example is not discussed in the lecture (but there is an old lecture video about it).

# Laplace equation in the unit disc

- ▶ Laplace equation:

$$\Delta u = 0$$

- ▶ Laplace equation models heat distribution when the system has reached thermal equilibrium.
- ▶ Appears in many other places as well.

# Dirichlet problem

- Consider 2-dimensional unit disc in  $\mathbb{R}^2$ :

$$\Omega = \{(x, y) \in \mathbb{R}^2 : (x^2 + y^2)^{\frac{1}{2}} < 1\}$$

# Dirichlet problem

- ▶ Consider 2-dimensional unit disc in  $\mathbb{R}^2$ :

$$\Omega = \{(x, y) \in \mathbb{R}^2 : (x^2 + y^2)^{\frac{1}{2}} < 1\}$$

- ▶ Let  $g \in C(\partial\Omega)$  be a continuous function on the boundary

$$\partial\Omega = \{(x, y) \in \mathbb{R}^2 : (x^2 + y^2)^{\frac{1}{2}} = 1\}.$$

# Dirichlet problem

- Consider 2-dimensional unit disc in  $\mathbb{R}^2$ :

$$\Omega = \{(x, y) \in \mathbb{R}^2 : (x^2 + y^2)^{\frac{1}{2}} < 1\}$$

- Let  $g \in C(\partial\Omega)$  be a continuous function on the boundary

$$\partial\Omega = \{(x, y) \in \mathbb{R}^2 : (x^2 + y^2)^{\frac{1}{2}} = 1\}.$$

- The problem is to find  $u \in C^2(\Omega) \cap C(\overline{\Omega})$  such that

$$\begin{cases} \Delta u(x, y) = \frac{\partial^2 u}{\partial x^2}(x, y) + \frac{\partial^2 u}{\partial y^2}(x, y) = 0, & (x, y) \in \Omega, \\ u(x, y) = g(x, y), & (x, y) \in \partial\Omega. \end{cases}$$

# Solving the Dirichlet problem

- ▶ We will solve this problem using Fourier series together with technique called separation of variables.

# Solving the Dirichlet problem

- ▶ We will solve this problem using Fourier series together with technique called separation of variables.
- ▶ This means that we (first forget about the boundary values and) look for solutions of the form

$$u(x, y) = A(x)B(y),$$



# Solving the Dirichlet problem

- ▶ We will solve this problem using Fourier series together with technique called separation of variables.
- ▶ This means that we (first forget about the boundary values and) look for solutions of the form

$$u(x, y) = A(x)B(y),$$

- ▶ This would work well in rectangular areas, but for the disc, we switch to polar coordinates:

$$(x, y) = (r \cos \theta, r \sin \theta), \quad (x, y) \in \mathbb{R}^2, \quad 0 \leq r < \infty, \quad -\pi \leq \theta < \pi,$$

where  $r^2 = x^2 + y^2$  and  $\tan \theta = \frac{y}{x}$ . In polar coordinates, we have

$$\Omega = \{(r, \theta) : 0 \leq r < 1, -\pi \leq \theta < \pi\}$$

and

$$\partial\Omega = \{(1, \theta) : -\pi \leq \theta < \pi\}.$$

# Solving the Dirichlet problem

- ▶ We will solve this problem using Fourier series together with technique called separation of variables.
- ▶ This means that we (first forget about the boundary values and) look for solutions of the form

$$u(x, y) = A(x)B(y),$$

- ▶ This would work well in rectangular areas, but for the disc, we switch to polar coordinates:

$$(x, y) = (r \cos \theta, r \sin \theta), \quad (x, y) \in \mathbb{R}^2, \quad 0 \leq r < \infty, \quad -\pi \leq \theta < \pi,$$

where  $r^2 = x^2 + y^2$  and  $\tan \theta = \frac{y}{x}$ . In polar coordinates, we have

$$\Omega = \{(r, \theta) : 0 \leq r < 1, -\pi \leq \theta < \pi\}$$

and

$$\partial\Omega = \{(1, \theta) : -\pi \leq \theta < \pi\}.$$

- ▶ Note that unit disc is a rectangular set in polar coordinates and this is compatible with separation of variables.

# Laplace equation in polar coordinates

## Lemma

*The two-dimensional Laplace operator in polar coordinates is*

$$\Delta u = \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2}, \quad 0 < r < \infty, \quad -\pi \leq \theta < \pi.$$

[ You need to apply chain rule to prove this. Details are in lecture notes. ]

# Laplace equation in polar coordinates

## Lemma

*The two-dimensional Laplace operator in polar coordinates is*

$$\Delta u = \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2}, \quad 0 < r < \infty, \quad -\pi \leq \theta < \pi.$$

[ You need to apply chain rule to prove this. Details are in lecture notes. ]

Thus the Dirichlet problem assumes the following form:

$$\begin{cases} \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0, & 0 < r < 1, \quad -\pi \leq \theta < \pi, \\ u(1, \theta) = g(\theta), & -\pi \leq \theta < \pi, \end{cases}$$

for  $u = u(r, \theta)$ .

# Solving the Dirichlet problem: Step 1

- **Separation of variables:** We look for a product solution (to  $\Delta u = 0$ ) of the form

$$u(r, \theta) = A(\theta)B(r)$$

# Solving the Dirichlet problem: Step 1

- **Separation of variables:** We look for a product solution (to  $\Delta u = 0$ ) of the form

$$u(r, \theta) = A(\theta)B(r)$$

- We insert this into the PDE and write it in a form, where one side depends on  $r$  and the other on  $\theta$ :

$$\frac{r^2 B''(r) + r B'(r)}{B(r)} = - \frac{A''(\theta)}{A(\theta)}$$

# Solving the Dirichlet problem: Step 1

- **Separation of variables:** We look for a product solution (to  $\Delta u = 0$ ) of the form

$$u(r, \theta) = A(\theta)B(r)$$

- We insert this into the PDE and write it in a form, where one side depends on  $r$  and the other on  $\theta$ :

$$\frac{r^2 B''(r) + r B'(r)}{B(r)} = -\frac{A''(\theta)}{A(\theta)} = \lambda$$

- This can only happen, if both sides are equal to a constant, say equal to  $\lambda$ .

# Solving the Dirichlet problem: Step 1

- **Separation of variables:** We look for a product solution (to  $\Delta u = 0$ ) of the form

$$u(r, \theta) = A(\theta)B(r)$$

- We insert this into the PDE and write it in a form, where one side depends on  $r$  and the other on  $\theta$ :

$$\frac{r^2 B''(r) + r B'(r)}{B(r)} = -\frac{A''(\theta)}{A(\theta)} = \lambda$$

- This can only happen, if both sides are equal to a constant, say equal to  $\lambda$ .
- The PDE has been reduced to a system of two ODEs:

$$\begin{cases} A''(\theta) + \lambda A(\theta) = 0, \\ r^2 B''(r) + r B'(r) - \lambda B(r) = 0. \end{cases}$$



# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ Take into account the boundary data:

$$u(1, \theta) = A(\theta)B(1) = g(\theta)."$$

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ Take into account the boundary data:

$$u(1, \theta) = A(\theta)B(1) = g(\theta).$$

- ▶  $g$  is defined on the circle and it can be identified with a  $2\pi$ -periodic function. Thus  $A$  must be  $2\pi$ -periodic.

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ Take into account the boundary data:

$$u(1, \theta) = A(\theta)B(1) = g(\theta).$$

- ▶  $g$  is defined on the circle and it can be identified with a  $2\pi$ -periodic function. Thus  $A$  must be  $2\pi$ -periodic.
- ▶ For which  $\lambda$  do we have nontrivial solutions to the ODE:

$$A''(\theta) + \lambda A(\theta) = 0 \iff -A''(\theta) = \lambda A(\theta)?$$

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ Take into account the boundary data:

$$u(1, \theta) = A(\theta)B(1) = g(\theta).$$

- ▶  $g$  is defined on the circle and it can be identified with a  $2\pi$ -periodic function. Thus  $A$  must be  $2\pi$ -periodic.
- ▶ For which  $\lambda$  do we have nontrivial solutions to the ODE:

$$A''(\theta) + \lambda A(\theta) = 0 \iff -A''(\theta) = \lambda A(\theta)?$$

- ▶ We are interested in **eigenvalues  $\lambda$  and eigenfunctions  $A$** .

## Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ Take into account the boundary data:

$$u(1, \theta) = A(\theta)B(1) = g(\theta).$$

- ▶  $g$  is defined on the circle and it can be identified with a  $2\pi$ -periodic function. Thus  $A$  must be  $2\pi$ -periodic.
- ▶ For which  $\lambda$  do we have nontrivial solutions to the ODE:

$$A''(\theta) + \lambda A(\theta) = 0 \iff -A''(\theta) = \lambda A(\theta)?$$

- ▶ We are interested in eigenvalues  $\lambda$  and eigenfunctions  $A$ .
- ▶ There are no complex eigenvalues (proof in lecture notes).

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ Take into account the boundary data:

$$u(1, \theta) = A(\theta)B(1) = g(\theta)."$$

- ▶  $g$  is defined on the circle and it can be identified with a  $2\pi$ -periodic function. Thus  $A$  must be  $2\pi$ -periodic.
- ▶ For which  $\lambda$  do we have nontrivial solutions to the ODE:

$$A''(\theta) + \lambda A(\theta) = 0 \iff -A''(\theta) = \lambda A(\theta)?$$

- ▶ We are interested in eigenvalues  $\lambda$  and eigenfunctions  $A$ .
- ▶ There are no complex eigenvalues (proof in lecture notes).
- ▶ **Case:**  $\lambda < 0$ :

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ Take into account the boundary data:

$$u(1, \theta) = A(\theta)B(1) = g(\theta)."$$

- ▶  $g$  is defined on the circle and it can be identified with a  $2\pi$ -periodic function. Thus  $A$  must be  $2\pi$ -periodic.
- ▶ For which  $\lambda$  do we have nontrivial solutions to the ODE:

$$A''(\theta) + \lambda A(\theta) = 0 \iff -A''(\theta) = \lambda A(\theta)?$$

- ▶ We are interested in eigenvalues  $\lambda$  and eigenfunctions  $A$ .
- ▶ There are no complex eigenvalues (proof in lecture notes).
- ▶ **Case:**  $\lambda = -\mu^2 < 0$ :

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ Take into account the boundary data:

$$u(1, \theta) = A(\theta)B(1) = g(\theta).$$

- ▶  $g$  is defined on the circle and it can be identified with a  $2\pi$ -periodic function. Thus  $A$  must be  $2\pi$ -periodic.
- ▶ For which  $\lambda$  do we have nontrivial solutions to the ODE:

$$A''(\theta) + \lambda A(\theta) = 0 \iff -A''(\theta) = \lambda A(\theta)?$$

- ▶ We are interested in eigenvalues  $\lambda$  and eigenfunctions  $A$ .
- ▶ There are no complex eigenvalues (proof in lecture notes).
- ▶ **Case:**  $\lambda = -\mu^2 < 0$ :

$$A(\theta) = c_1 e^{\mu\theta} + c_2 e^{-\mu\theta}.$$



## Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ Take into account the boundary data:

$$u(1, \theta) = A(\theta)B(1) = g(\theta)."$$

- ▶  $g$  is defined on the circle and it can be identified with a  $2\pi$ -periodic function. Thus  $A$  must be  $2\pi$ -periodic.
- ▶ For which  $\lambda$  do we have nontrivial solutions to the ODE:

$$A''(\theta) + \lambda A(\theta) = 0 \iff -A''(\theta) = \lambda A(\theta)?$$

- ▶ We are interested in eigenvalues  $\lambda$  and eigenfunctions  $A$ .
- ▶ There are no complex eigenvalues (proof in lecture notes).
- ▶ **Case:**  $\lambda = -\mu^2 < 0$ :

$$A(\theta) = c_1 e^{\mu\theta} + c_2 e^{-\mu\theta}.$$

The only periodic solution is  $A = 0$ . Then also  $u = AB = 0$ .

# Solving the Dirichlet problem: Step 2: Solving separated equations

► **Case**  $\lambda = 0$ :

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **Case**  $\lambda = 0$ :
- ▶ The equation reduces to  $A'' = 0$  with the general solution

$$A(\theta) = c_1\theta + c_2.$$

# Solving the Dirichlet problem: Step 2: Solving separated equations

► **Case**  $\lambda = 0$ :

► The equation reduces to  $A'' = 0$  with the general solution

$$A(\theta) = c_1\theta + c_2.$$

► The only periodic solution is  $A(\theta) = c_2$ .

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **Case**  $\lambda = 0$ :

- ▶ The equation reduces to  $A'' = 0$  with the general solution

$$A(\theta) = c_1\theta + c_2.$$

- ▶ The only periodic solution is  $A(\theta) = c_2$ .
- ▶ In this case, the ODE for  $B$  becomes

$$rB'' + B' = 0.$$

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **Case**  $\lambda = 0$ :

- ▶ The equation reduces to  $A'' = 0$  with the general solution

$$A(\theta) = c_1\theta + c_2.$$

- ▶ The only periodic solution is  $A(\theta) = c_2$ .
- ▶ In this case, the ODE for  $B$  becomes

$$rB'' + B' = 0.$$

- ▶ Its general solution is

$$B(r) = d_1 + d_2 \ln r.$$

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **Case**  $\lambda = 0$ :

- ▶ The equation reduces to  $A'' = 0$  with the general solution

$$A(\theta) = c_1\theta + c_2.$$

- ▶ The only periodic solution is  $A(\theta) = c_2$ .
- ▶ In this case, the ODE for  $B$  becomes

$$rB'' + B' = 0.$$

- ▶ Its general solution is

$$B(r) = d_1 + d_2 \ln r.$$

- ▶ Thus we get:

$$u(r, \theta) = A(\theta)B(r) = c_2(d_1 + d_2 \ln r).$$

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **Case**  $\lambda = 0$ :

- ▶ The equation reduces to  $A'' = 0$  with the general solution

$$A(\theta) = c_1\theta + c_2.$$

- ▶ The only periodic solution is  $A(\theta) = c_2$ .
- ▶ In this case, the ODE for  $B$  becomes

$$rB'' + B' = 0.$$

- ▶ Its general solution is

$$B(r) = d_1 + d_2 \ln r.$$

- ▶ Thus we get:

$$u(r, \theta) = A(\theta)B(r) = c_2(d_1 + d_2 \ln r).$$

- ▶ Solution becomes unbounded as  $r \rightarrow 0$ . This is against physical intuition and these solutions are excluded. (Not a solution at 0.)



# Solving the Dirichlet problem: Step 2: Solving separated equations

# Solving the Dirichlet problem: Step 2: Solving separated equations

- The case  $\lambda > 0$ :

# Solving the Dirichlet problem: Step 2: Solving separated equations

- **The case**  $\lambda = \mu^2 > 0$ :  $A'' + \mu^2 A = 0$

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **The case**  $\lambda = \mu^2 > 0$ :  $A'' + \mu^2 A = 0$
- ▶ The general complex solution of this equation is

$$A(\theta) = c_1 e^{i\mu\theta} + c_2 e^{-i\mu\theta}.$$

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **The case**  $\lambda = \mu^2 > 0$ :  $A'' + \mu^2 A = 0$
- ▶ The general complex solution of this equation is

$$A(\theta) = c_1 e^{i\mu\theta} + c_2 e^{-i\mu\theta}.$$

- ▶ This is  $2\pi$ -periodic if  $\mu \in \mathbb{Z}$ . Thus  $\lambda = \mu^2 = j^2$ ,  $j \in \mathbb{Z} \setminus \{0\}$ .

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **The case**  $\lambda = \mu^2 > 0$ :  $A'' + \mu^2 A = 0$
- ▶ The general complex solution of this equation is

$$A(\theta) = c_1 e^{i\mu\theta} + c_2 e^{-i\mu\theta}.$$

- ▶ This is  $2\pi$ -periodic if  $\mu \in \mathbb{Z}$ . Thus  $\lambda = \mu^2 = j^2$ ,  $j \in \mathbb{Z} \setminus \{0\}$ .
- ▶ Now the ODE for  $B$  is Euler equation:

$$r^2 B'' + r B' - j^2 B = 0.$$

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **The case**  $\lambda = \mu^2 > 0$ :  $A'' + \mu^2 A = 0$
- ▶ The general complex solution of this equation is

$$A(\theta) = c_1 e^{i\mu\theta} + c_2 e^{-i\mu\theta}.$$

- ▶ This is  $2\pi$ -periodic if  $\mu \in \mathbb{Z}$ . Thus  $\lambda = \mu^2 = j^2$ ,  $j \in \mathbb{Z} \setminus \{0\}$ .
- ▶ Now the ODE for  $B$  is Euler equation:

$$r^2 B'' + r B' - j^2 B = 0.$$

- ▶ Its general solution is

$$B(r) = d_1 r^j + d_2 r^{-j}$$

# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **The case**  $\lambda = \mu^2 > 0$ :  $A'' + \mu^2 A = 0$
- ▶ The general complex solution of this equation is

$$A(\theta) = c_1 e^{i\mu\theta} + c_2 e^{-i\mu\theta}.$$

- ▶ This is  $2\pi$ -periodic if  $\mu \in \mathbb{Z}$ . Thus  $\lambda = \mu^2 = j^2$ ,  $j \in \mathbb{Z} \setminus \{0\}$ .
- ▶ Now the ODE for  $B$  is Euler equation:

$$r^2 B'' + r B' - j^2 B = 0.$$

- ▶ Its general solution is

$$B(r) = d_1 r^j + d_2 r^{-j}$$

- ▶ The term with  $r^{-j}$ ,  $j > 0$  blows up at 0 and is therefore excluded.



# Solving the Dirichlet problem: Step 2: Solving separated equations

- ▶ **The case**  $\lambda = \mu^2 > 0$ :  $A'' + \mu^2 A = 0$
- ▶ The general complex solution of this equation is

$$A(\theta) = c_1 e^{i\mu\theta} + c_2 e^{-i\mu\theta}.$$

- ▶ This is  $2\pi$ -periodic if  $\mu \in \mathbb{Z}$ . Thus  $\lambda = \mu^2 = j^2$ ,  $j \in \mathbb{Z} \setminus \{0\}$ .
- ▶ Now the ODE for  $B$  is Euler equation:

$$r^2 B'' + r B' - j^2 B = 0.$$

- ▶ Its general solution is

$$B(r) = d_1 r^j + d_2 r^{-j}$$

- ▶ The term with  $r^{-j}$ ,  $j > 0$  blows up at 0 and is therefore excluded.
- ▶ Thus  $\lambda > 0$  gives relevant solutions of form

$$u(r, \theta) = A(\theta)B(r) = r^{|j|} e^{ij\theta}.$$

# Solving the Dirichlet problem: Step 3: Fourier series solution for the whole problem

- Now we have found solutions

$$u_j(r, \theta) = A(\theta)B(r) = r^{|j|} e^{ij\theta}.$$

which solve the problem with boundary values

$$u_j(1, \theta) = e^{ij\theta}, \quad j \in \mathbb{Z}.$$

# Solving the Dirichlet problem: Step 3: Fourier series solution for the whole problem

- Now we have found solutions

$$u_j(r, \theta) = A(\theta)B(r) = r^{|j|}e^{ij\theta}.$$

which solve the problem with boundary values

$$u_j(1, \theta) = e^{ij\theta}, \quad j \in \mathbb{Z}.$$

- Laplace operator is linear. Thus all linear combinations are also solutions:

$$u(r, \theta) = \sum_{j=-\infty}^{\infty} a_j r^{|j|} e^{ij\theta}, \quad 0 \leq r < 1, \quad -\pi < \theta \leq \pi.$$

# Solving the Dirichlet problem: Step 3: Fourier series solution for the whole problem

- Now we have found solutions

$$u_j(r, \theta) = A(\theta)B(r) = r^{|j|}e^{ij\theta}.$$

which solve the problem with boundary values

$$u_j(1, \theta) = e^{ij\theta}, \quad j \in \mathbb{Z}.$$

- Laplace operator is linear. Thus all linear combinations are also solutions:

$$u(r, \theta) = \sum_{j=-\infty}^{\infty} a_j r^{|j|} e^{ij\theta}, \quad 0 \leq r < 1, \quad -\pi < \theta \leq \pi.$$

- This should be compatible with the boundary data when  $r = 1$ :

$$u(1, \theta) = \sum_{j=-\infty}^{\infty} a_j e^{ij\theta} = g(\theta).$$

# Solving the Dirichlet problem: Step 3: Fourier series solution for the whole problem

- How to choose  $a_j$ 's so that:

$$u(1, \theta) = \sum_{j=-\infty}^{\infty} a_j e^{ij\theta} = g(\theta)?$$

# Solving the Dirichlet problem: Step 3: Fourier series solution for the whole problem

- ▶ How to choose  $a_j$ 's so that:

$$u(1, \theta) = \sum_{j=-\infty}^{\infty} a_j e^{ij\theta} = g(\theta)?$$

- ▶ If  $g \in L^2([-\pi, \pi])$  then we know that there is a solution if we interpret the equality in  $L^2$ -sense.

# Solving the Dirichlet problem: Step 3: Fourier series solution for the whole problem

- ▶ How to choose  $a_j$ 's so that:

$$u(1, \theta) = \sum_{j=-\infty}^{\infty} a_j e^{ij\theta} = g(\theta)?$$

- ▶ If  $g \in L^2([-\pi, \pi])$  then we know that there is a solution if we interpret the equality in  $L^2$ -sense.
- ▶ If  $g \in C^1([-\pi, \pi])$ , then the Fourier series converges uniformly and

$$a_j = \hat{g}(j).$$

# Things to think about:

1. Does the series solution we obtained converge?



# Things to think about:

1. Does the series solution we obtained converge?
2. Is the function really a solution to our PDE?

# Things to think about:

1. Does the series solution we obtained converge?
2. Is the function really a solution to our PDE?
3. Does the solution attain the correct boundary values?

# Things to think about:

1. Does the series solution we obtained converge?
2. Is the function really a solution to our PDE?
3. Does the solution attain the correct boundary values?
4. Are there other solutions than the solution we found?

# Things to think about:

1. Does the series solution we obtained converge?
2. Is the function really a solution to our PDE?
3. Does the solution attain the correct boundary values?
4. Are there other solutions than the solution we found?

On formal level (2) and (3) seem to be ok. One needs to be careful when switching the order of the limit and the infinite series. We will return later to this and to the question of uniqueness.

## Step 4: Explicit representation formula

- We can plug in the formula for the Fourier coefficients  $\hat{g}(j)$ .

$$u(r, \theta) = \sum_{j=-\infty}^{\infty} \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} g(t) e^{-ijt} dt \right) r^{|j|} e^{ij\theta}$$

## Step 4: Explicit representation formula

- ▶ We can plug in the formula for the Fourier coefficients  $\hat{g}(j)$ .
- ▶ One has to be careful when changing the order of summation and integration.

$$u(r, \theta) = \sum_{j=-\infty}^{\infty} \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} g(t) e^{-ijt} dt \right) r^{|j|} e^{ij\theta}$$

## Step 4: Explicit representation formula

- ▶ We can plug in the formula for the Fourier coefficients  $\hat{g}(j)$ .
- ▶ One has to be careful when changing the order of summation and integration.

$$\begin{aligned} u(r, \theta) &= \sum_{j=-\infty}^{\infty} \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} g(t) e^{-ijt} dt \right) r^{|j|} e^{ij\theta} \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} g(t) \left( \sum_{j=-\infty}^{\infty} r^{|j|} e^{ij(\theta-t)} \right) dt \end{aligned}$$

## Step 4: Explicit representation formula

- ▶ We can plug in the formula for the Fourier coefficients  $\hat{g}(j)$ .
- ▶ One has to be careful when changing the order of summation and integration.

$$\begin{aligned}u(r, \theta) &= \sum_{j=-\infty}^{\infty} \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} g(t) e^{-ijt} dt \right) r^{|j|} e^{ij\theta} \\&= \frac{1}{2\pi} \int_{-\pi}^{\pi} g(t) \left( \sum_{j=-\infty}^{\infty} r^{|j|} e^{ij(\theta-t)} \right) dt \\&= (g * P_r)(\theta) = \frac{1}{2\pi} \int_{-\pi}^{\pi} g(t) P_r(\theta - t) dt,\end{aligned}$$

where we have the Poisson kernel:

$$P_r(\theta) = P(r, \theta) = \sum_{j=-\infty}^{\infty} r^{|j|} e^{ij\theta}.$$



# Poisson kernel

$$P_r(\theta) = P(r, \theta) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2}, \quad 0 \leq r < 1, \quad -\pi \leq \theta < \pi.$$

Some properties:

# Poisson kernel

$$P_r(\theta) = P(r, \theta) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2}, \quad 0 \leq r < 1, \quad -\pi \leq \theta < \pi.$$

Some properties:

►  $P_r(\theta) \geq 0$

# Poisson kernel

$$P_r(\theta) = P(r, \theta) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2}, \quad 0 \leq r < 1, \quad -\pi \leq \theta < \pi.$$

Some properties:

►  $P_r(\theta) \geq 0$



$$\frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(\theta) d\theta = 1$$

# Poisson kernel

$$P_r(\theta) = P(r, \theta) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2}, \quad 0 \leq r < 1, \quad -\pi \leq \theta < \pi.$$

Some properties:

►  $P_r(\theta) \geq 0$



$$\frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(\theta) d\theta = 1$$

► Observe that

$$P(r, 0) = \frac{1 - r^2}{1 - 2r + r^2}, \quad 0 \leq r < 1,$$

consequently  $\lim_{r \rightarrow 1} P(r, 0) = \infty$ .

However  $\lim_{r \rightarrow 1} P(r, \theta) = 0$  when  $\theta \neq 0$ .

# Poisson kernel

$$P_r(\theta) = P(r, \theta) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2}, \quad 0 \leq r < 1, \quad -\pi \leq \theta < \pi.$$

Some properties:

►  $P_r(\theta) \geq 0$



$$\frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(\theta) d\theta = 1$$

► Observe that

$$P(r, 0) = \frac{1 - r^2}{1 - 2r + r^2}, \quad 0 \leq r < 1,$$

consequently  $\lim_{r \rightarrow 1} P(r, 0) = \infty$ .

However  $\lim_{r \rightarrow 1} P(r, \theta) = 0$  when  $\theta \neq 0$ .

► This formula does not work when  $r = 1$ !