

MS-C2111 Stochastic Processes



Lecture 2

Long-term behavior of Markov chains

Jukka Kohonen
Aalto University

Contents

Invariant and limiting distributions

Examples

Connectivity = Irreducibility

Periodicity

Convergence

Reducible MCs

Long-term behavior

What can we say about the state of a Markov chain:

$$\mathbb{P}_x(X_t = y) \approx ? \quad \text{for } t \text{ large?}$$

Long-term behavior

What can we say about the state of a Markov chain:

$$\mathbb{P}_x(X_t = y) \approx ? \quad \text{for } t \text{ large?}$$

- Does the distribution of X_t have a limit as $t \rightarrow \infty$?

Long-term behavior

What can we say about the state of a Markov chain:

$$\mathbb{P}_x(X_t = y) \approx ? \quad \text{for } t \text{ large?}$$

- Does the distribution of X_t have a limit as $t \rightarrow \infty$?
- If yes, does it depend on the initial state x ?

Long-term behavior

What can we say about the state of a Markov chain:

$$\mathbb{P}_x(X_t = y) \approx ? \quad \text{for } t \text{ large?}$$

- Does the distribution of X_t have a limit as $t \rightarrow \infty$?
- If yes, does it depend on the initial state x ?
- If yes, how can it be computed?

Invariant distribution

An **invariant distribution** (or *stationary* or *equilibrium* distribution) of a transition matrix P and a corresponding Markov chain is a row vector π such that $\pi(x) \geq 0$, $\sum_x \pi(x) = 1$, and

$$\pi = \pi P,$$

Invariant distribution

An **invariant distribution** (or *stationary* or *equilibrium* distribution) of a transition matrix **P** and a corresponding Markov chain is a row vector **π** such that $\pi(x) \geq 0$, $\sum_x \pi(x) = 1$, and

$$\pi = \pi P,$$

that is,

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad y \in S.$$

Invariant distribution

An **invariant distribution** (or *stationary* or *equilibrium* distribution) of a transition matrix **P** and a corresponding Markov chain is a row vector **π** such that $\pi(x) \geq 0$, $\sum_x \pi(x) = 1$, and

$$\pi = \pi P,$$

that is,

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad y \in S.$$

Note

If the initial state is random and such that $\mu_0 = \pi$, then

$$\mu_t =$$

Invariant distribution

An **invariant distribution** (or *stationary* or *equilibrium* distribution) of a transition matrix P and a corresponding Markov chain is a row vector π such that $\pi(x) \geq 0$, $\sum_x \pi(x) = 1$, and

$$\pi = \pi P,$$

that is,

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad y \in S.$$

Note

If the initial state is random and such that $\mu_0 = \pi$, then

$$\mu_t = \pi P^t$$

Invariant distribution

An **invariant distribution** (or *stationary* or *equilibrium* distribution) of a transition matrix **P** and a corresponding Markov chain is a row vector **π** such that $\pi(x) \geq 0$, $\sum_x \pi(x) = 1$, and

$$\pi = \pi P,$$

that is,

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad y \in S.$$

Note

If the initial state is random and such that $\mu_0 = \pi$, then

$$\mu_t = \pi P^t = (\pi P) P^{t-1}$$

Invariant distribution

An **invariant distribution** (or *stationary* or *equilibrium* distribution) of a transition matrix P and a corresponding Markov chain is a row vector π such that $\pi(x) \geq 0$, $\sum_x \pi(x) = 1$, and

$$\pi = \pi P,$$

that is,

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad y \in S.$$

Note

If the initial state is random and such that $\mu_0 = \pi$, then

$$\mu_t = \pi P^t = (\pi P) P^{t-1} = \pi P^{t-1}$$

Invariant distribution

An **invariant distribution** (or *stationary* or *equilibrium* distribution) of a transition matrix P and a corresponding Markov chain is a row vector π such that $\pi(x) \geq 0$, $\sum_x \pi(x) = 1$, and

$$\pi = \pi P,$$

that is,

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad y \in S.$$

Note

If the initial state is random and such that $\mu_0 = \pi$, then

$$\mu_t = \pi P^t = (\pi P) P^{t-1} = \pi P^{t-1} = \dots = \pi P$$

Invariant distribution

An **invariant distribution** (or *stationary* or *equilibrium* distribution) of a transition matrix P and a corresponding Markov chain is a row vector π such that $\pi(x) \geq 0$, $\sum_x \pi(x) = 1$, and

$$\pi = \pi P,$$

that is,

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad y \in S.$$

Note

If the initial state is random and such that $\mu_0 = \pi$, then

$$\mu_t = \pi P^t = (\pi P) P^{t-1} = \pi P^{t-1} = \dots = \pi P = \pi.$$

Invariant distribution

An **invariant distribution** (or *stationary* or *equilibrium* distribution) of a transition matrix P and a corresponding Markov chain is a row vector π such that $\pi(x) \geq 0$, $\sum_x \pi(x) = 1$, and

$$\pi = \pi P,$$

that is,

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad y \in S.$$

Note

If the initial state is random and such that $\mu_0 = \pi$, then

$$\mu_t = \pi P^t = (\pi P) P^{t-1} = \pi P^{t-1} = \dots = \pi P = \pi.$$

Hence X_t is π -distributed for all t .

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1}$$

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1} = (\mu_0 P^t)P$$

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1} = (\mu_0 P^t)P = \mu_t P,$$

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1} = (\mu_0 P^t) P = \mu_t P,$$

that is
$$\mu_{t+1}(y) = \sum_{x \in S} \mu_t(x) P(x, y).$$

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1} = (\mu_0 P^t) P = \mu_t P,$$

$$\text{that is} \quad \mu_{t+1}(y) = \sum_{x \in S} \mu_t(x) P(x, y).$$

If $\mu_t(x) \rightarrow \pi(x)$ for all x , then

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1} = (\mu_0 P^t) P = \mu_t P,$$

$$\text{that is} \quad \mu_{t+1}(y) = \sum_{x \in S} \mu_t(x) P(x, y).$$

If $\mu_t(x) \rightarrow \pi(x)$ for all x , then

$$\pi(y) =$$

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1} = (\mu_0 P^t) P = \mu_t P,$$

$$\text{that is} \quad \mu_{t+1}(y) = \sum_{x \in S} \mu_t(x) P(x, y).$$

If $\mu_t(x) \rightarrow \pi(x)$ for all x , then

$$\pi(y) = \lim_{t \rightarrow \infty} \mu_{t+1}(y)$$

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1} = (\mu_0 P^t) P = \mu_t P,$$

$$\text{that is} \quad \mu_{t+1}(y) = \sum_{x \in S} \mu_t(x) P(x, y).$$

If $\mu_t(x) \rightarrow \pi(x)$ for all x , then

$$\pi(y) = \lim_{t \rightarrow \infty} \mu_{t+1}(y) = \sum_{x \in S} \lim_{t \rightarrow \infty} \mu_t(x) P(x, y)$$

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1} = (\mu_0 P^t) P = \mu_t P,$$

$$\text{that is} \quad \mu_{t+1}(y) = \sum_{x \in S} \mu_t(x) P(x, y).$$

If $\mu_t(x) \rightarrow \pi(x)$ for all x , then

$$\pi(y) = \lim_{t \rightarrow \infty} \mu_{t+1}(y) = \sum_{x \in S} \lim_{t \rightarrow \infty} \mu_t(x) P(x, y) = \sum_{x \in S} \pi(x) P(x, y).$$

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1} = (\mu_0 P^t) P = \mu_t P,$$

$$\text{that is} \quad \mu_{t+1}(y) = \sum_{x \in S} \mu_t(x) P(x, y).$$

If $\mu_t(x) \rightarrow \pi(x)$ for all x , then

$$\pi(y) = \lim_{t \rightarrow \infty} \mu_{t+1}(y) = \sum_{x \in S} \lim_{t \rightarrow \infty} \mu_t(x) P(x, y) = \sum_{x \in S} \pi(x) P(x, y).$$

Hence $\pi P = \pi$.

Limiting distribution is an invariant distribution

Theorem

Any limiting distribution π of a finite-state Markov chain is also an invariant distribution.

Proof.

$$\mu_{t+1} = \mu_0 P^{t+1} = (\mu_0 P^t) P = \mu_t P,$$

$$\text{that is } \mu_{t+1}(y) = \sum_{x \in S} \mu_t(x) P(x, y).$$

If $\mu_t(x) \rightarrow \pi(x)$ for all x , then

$$\pi(y) = \lim_{t \rightarrow \infty} \mu_{t+1}(y) = \sum_{x \in S} \lim_{t \rightarrow \infty} \mu_t(x) P(x, y) = \sum_{x \in S} \pi(x) P(x, y).$$

Hence $\pi P = \pi$. Analogously $\implies \sum_{x \in S} \pi(x) = 1$. □

Contents

Invariant and limiting distributions

Examples

Connectivity = Irreducibility

Periodicity

Convergence

Reducible MCs

Example 1: Brand loyalty

A smartphone market is dominated by three manufacturers. When buying a new phone, a customer chooses a phone from the same manufacturer i as the previous one with probability β_i , and otherwise the customer randomly chooses one of the other manufacturers.

Company i	1	2	3
Loyalty β_i	0.8	0.6	0.4

Example 1: Brand loyalty

A smartphone market is dominated by three manufacturers. When buying a new phone, a customer chooses a phone from the same manufacturer i as the previous one with probability β_i , and otherwise the customer randomly chooses one of the other manufacturers.

Company i	1	2	3
Loyalty β_i	0.8	0.6	0.4

Assume that all smartphones have the same lifetime regardless of the manufacturer. Will the market shares of the manufacturers stabilize in the long run?

Example 1: Brand loyalty

Company i	1	2	3
Loyalty β_i	0.8	0.6	0.4

Let us model the manufacturer of a typical customer's phone after the t -th purchase instant by a Markov chain (X_0, X_1, \dots) with state space $S = \{1, 2, 3\}$ and transition matrix

$$P = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$$

Example 1: Brand loyalty

Company i	1	2	3
Loyalty β_i	0.8	0.6	0.4

Let us model the manufacturer of a typical customer's phone after the t -th purchase instant by a Markov chain (X_0, X_1, \dots) with state space $S = \{1, 2, 3\}$ and transition matrix

$$P = \begin{bmatrix} 0.8 & & \\ & & \\ & & \end{bmatrix}$$

Example 1: Brand loyalty

Company i	1	2	3
Loyalty β_i	0.8	0.6	0.4

Let us model the manufacturer of a typical customer's phone after the t -th purchase instant by a Markov chain (X_0, X_1, \dots) with state space $S = \{1, 2, 3\}$ and transition matrix

$$P = \begin{bmatrix} 0.8 & 0.1 \\ & & \end{bmatrix}$$

Example 1: Brand loyalty

Company i	1	2	3
Loyalty β_i	0.8	0.6	0.4

Let us model the manufacturer of a typical customer's phone after the t -th purchase instant by a Markov chain (X_0, X_1, \dots) with state space $S = \{1, 2, 3\}$ and transition matrix

$$P = \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ & & \end{bmatrix}$$

Example 1: Brand loyalty

Company i	1	2	3
Loyalty β_i	0.8	0.6	0.4

Let us model the manufacturer of a typical customer's phone after the t -th purchase instant by a Markov chain (X_0, X_1, \dots) with state space $S = \{1, 2, 3\}$ and transition matrix

$$P = \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \end{bmatrix}$$

Example 1: Brand loyalty

Company i	1	2	3
Loyalty β_i	0.8	0.6	0.4

Let us model the manufacturer of a typical customer's phone after the t -th purchase instant by a Markov chain (X_0, X_1, \dots) with state space $S = \{1, 2, 3\}$ and transition matrix

$$P = \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

Example 1: Brand loyalty

Company i	1	2	3
Loyalty β_i	0.8	0.6	0.4

Let us model the manufacturer of a typical customer's phone after the t -th purchase instant by a Markov chain (X_0, X_1, \dots) with state space $S = \{1, 2, 3\}$ and transition matrix

$$P = \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

Question

What happens to $\mu_t = \mu_0 P^t$ in the long run as $t \rightarrow \infty$?

Example 1: Brand loyalty

Compute powers of the transition matrix:

$$P = \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

Example 1: Brand loyalty

Compute powers of the transition matrix:

$$P = \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

$$P^2 = \begin{bmatrix} 0.69 & 0.17 & 0.14 \\ 0.34 & 0.44 & 0.22 \\ 0.42 & 0.33 & 0.25 \end{bmatrix}$$

Example 1: Brand loyalty

Compute powers of the transition matrix:

$$P = \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

$$P^2 = \begin{bmatrix} 0.69 & 0.17 & 0.14 \\ 0.34 & 0.44 & 0.22 \\ 0.42 & 0.33 & 0.25 \end{bmatrix}$$

$$P^{10} = \begin{bmatrix} 0.5471287 & 0.2715017 & 0.1813696 \\ 0.5430034 & 0.2745217 & 0.1824748 \\ 0.5441087 & 0.2737123 & 0.1821790 \end{bmatrix}$$

Example 1: Brand loyalty

Compute powers of the transition matrix:

$$P = \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

$$P^2 = \begin{bmatrix} 0.69 & 0.17 & 0.14 \\ 0.34 & 0.44 & 0.22 \\ 0.42 & 0.33 & 0.25 \end{bmatrix}$$

$$P^{10} = \begin{bmatrix} 0.5471287 & 0.2715017 & 0.1813696 \\ 0.5430034 & 0.2745217 & 0.1824748 \\ 0.5441087 & 0.2737123 & 0.1821790 \end{bmatrix}$$

$$P^{20} = \begin{bmatrix} 0.5454610 & 0.2727226 & 0.1818165 \\ 0.5454452 & 0.2727341 & 0.1818207 \\ 0.5454494 & 0.2727310 & 0.1818196 \end{bmatrix}$$

Example 1: Brand loyalty

$$P^{20} = \begin{bmatrix} 0.5454610 & 0.2727226 & 0.1818165 \\ 0.5454452 & 0.2727341 & 0.1818207 \\ 0.5454494 & 0.2727310 & 0.1818196 \end{bmatrix}$$

After 20 phone purchases:

- An initial customer of brand 1 is a customer of brand 2 with probability

Example 1: Brand loyalty

$$P^{20} = \begin{bmatrix} 0.5454610 & 0.2727226 & 0.1818165 \\ 0.5454452 & 0.2727341 & 0.1818207 \\ 0.5454494 & 0.2727310 & 0.1818196 \end{bmatrix}$$

After 20 phone purchases:

- An initial customer of brand 1 is a customer of brand 2 with probability $P^{20}(1, 2) \approx 0.273$

Example 1: Brand loyalty

$$P^{20} = \begin{bmatrix} 0.5454610 & 0.2727226 & 0.1818165 \\ 0.5454452 & 0.2727341 & 0.1818207 \\ 0.5454494 & 0.2727310 & 0.1818196 \end{bmatrix}$$

After 20 phone purchases:

- An initial customer of brand 1 is a customer of brand 2 with probability $P^{20}(1, 2) \approx 0.273$
- An initial customer of brand 2 is a customer of brand 2 with probability

Example 1: Brand loyalty

$$P^{20} = \begin{bmatrix} 0.5454610 & 0.2727226 & 0.1818165 \\ 0.5454452 & 0.2727341 & 0.1818207 \\ 0.5454494 & 0.2727310 & 0.1818196 \end{bmatrix}$$

After 20 phone purchases:

- An initial customer of brand 1 is a customer of brand 2 with probability $P^{20}(1, 2) \approx 0.273$
- An initial customer of brand 2 is a customer of brand 2 with probability $P^{20}(2, 2) \approx 0.273$

Example 1: Brand loyalty

$$P^{20} = \begin{bmatrix} 0.5454610 & 0.2727226 & 0.1818165 \\ 0.5454452 & 0.2727341 & 0.1818207 \\ 0.5454494 & 0.2727310 & 0.1818196 \end{bmatrix}$$

After 20 phone purchases:

- An initial customer of brand 1 is a customer of brand 2 with probability $P^{20}(1, 2) \approx 0.273$
- An initial customer of brand 2 is a customer of brand 2 with probability $P^{20}(2, 2) \approx 0.273$
- An initial customer of brand 3 is a customer of brand 2 with probability

Example 1: Brand loyalty

$$P^{20} = \begin{bmatrix} 0.5454610 & 0.2727226 & 0.1818165 \\ 0.5454452 & 0.2727341 & 0.1818207 \\ 0.5454494 & 0.2727310 & 0.1818196 \end{bmatrix}$$

After 20 phone purchases:

- An initial customer of brand 1 is a customer of brand 2 with probability $P^{20}(1, 2) \approx 0.273$
- An initial customer of brand 2 is a customer of brand 2 with probability $P^{20}(2, 2) \approx 0.273$
- An initial customer of brand 3 is a customer of brand 2 with probability $P^{20}(3, 2) \approx 0.273$

Example 1: Brand loyalty

$$P^{20} = \begin{bmatrix} 0.5454610 & 0.2727226 & 0.1818165 \\ 0.5454452 & 0.2727341 & 0.1818207 \\ 0.5454494 & 0.2727310 & 0.1818196 \end{bmatrix}$$

After 20 phone purchases:

- An initial customer of brand 1 is a customer of brand 2 with probability $P^{20}(1, 2) \approx 0.273$
- An initial customer of brand 2 is a customer of brand 2 with probability $P^{20}(2, 2) \approx 0.273$
- An initial customer of brand 3 is a customer of brand 2 with probability $P^{20}(3, 2) \approx 0.273$

The relevance of the initial state is negligible after 20 purchases.

Example 1: Brand loyalty

$$P^{20} = \begin{bmatrix} 0.5454610 & 0.2727226 & 0.1818165 \\ 0.5454452 & 0.2727341 & 0.1818207 \\ 0.5454494 & 0.2727310 & 0.1818196 \end{bmatrix}$$

After 20 phone purchases:

- An initial customer of brand 1 is a customer of brand 2 with probability $P^{20}(1, 2) \approx 0.273$
- An initial customer of brand 2 is a customer of brand 2 with probability $P^{20}(2, 2) \approx 0.273$
- An initial customer of brand 3 is a customer of brand 2 with probability $P^{20}(3, 2) \approx 0.273$

The relevance of the initial state is negligible after 20 purchases.

\implies Market shares seem to stabilize to

$$[0.545, 0.273, 0.182]$$

Example 1: Brand loyalty

Invariant distribution is a solution of

$$[\pi(1), \pi(2), \pi(3)] = [\pi(1), \pi(2), \pi(3)] \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

Example 1: Brand loyalty

Invariant distribution is a solution of

$$[\pi(1), \pi(2), \pi(3)] = [\pi(1), \pi(2), \pi(3)] \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

$$\pi(1) = \frac{8}{10}\pi(1) + \frac{2}{10}\pi(2) + \frac{3}{10}\pi(3)$$

$$\pi(2) = \frac{1}{10}\pi(1) + \frac{6}{10}\pi(2) + \frac{3}{10}\pi(3)$$

$$\pi(3) = \frac{1}{10}\pi(1) + \frac{2}{10}\pi(2) + \frac{4}{10}\pi(3)$$

Example 1: Brand loyalty

Invariant distribution is a solution of

$$[\pi(1), \pi(2), \pi(3)] = [\pi(1), \pi(2), \pi(3)] \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

$$\pi(1) = \frac{8}{10}\pi(1) + \frac{2}{10}\pi(2) + \frac{3}{10}\pi(3)$$

$$\pi(2) = \frac{1}{10}\pi(1) + \frac{6}{10}\pi(2) + \frac{3}{10}\pi(3)$$

$$\pi(3) = \frac{1}{10}\pi(1) + \frac{2}{10}\pi(2) + \frac{4}{10}\pi(3)$$

(recall also: $\pi(1) + \pi(2) + \pi(3) = 1$)

Example 1: Brand loyalty

Invariant distribution is a solution of

$$[\pi(1), \pi(2), \pi(3)] = [\pi(1), \pi(2), \pi(3)] \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

$$\pi(1) = \frac{8}{10}\pi(1) + \frac{2}{10}\pi(2) + \frac{3}{10}\pi(3)$$

$$\pi(2) = \frac{1}{10}\pi(1) + \frac{6}{10}\pi(2) + \frac{3}{10}\pi(3)$$

$$\pi(3) = \frac{1}{10}\pi(1) + \frac{2}{10}\pi(2) + \frac{4}{10}\pi(3)$$

(recall also: $\pi(1) + \pi(2) + \pi(3) = 1$)

$$\implies \pi = \left[\frac{6}{11}, \frac{3}{11}, \frac{2}{11} \right]$$

Example 1: Brand loyalty

Invariant distribution is a solution of

$$[\pi(1), \pi(2), \pi(3)] = [\pi(1), \pi(2), \pi(3)] \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

$$\pi(1) = \frac{8}{10}\pi(1) + \frac{2}{10}\pi(2) + \frac{3}{10}\pi(3)$$

$$\pi(2) = \frac{1}{10}\pi(1) + \frac{6}{10}\pi(2) + \frac{3}{10}\pi(3)$$

$$\pi(3) = \frac{1}{10}\pi(1) + \frac{2}{10}\pi(2) + \frac{4}{10}\pi(3)$$

(recall also: $\pi(1) + \pi(2) + \pi(3) = 1$)

$$\Rightarrow \pi = \left[\frac{6}{11}, \frac{3}{11}, \frac{2}{11} \right] \approx [0.5454, 0.2727, 0.1818]$$

Example 1: Brand loyalty

Invariant distribution is a solution of

$$[\pi(1), \pi(2), \pi(3)] = [\pi(1), \pi(2), \pi(3)] \begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

$$\pi(1) = \frac{8}{10}\pi(1) + \frac{2}{10}\pi(2) + \frac{3}{10}\pi(3)$$

$$\pi(2) = \frac{1}{10}\pi(1) + \frac{6}{10}\pi(2) + \frac{3}{10}\pi(3)$$

$$\pi(3) = \frac{1}{10}\pi(1) + \frac{2}{10}\pi(2) + \frac{4}{10}\pi(3)$$

(recall also: $\pi(1) + \pi(2) + \pi(3) = 1$)


$$\implies \pi = \left[\frac{6}{11}, \frac{3}{11}, \frac{2}{11}\right] \approx [0.5454, 0.2727, 0.1818]$$

Conclusion: Market shares appear to stabilize to the invariant distribution π

Example 2: Ehrenfest¹ chain

N gas particles in two containers (left, right) connected by a tiny hole.

- Each round a random particle jumps from a container to another


¹Paul (1880–1993) and Tatyana (1867–1964) Ehrenfest 

Example 2: Ehrenfest¹ chain

N gas particles in two containers (left, right) connected by a tiny hole.

- Each round a random particle jumps from a container to another
- X_t = number of particles in the left container after t rounds is a Markov chain with transition probabilities

$$P = \begin{bmatrix} 0 & 4/4 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 4/4 & 0 \end{bmatrix} \quad N = 4$$




¹Paul (1880–1993) and Tatyana (1867–1964) Ehrenfest

Example 2: Ehrenfest¹ chain

N gas particles in two containers (left, right) connected by a tiny hole.

- Each round a random particle jumps from a container to another
- X_t = number of particles in the left container after t rounds is a Markov chain with transition probabilities

$$P = \begin{bmatrix} 0 & 4/4 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 4/4 & 0 \end{bmatrix} \quad N = 4$$



$$P(i, i-1) = \mathbb{P}(X_{t+1} = i-1 | X_t = i) =$$


¹Paul (1880–1993) and Tatyana (1867–1964) Ehrenfest

Example 2: Ehrenfest¹ chain

N gas particles in two containers (left, right) connected by a tiny hole.

- Each round a random particle jumps from a container to another
- X_t = number of particles in the left container after t rounds is a Markov chain with transition probabilities

$$P = \begin{bmatrix} 0 & 4/4 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 4/4 & 0 \end{bmatrix} \quad N = 4$$



$$P(i, i-1) = \mathbb{P}(X_{t+1} = i-1 | X_t = i) = \frac{i}{N}, \quad 1 \leq i \leq N$$


¹Paul (1880–1993) and Tatyana (1867–1964) Ehrenfest

Example 2: Ehrenfest¹ chain

N gas particles in two containers (left, right) connected by a tiny hole.

- Each round a random particle jumps from a container to another
- X_t = number of particles in the left container after t rounds is a Markov chain with transition probabilities

$$P = \begin{bmatrix} 0 & 4/4 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 4/4 & 0 \end{bmatrix} \quad N = 4$$



$$P(i, i-1) = \mathbb{P}(X_{t+1} = i-1 | X_t = i) = \frac{i}{N}, \quad 1 \leq i \leq N$$

$$P(i, i+1) = \mathbb{P}(X_{t+1} = i+1 | X_t = i) =$$


¹Paul (1880–1993) and Tatyana (1867–1964) Ehrenfest

Example 2: Ehrenfest¹ chain

N gas particles in two containers (left, right) connected by a tiny hole.

- Each round a random particle jumps from a container to another
- X_t = number of particles in the left container after t rounds is a Markov chain with transition probabilities

$$P = \begin{bmatrix} 0 & 4/4 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 4/4 & 0 \end{bmatrix} \quad N = 4$$



$$P(i, i-1) = \mathbb{P}(X_{t+1} = i-1 | X_t = i) = \frac{i}{N}, \quad 1 \leq i \leq N$$

$$P(i, i+1) = \mathbb{P}(X_{t+1} = i+1 | X_t = i) = \frac{N-i}{N}, \quad 0 \leq i \leq N-1$$

¹Paul (1880–1993) and Tatyana (1867–1964) Ehrenfest

... Example 2: Ehrenfest chain

Compute powers of P ...

$$P = \frac{1}{4} \begin{bmatrix} 0 & 4 & 0 & 0 & 0 \\ 1 & 0 & 3 & 0 & 0 \\ 0 & 2 & 0 & 2 & 0 \\ 0 & 0 & 3 & 0 & 1 \\ 0 & 0 & 0 & 4 & 0 \end{bmatrix} \quad P^2 = \left(\frac{1}{4}\right)^2 \begin{bmatrix} 4 & 0 & 12 & 0 & 0 \\ 0 & 10 & 0 & 6 & 0 \\ 2 & 0 & 12 & 0 & 2 \\ 0 & 6 & 0 & 10 & 0 \\ 0 & 0 & 12 & 0 & 4 \end{bmatrix}$$

... Example 2: Ehrenfest chain

Compute powers of P ...

$$P = \frac{1}{4} \begin{bmatrix} 0 & 4 & 0 & 0 & 0 \\ 1 & 0 & 3 & 0 & 0 \\ 0 & 2 & 0 & 2 & 0 \\ 0 & 0 & 3 & 0 & 1 \\ 0 & 0 & 0 & 4 & 0 \end{bmatrix} \quad P^2 = \left(\frac{1}{4}\right)^2 \begin{bmatrix} 4 & 0 & 12 & 0 & 0 \\ 0 & 10 & 0 & 6 & 0 \\ 2 & 0 & 12 & 0 & 2 \\ 0 & 6 & 0 & 10 & 0 \\ 0 & 0 & 12 & 0 & 4 \end{bmatrix}$$

$$P^3 = \left(\frac{1}{4}\right)^3 \begin{bmatrix} 0 & 40 & 0 & 24 & 0 \\ 10 & 0 & 48 & 0 & 6 \\ 0 & 32 & 0 & 32 & 0 \\ 6 & 0 & 48 & 0 & 10 \\ 0 & 24 & 0 & 40 & 0 \end{bmatrix} \quad P^4 = \left(\frac{1}{4}\right)^4 \begin{bmatrix} 40 & 0 & 192 & 0 & 24 \\ 0 & 136 & 0 & 120 & 0 \\ 32 & 0 & 192 & 0 & 32 \\ 0 & 120 & 0 & 136 & 0 \\ 24 & 0 & 192 & 0 & 40 \end{bmatrix}$$

... Example 2: Ehrenfest chain

Compute powers of P ...

$$P = \frac{1}{4} \begin{bmatrix} 0 & 4 & 0 & 0 & 0 \\ 1 & 0 & 3 & 0 & 0 \\ 0 & 2 & 0 & 2 & 0 \\ 0 & 0 & 3 & 0 & 1 \\ 0 & 0 & 0 & 4 & 0 \end{bmatrix} \quad P^2 = \left(\frac{1}{4}\right)^2 \begin{bmatrix} 4 & 0 & 12 & 0 & 0 \\ 0 & 10 & 0 & 6 & 0 \\ 2 & 0 & 12 & 0 & 2 \\ 0 & 6 & 0 & 10 & 0 \\ 0 & 0 & 12 & 0 & 4 \end{bmatrix}$$

$$P^3 = \left(\frac{1}{4}\right)^3 \begin{bmatrix} 0 & 40 & 0 & 24 & 0 \\ 10 & 0 & 48 & 0 & 6 \\ 0 & 32 & 0 & 32 & 0 \\ 6 & 0 & 48 & 0 & 10 \\ 0 & 24 & 0 & 40 & 0 \end{bmatrix} \quad P^4 = \left(\frac{1}{4}\right)^4 \begin{bmatrix} 40 & 0 & 192 & 0 & 24 \\ 0 & 136 & 0 & 120 & 0 \\ 32 & 0 & 192 & 0 & 32 \\ 0 & 120 & 0 & 136 & 0 \\ 24 & 0 & 192 & 0 & 40 \end{bmatrix}$$

- $P^t(x, x) = 0$ for odd $t = 1, 3, 5, \dots$ and $P^t(x, x) > 0$ for even $t = 2, 4, 6, \dots$

... Example 2: Ehrenfest chain

Compute powers of P ...

$$P = \frac{1}{4} \begin{bmatrix} 0 & 4 & 0 & 0 & 0 \\ 1 & 0 & 3 & 0 & 0 \\ 0 & 2 & 0 & 2 & 0 \\ 0 & 0 & 3 & 0 & 1 \\ 0 & 0 & 0 & 4 & 0 \end{bmatrix}$$

$$P^2 = \left(\frac{1}{4}\right)^2 \begin{bmatrix} 4 & 0 & 12 & 0 & 0 \\ 0 & 10 & 0 & 6 & 0 \\ 2 & 0 & 12 & 0 & 2 \\ 0 & 6 & 0 & 10 & 0 \\ 0 & 0 & 12 & 0 & 4 \end{bmatrix}$$

$$P^3 = \left(\frac{1}{4}\right)^3 \begin{bmatrix} 0 & 40 & 0 & 24 & 0 \\ 10 & 0 & 48 & 0 & 6 \\ 0 & 32 & 0 & 32 & 0 \\ 6 & 0 & 48 & 0 & 10 \\ 0 & 24 & 0 & 40 & 0 \end{bmatrix}$$

$$P^4 = \left(\frac{1}{4}\right)^4 \begin{bmatrix} 40 & 0 & 192 & 0 & 24 \\ 0 & 136 & 0 & 120 & 0 \\ 32 & 0 & 192 & 0 & 32 \\ 0 & 120 & 0 & 136 & 0 \\ 24 & 0 & 192 & 0 & 40 \end{bmatrix}$$

- $P^t(x, x) = 0$ for odd $t = 1, 3, 5, \dots$ and $P^t(x, x) > 0$ for even $t = 2, 4, 6, \dots$
- Periodic structure: If X_0 is even, then X_1 is odd, X_2 is even, \dots

... Example 2: Ehrenfest chain

Compute powers of P ...

$$P = \frac{1}{4} \begin{bmatrix} 0 & 4 & 0 & 0 & 0 \\ 1 & 0 & 3 & 0 & 0 \\ 0 & 2 & 0 & 2 & 0 \\ 0 & 0 & 3 & 0 & 1 \\ 0 & 0 & 0 & 4 & 0 \end{bmatrix} \quad P^2 = \left(\frac{1}{4}\right)^2 \begin{bmatrix} 4 & 0 & 12 & 0 & 0 \\ 0 & 10 & 0 & 6 & 0 \\ 2 & 0 & 12 & 0 & 2 \\ 0 & 6 & 0 & 10 & 0 \\ 0 & 0 & 12 & 0 & 4 \end{bmatrix}$$

$$P^3 = \left(\frac{1}{4}\right)^3 \begin{bmatrix} 0 & 40 & 0 & 24 & 0 \\ 10 & 0 & 48 & 0 & 6 \\ 0 & 32 & 0 & 32 & 0 \\ 6 & 0 & 48 & 0 & 10 \\ 0 & 24 & 0 & 40 & 0 \end{bmatrix} \quad P^4 = \left(\frac{1}{4}\right)^4 \begin{bmatrix} 40 & 0 & 192 & 0 & 24 \\ 0 & 136 & 0 & 120 & 0 \\ 32 & 0 & 192 & 0 & 32 \\ 0 & 120 & 0 & 136 & 0 \\ 24 & 0 & 192 & 0 & 40 \end{bmatrix}$$

- $P^t(x, x) = 0$ for odd $t = 1, 3, 5, \dots$ and $P^t(x, x) > 0$ for even $t = 2, 4, 6, \dots$
- Periodic structure: If X_0 is even, then X_1 is odd, X_2 is even, ...
- $P^t(x, x)$ does not converge, so neither does μ_t

Example 3: Many limits

The transition matrix

$$P = \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

satisfies $P^t = P$ for all $t = 1, 2, \dots$

Example 3: Many limits

The transition matrix

$$P = \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

satisfies $P^t = P$ for all $t = 1, 2, \dots$

When $X_0 = 1$ i.e. $\mu_0 = [1, 0, 0]$,

$$\mu_t = \mu_0 P^t = [0.5, 0.5, 0] \quad \text{for all } t \geq 1$$

Example 3: Many limits

The transition matrix

$$P = \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

satisfies $P^t = P$ for all $t = 1, 2, \dots$

When $X_0 = 1$ i.e. $\mu_0 = [1, 0, 0]$,

$$\mu_t = \mu_0 P^t = [0.5, 0.5, 0] \quad \text{for all } t \geq 1$$

When $X_0 = 3$ i.e. $\mu_0 = [0, 0, 1]$,

$$\mu_t = [0, 0, 1] P^t = [0, 0, 1] \quad \text{for all } t \geq 0$$

Example 3: Many limits

The transition matrix

$$P = \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

satisfies $P^t = P$ for all $t = 1, 2, \dots$

When $X_0 = 1$ i.e. $\mu_0 = [1, 0, 0]$,

$$\mu_t = \mu_0 P^t = [0.5, 0.5, 0] \quad \text{for all } t \geq 1$$

When $X_0 = 3$ i.e. $\mu_0 = [0, 0, 1]$,

$$\mu_t = [0, 0, 1] P^t = [0, 0, 1] \quad \text{for all } t \geq 0$$

This MC has many limiting distributions, depending on initial state.

Example 3: Many limits

The transition matrix

$$P = \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

satisfies $P^t = P$ for all $t = 1, 2, \dots$

When $X_0 = 1$ i.e. $\mu_0 = [1, 0, 0]$,

$$\mu_t = \mu_0 P^t = [0.5, 0.5, 0] \quad \text{for all } t \geq 1$$

When $X_0 = 3$ i.e. $\mu_0 = [0, 0, 1]$,

$$\mu_t = [0, 0, 1] P^t = [0, 0, 1] \quad \text{for all } t \geq 0$$

This MC has many limiting distributions, depending on initial state.

For all $\alpha \in [0, 1]$, the distribution

$$\pi = \alpha[0.5, 0.5, 0] + (1 - \alpha)[0, 0, 1]$$

is an invariant distribution.

Examples: Summary

Unique limiting
distribution

$$\begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

(Brand loyalty)

No limiting distribution

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

(Ehrenfest)

Many limiting
distributions

$$\begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Contents

Invariant and limiting distributions

Examples

Connectivity = Irreducibility

Periodicity

Convergence

Reducible MCs

Irreducible Markov chain

$x \rightsquigarrow y$ if the transition diagram contains a path from x to y

Irreducible Markov chain

$x \rightsquigarrow y$ if the transition diagram contains a path from x to y

The matrix and the MC is **irreducible** if $x \rightsquigarrow y$ for all x, y .

(In graph theory terms: **strongly connected**)

Irreducible Markov chain

$x \rightsquigarrow y$ if the transition diagram contains a path from x to y

The matrix and the MC is **irreducible** if $x \rightsquigarrow y$ for all x, y .

(In graph theory terms: **strongly connected**)

Which of the following are irreducible?

$$\begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer:

Irreducible Markov chain

$x \rightsquigarrow y$ if the transition diagram contains a path from x to y

The matrix and the MC is **irreducible** if $x \rightsquigarrow y$ for all x, y .

(In graph theory terms: **strongly connected**)

Which of the following are irreducible?

$$\begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer: 2 first are irreducible, 3rd is not.

Determining irreducibility

Theorem

For all $x \neq y$: $x \rightsquigarrow y$ if and only if $P^t(x, y) > 0$ for some $t \geq 1$.

Determining irreducibility

Theorem

For all $x \neq y$: $x \rightsquigarrow y$ if and only if $P^t(x, y) > 0$ for some $t \geq 1$.

Proof.

If $x \rightsquigarrow y$, then there is a path $x = x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_t = y$. Hence

$$P(x_0, x_1)P(x_1, x_2) \cdots P(x_{t-1}, x_t) > 0.$$

Determining irreducibility

Theorem

For all $x \neq y$: $x \rightsquigarrow y$ if and only if $P^t(x, y) > 0$ for some $t \geq 1$.

Proof.

If $x \rightsquigarrow y$, then there is a path $x = x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_t = y$. Hence

$$P(x_0, x_1)P(x_1, x_2) \cdots P(x_{t-1}, x_t) > 0.$$

Hence $P^t(x, y) = \mathbb{P}(X_t = y \mid X_0 = x)$



Determining irreducibility

Theorem

For all $x \neq y$: $x \rightsquigarrow y$ if and only if $P^t(x, y) > 0$ for some $t \geq 1$.

Proof.

If $x \rightsquigarrow y$, then there is a path $x = x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_t = y$. Hence

$$P(x_0, x_1)P(x_1, x_2) \cdots P(x_{t-1}, x_t) > 0.$$

Hence

$$\begin{aligned} P^t(x, y) &= \mathbb{P}(X_t = y \mid X_0 = x) \\ &= \mathbb{P}(X_t = x_t \mid X_0 = x_0) \end{aligned}$$



Determining irreducibility

Theorem

For all $x \neq y$: $x \rightsquigarrow y$ if and only if $P^t(x, y) > 0$ for some $t \geq 1$.

Proof.

If $x \rightsquigarrow y$, then there is a path $x = x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_t = y$. Hence

$$P(x_0, x_1)P(x_1, x_2) \cdots P(x_{t-1}, x_t) > 0.$$

Hence

$$\begin{aligned} P^t(x, y) &= \mathbb{P}(X_t = y \mid X_0 = x) \\ &= \mathbb{P}(X_t = x_t \mid X_0 = x_0) \\ &\geq \mathbb{P}(X_t = x_t, X_{t-1} = x_{t-1}, \dots, X_1 = x_1 \mid X_0 = x_0) \end{aligned}$$



Determining irreducibility

Theorem

For all $x \neq y$: $x \rightsquigarrow y$ if and only if $P^t(x, y) > 0$ for some $t \geq 1$.

Proof.

If $x \rightsquigarrow y$, then there is a path $x = x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_t = y$. Hence

$$P(x_0, x_1)P(x_1, x_2) \cdots P(x_{t-1}, x_t) > 0.$$

Hence

$$\begin{aligned} P^t(x, y) &= \mathbb{P}(X_t = y \mid X_0 = x) \\ &= \mathbb{P}(X_t = x_t \mid X_0 = x_0) \\ &\geq \mathbb{P}(X_t = x_t, X_{t-1} = x_{t-1}, \dots, X_1 = x_1 \mid X_0 = x_0) \\ &= P(x_0, x_1)P(x_1, x_2) \cdots P(x_{t-1}, x_t) \end{aligned}$$



Determining irreducibility

Theorem

For all $x \neq y$: $x \rightsquigarrow y$ if and only if $P^t(x, y) > 0$ for some $t \geq 1$.

Proof.

If $x \rightsquigarrow y$, then there is a path $x = x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_t = y$. Hence

$$P(x_0, x_1)P(x_1, x_2) \cdots P(x_{t-1}, x_t) > 0.$$

Hence

$$\begin{aligned} P^t(x, y) &= \mathbb{P}(X_t = y \mid X_0 = x) \\ &= \mathbb{P}(X_t = x_t \mid X_0 = x_0) \\ &\geq \mathbb{P}(X_t = x_t, X_{t-1} = x_{t-1}, \dots, X_1 = x_1 \mid X_0 = x_0) \\ &= P(x_0, x_1)P(x_1, x_2) \cdots P(x_{t-1}, x_t) \\ &> 0. \end{aligned}$$



Determining irreducibility

Theorem

For $x \neq y$: $x \rightsquigarrow y$ if and only if $P^t(x, y) > 0$ for some $t \geq 1$.

... Proof (converse).

If $P^t(x, y) > 0$ for some $t \geq 1$,

Determining irreducibility

Theorem

For $x \neq y$: $x \rightsquigarrow y$ if and only if $P^t(x, y) > 0$ for some $t \geq 1$.

... Proof (converse).

If $P^t(x, y) > 0$ for some $t \geq 1$, then $\mathbb{P}(X_t = y \mid X_0 = x) > 0$, that a MC starting from x is after t time steps at possibly at y .

Determining irreducibility

Theorem

For $x \neq y$: $x \rightsquigarrow y$ if and only if $P^t(x, y) > 0$ for some $t \geq 1$.

... Proof (converse).

If $P^t(x, y) > 0$ for some $t \geq 1$, then $\mathbb{P}(X_t = y \mid X_0 = x) > 0$, that a MC starting from x is after t time steps at possibly at y .

Hence there must be a path in the transition diagram, of length t , so $x \rightsquigarrow y$. □

Determining irreducibility

Theorem

A transition matrix P is irreducible if and only if for all $x \neq y$ there exists an integer $t \geq 1$, such that $P^t(x, y) > 0$.

Determining irreducibility

Theorem

A transition matrix P is irreducible if and only if for all $x \neq y$ there exists an integer $t \geq 1$, such that $P^t(x, y) > 0$.

Proof.

Directly from previous result, according to which
 $x \rightsquigarrow y$ iff $P^t(x, y) > 0$ for some $t \geq 1$.



Components

$x \leftrightarrow y$, if $x \rightsquigarrow y$ and $y \rightsquigarrow x$.

Components

$x \longleftrightarrow y$, if $x \rightsquigarrow y$ and $y \rightsquigarrow x$.

$C(x) = \{y : y \longleftrightarrow x\}$ is the component of state x

Components

$x \longleftrightarrow y$, if $x \rightsquigarrow y$ and $y \rightsquigarrow x$.

$C(x) = \{y : y \longleftrightarrow x\}$ is the component of state x

The components partition the state space.

$x \longleftrightarrow y$ is an equivalence relation (reflexive, symmetric, transitive)

Components

$x \longleftrightarrow y$, if $x \rightsquigarrow y$ and $y \rightsquigarrow x$.

$C(x) = \{y : y \longleftrightarrow x\}$ is the component of state x

The components partition the state space.

$x \longleftrightarrow y$ is an equivalence relation (reflexive, symmetric, transitive)

Determine the components of the following chains:

$$\begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer:

Components

$x \longleftrightarrow y$, if $x \rightsquigarrow y$ and $y \rightsquigarrow x$.

$C(x) = \{y : y \longleftrightarrow x\}$ is the component of state x

The components partition the state space.

$x \longleftrightarrow y$ is an equivalence relation (reflexive, symmetric, transitive)

Determine the components of the following chains:

$$\begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer: First two have only one component (full state space).

Components

$x \longleftrightarrow y$, if $x \rightsquigarrow y$ and $y \rightsquigarrow x$.

$C(x) = \{y : y \longleftrightarrow x\}$ is the component of state x

The components partition the state space.

$x \longleftrightarrow y$ is an equivalence relation (reflexive, symmetric, transitive)

Determine the components of the following chains:

$$\begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer: First two have only one component (full state space).

Third has two components $C(1) = C(2) = \{1, 2\}$ ja $C(3) = \{3\}$.

Invariant distribution

Theorem

Every irreducible finite-state MC has a unique invariant distribution π , for which $\pi(x) > 0$ for all x .

Proof.

Precise proof: [Dur12, Thm. 1.14] or [LPW08, Sec 1.5].

Invariant distribution

Theorem

Every irreducible finite-state MC has a unique invariant distribution π , for which $\pi(x) > 0$ for all x .

Proof.

Precise proof: [Dur12, Thm. 1.14] or [LPW08, Sec 1.5]. Existence:

$$\pi(x) = \frac{1}{\mathbb{E}(T_x^+ | X_0 = x)}, \quad T_x^+ = \min\{t \geq 1 : X_t = x\}.$$

Invariant distribution

Theorem

Every irreducible finite-state MC has a unique invariant distribution π , for which $\pi(x) > 0$ for all x .

Proof.

Precise proof: [Dur12, Thm. 1.14] or [LPW08, Sec 1.5]. Existence:

$$\pi(x) = \frac{1}{\mathbb{E}(T_x^+ | X_0 = x)}, \quad T_x^+ = \min\{t \geq 1 : X_t = x\}.$$

Uniqueness: for an irreducible chain the column vectors solving $Ph = h$ are of the form $h = [c, c, \dots, c]^T$, so the kernel of matrix $P - I$ is one-dimensional.

Invariant distribution

Theorem

Every irreducible finite-state MC has a unique invariant distribution π , for which $\pi(x) > 0$ for all x .

Proof.

Precise proof: [Dur12, Thm. 1.14] or [LPW08, Sec 1.5]. Existence:

$$\pi(x) = \frac{1}{\mathbb{E}(T_x^+ | X_0 = x)}, \quad T_x^+ = \min\{t \geq 1 : X_t = x\}.$$

Uniqueness: for an irreducible chain the column vectors solving $Ph = h$ are of the form $h = [c, c, \dots, c]^T$, so the kernel of matrix $P - I$ is one-dimensional. \implies also the solution space of the row-vector equation $\pi(P - I) = 0$ has dimension one.

Invariant distribution

Theorem

Every irreducible finite-state MC has a unique invariant distribution π , for which $\pi(x) > 0$ for all x .

Proof.

Precise proof: [Dur12, Thm. 1.14] or [LPW08, Sec 1.5]. Existence:

$$\pi(x) = \frac{1}{\mathbb{E}(T_x^+ | X_0 = x)}, \quad T_x^+ = \min\{t \geq 1 : X_t = x\}.$$

Uniqueness: for an irreducible chain the column vectors solving $Ph = h$ are of the form $h = [c, c, \dots, c]^T$, so the kernel of matrix $P - I$ is one-dimensional. \implies also the solution space of the row-vector equation $\pi(P - I) = 0$ has dimension one. Condition $\sum_x \pi(x) = 1 \implies$ that the equation $\pi = \pi P$ has at most one solution. □

When do the distributions μ_t converge

$$\mu_0 P^t \rightarrow \pi \quad \text{as } t \rightarrow \infty?$$

Contents

Invariant and limiting distributions

Examples

Connectivity = Irreducibility

Periodicity

Convergence

Reducible MCs

Period of a state

The **period** of state x is the gcd of the set of possible return times

$$\mathcal{T}_x = \{t \geq 1 : P^t(x, x) > 0\}$$

Period of a state

The **period** of state x is the gcd of the set of possible return times

$$\mathcal{T}_x = \{t \geq 1 : P^t(x, x) > 0\}$$

A transition matrix P and a corresponding MC is **aperiodic**, if all states have period one.

Period of a state

The **period** of state x is the gcd of the set of possible return times

$$\mathcal{T}_x = \{t \geq 1 : P^t(x, x) > 0\}$$

A transition matrix P and a corresponding MC is **aperiodic**, if all states have period one.

Which of the following are aperiodic?

$$\begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer:

Period of a state

The **period** of state x is the gcd of the set of possible return times $\mathcal{T}_x = \{t \geq 1 : P^t(x, x) > 0\}$

A transition matrix P and a corresponding MC is **aperiodic**, if all states have period one.

Which of the following are aperiodic?

$$\begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer: For the second chain $\mathcal{T}_x = \{2, 4, 6, \dots\}$ for all x , and every state has period 2.

Period of a state

The **period** of state x is the gcd of the set of possible return times $\mathcal{T}_x = \{t \geq 1 : P^t(x, x) > 0\}$

A transition matrix P and a corresponding MC is **aperiodic**, if all states have period one.

Which of the following are aperiodic?

$$\begin{bmatrix} 0.8 & 0.1 & 0.1 \\ 0.2 & 0.6 & 0.2 \\ 0.3 & 0.3 & 0.4 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1/4 & 0 & 3/4 & 0 & 0 \\ 0 & 2/4 & 0 & 2/4 & 0 \\ 0 & 0 & 3/4 & 0 & 1/4 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0.5 & 0.5 & 0 \\ 0.5 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer: For the second chain $\mathcal{T}_x = \{2, 4, 6, \dots\}$ for all x , and every state has period 2. The other chains are aperiodic.

Aperiodicity

Theorem

An irreducible MC is aperiodic if $P(x, x) > 0$ for some x .

Aperiodicity

Theorem

An irreducible MC is aperiodic if $P(x, x) > 0$ for some x .

Lemma

If $P(x, x) > 0$, then the period of x is 1.

Lemma

Every state of an irreducible chain has the same period.

Aperiodicity

Theorem

An irreducible MC is aperiodic if $P(x, x) > 0$ for some x .

Lemma

If $P(x, x) > 0$, then the period of x is 1.

Lemma

Every state of an irreducible chain has the same period.


Proof.

Exercise.




Example 4

What is the period of state 0?



Example 4

What is the period of state 0?




- From state 0 we move either to the left (L) or right (R) circuit

$$\mathcal{T}_0 = \left\{ \underbrace{3}_L, \underbrace{4}_R, \underbrace{6}_{LL}, \underbrace{7}_{LR}, \underbrace{8}_{RR}, \underbrace{9}_{LLL}, \dots \right\}$$

Example 4

What is the period of state 0?




- From state 0 we move either to the left (L) or right (R) circuit

$$\mathcal{T}_0 = \left\{ \underbrace{3}_L, \underbrace{4}_R, \underbrace{6}_{LL}, \underbrace{7}_{LR}, \underbrace{8}_{RR}, \underbrace{9}_{LLL}, \dots \right\}$$

- The gcd of \mathcal{T}_0 is 1 so the period of state 0 is 1.

Example 4

What is the period of state 0?



- From state 0 we move either to the left (L) or right (R) circuit


$$\mathcal{T}_0 = \left\{ \underbrace{3}_L, \underbrace{4}_R, \underbrace{6}_{LL}, \underbrace{7}_{LR}, \underbrace{8}_{RR}, \underbrace{9}_{LLL}, \dots \right\}$$

- The gcd of \mathcal{T}_0 is 1 so the period of state 0 is 1.

What is the period of state 1?

Example 4

What is the period of state 0?



- From state 0 we move either to the left (L) or right (R) circuit

$$\mathcal{T}_0 = \left\{ \underbrace{3}_L, \underbrace{4}_R, \underbrace{6}_{LL}, \underbrace{7}_{LR}, \underbrace{8}_{RR}, \underbrace{9}_{LLL}, \dots \right\}$$


- The gcd of \mathcal{T}_0 is 1 so the period of state 0 is 1.

What is the period of state 1?

- The chain is irreducible, so all states have the same period.

Example 4

What is the period of state 0?



- From state 0 we move either to the left (L) or right (R) circuit

$$\mathcal{T}_0 = \left\{ \underbrace{3}_L, \underbrace{4}_R, \underbrace{6}_{LL}, \underbrace{7}_{LR}, \underbrace{8}_{RR}, \underbrace{9}_{LLL}, \dots \right\}$$

- The gcd of \mathcal{T}_0 is 1 so the period of state 0 is 1.

What is the period of state 1?

- The chain is irreducible, so all states have the same period.
- The MC is hence irreducible and aperiodic.

Contents

Invariant and limiting distributions

Examples

Connectivity = Irreducibility

Periodicity

Convergence

Reducible MCs

Convergence theorem

Theorem

Let P be an irreducible and aperiodic transition matrix. Then for all x and y ,

$$P^t(x, y) \rightarrow \pi(y), \quad t \rightarrow \infty,$$

where π is the unique invariant distribution of P .

Convergence theorem

Theorem

Let P be an irreducible and aperiodic transition matrix. Then for all x and y ,

$$P^t(x, y) \rightarrow \pi(y), \quad t \rightarrow \infty,$$

where π is the unique invariant distribution of P .

Corollary: For an arbitrary initial distribution μ_0 , $\mu_t = \mu_0 P^t \rightarrow \pi$:

Convergence theorem

Theorem

Let P be an irreducible and aperiodic transition matrix. Then for all x and y ,

$$P^t(x, y) \rightarrow \pi(y), \quad t \rightarrow \infty,$$

where π is the unique invariant distribution of P .

Corollary: For an arbitrary initial distribution μ_0 , $\mu_t = \mu_0 P^t \rightarrow \pi$:

$$\mu_t(y)$$

Convergence theorem

Theorem

Let P be an irreducible and aperiodic transition matrix. Then for all x and y ,

$$P^t(x, y) \rightarrow \pi(y), \quad t \rightarrow \infty,$$

where π is the unique invariant distribution of P .

Corollary: For an arbitrary initial distribution μ_0 , $\mu_t = \mu_0 P^t \rightarrow \pi$:

$$\mu_t(y) = \sum_x \mu_0(x) P^t(x, y)$$

Convergence theorem

Theorem

Let P be an irreducible and aperiodic transition matrix. Then for all x and y ,

$$P^t(x, y) \rightarrow \pi(y), \quad t \rightarrow \infty,$$

where π is the unique invariant distribution of P .

Corollary: For an arbitrary initial distribution μ_0 , $\mu_t = \mu_0 P^t \rightarrow \pi$:

$$\mu_t(y) = \sum_x \mu_0(x) P^t(x, y) \rightarrow \sum_x \mu_0(x) \pi(y)$$

Convergence theorem

Theorem

Let P be an irreducible and aperiodic transition matrix. Then for all x and y ,

$$P^t(x, y) \rightarrow \pi(y), \quad t \rightarrow \infty,$$

where π is the unique invariant distribution of P .

Corollary: For an arbitrary initial distribution μ_0 , $\mu_t = \mu_0 P^t \rightarrow \pi$:

$$\mu_t(y) = \sum_x \mu_0(x) P^t(x, y) \rightarrow \sum_x \mu_0(x) \pi(y) = \pi(y).$$

Convergence theorem

Theorem

Let P be an irreducible and aperiodic transition matrix. Then for all x and y ,

$$P^t(x, y) \rightarrow \pi(y), \quad t \rightarrow \infty,$$

where π is the unique invariant distribution of P .

Corollary: For an arbitrary initial distribution μ_0 , $\mu_t = \mu_0 P^t \rightarrow \pi$:

$$\mu_t(y) = \sum_x \mu_0(x) P^t(x, y) \rightarrow \sum_x \mu_0(x) \pi(y) = \pi(y).$$

In matrix form:

$$\lim_{t \rightarrow \infty} P^t = \begin{bmatrix} \pi \\ \vdots \\ \pi \end{bmatrix}$$

Convergence theorem

Theorem

Let P be an irreducible and aperiodic transition matrix. Then for all x and y ,

$$P^t(x, y) \rightarrow \pi(y), \quad t \rightarrow \infty,$$

where π is the unique invariant distribution of P .

Corollary: For an arbitrary initial distribution μ_0 , $\mu_t = \mu_0 P^t \rightarrow \pi$:

$$\mu_t(y) = \sum_x \mu_0(x) P^t(x, y) \rightarrow \sum_x \mu_0(x) \pi(y) = \pi(y).$$

In matrix form:

$$\lim_{t \rightarrow \infty} P^t = \begin{bmatrix} \pi \\ \vdots \\ \pi \end{bmatrix}$$

Proof.

See [LPW08, Sec. 4.3] (online book!) or [Dur12, Sec. 1.7].



Example 5

$$P = \begin{bmatrix} 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 1 & 0 & 0 \end{bmatrix}$$

- Is the chain irreducible?

Example 5

$$P = \begin{bmatrix} 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 1 & 0 & 0 \end{bmatrix}$$

- Is the chain irreducible? Yes.

Example 5

$$P = \begin{bmatrix} 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 1 & 0 & 0 \end{bmatrix}$$

- Is the chain irreducible? Yes.
- Is the chain aperiodic?

Example 5

$$P = \begin{bmatrix} 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 1 & 0 & 0 \end{bmatrix}$$

- Is the chain irreducible? Yes.
- Is the chain aperiodic? Yes.

Example 5

$$P = \begin{bmatrix} 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 1 & 0 & 0 \end{bmatrix}$$

- Is the chain irreducible? Yes.
- Is the chain aperiodic? Yes.

Find π such that $\pi P = \pi$:

$$\frac{1}{2}\pi_2 + \pi_3 = \pi_1$$

$$\pi_1 = \pi_2$$

$$\frac{1}{2}\pi_2 = \pi_3.$$

Hence $\pi_2 = \pi_1$ and further $\pi_3 = \frac{1}{2}\pi_1$.

Example 5

$$P = \begin{bmatrix} 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 1 & 0 & 0 \end{bmatrix}$$

- Is the chain irreducible? Yes.
- Is the chain aperiodic? Yes.

Find π such that $\pi P = \pi$:

$$\frac{1}{2}\pi_2 + \pi_3 = \pi_1$$

$$\pi_1 = \pi_2$$

$$\frac{1}{2}\pi_2 = \pi_3.$$

Hence $\pi_2 = \pi_1$ and further $\pi_3 = \frac{1}{2}\pi_1$.

Condition $1 = \pi_1 + \pi_2 + \pi_3 = \pi_1(1 + 1 + \frac{1}{2})$ implies $\pi_1 = \frac{2}{5}$.

Example 5

$$P = \begin{bmatrix} 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 1 & 0 & 0 \end{bmatrix}$$

- Is the chain irreducible? Yes.
- Is the chain aperiodic? Yes.

Find π such that $\pi P = \pi$:

$$\frac{1}{2}\pi_2 + \pi_3 = \pi_1$$

$$\pi_1 = \pi_2$$

$$\frac{1}{2}\pi_2 = \pi_3.$$

Hence $\pi_2 = \pi_1$ and further $\pi_3 = \frac{1}{2}\pi_1$.

Condition $1 = \pi_1 + \pi_2 + \pi_3 = \pi_1(1 + 1 + \frac{1}{2})$ implies $\pi_1 = \frac{2}{5}$. Hence the MC has invariant distribution $\pi = (\frac{2}{5}, \frac{2}{5}, \frac{1}{5})$. This is unique.

Example 5

Using the convergence theorem:

$$\lim_{t \rightarrow \infty} P^t = \begin{bmatrix} \frac{2}{5} & \frac{2}{5} & \frac{1}{5} \\ \frac{2}{5} & \frac{2}{5} & \frac{1}{5} \\ \frac{2}{5} & \frac{2}{5} & \frac{1}{5} \end{bmatrix} \quad \left(\text{vrt. } P^{15} = \begin{bmatrix} 0.3984 & 0.3984 & 0.2031 \\ 0.4023 & 0.3984 & 0.1992 \\ 0.3984 & 0.4062 & 0.1953 \end{bmatrix} \right)$$

Contents

Invariant and limiting distributions

Examples

Connectivity = Irreducibility

Periodicity

Convergence

Reducible MCs

Reducible MCs

What happens for reducible MCs in the long run?

$$\mathbb{P}_x(X_t = y) \rightarrow ?$$

Note

- Reducible MCs have several components.
- The MC cannot exit an absorbing component.

Reducible MCs

What happens for reducible MCs in the long run?

$$\mathbb{P}_x(X_t = y) \rightarrow ?$$


Note

- Reducible MCs have several components.
- The MC cannot exit an absorbing component.

The long-term behavior of a reducible MC depends on the initial state X_0 .


Example 6

$$P = \begin{matrix} & \begin{matrix} 0 & 1 & 2 & 3 & 4 \end{matrix} \\ \begin{pmatrix} 1 \\ \frac{1}{3} \\ 0 \\ 0 \\ 0 \end{pmatrix} & \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{2}{3} & 0 & 0 & \frac{1}{3} \\ 0 & 0 & 0 & \frac{2}{3} & \frac{1}{3} \\ 0 & 0 & 0 & \frac{2}{3} & \frac{1}{3} \end{pmatrix} \end{matrix}$$



Example 6

$$P = \begin{matrix} & \begin{matrix} 0 & 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ \frac{1}{3} \\ 0 \\ 0 \\ 0 \end{matrix} & \begin{pmatrix} 0 & 1 & 2 & 3 & 4 \\ 1 & 0 & 0 & 0 & 0 \\ \frac{1}{3} & 0 & \frac{1}{3} & \frac{1}{3} & 0 \\ 0 & \frac{2}{3} & 0 & 0 & \frac{1}{3} \\ 0 & 0 & 0 & \frac{2}{3} & \frac{1}{3} \\ 0 & 0 & 0 & \frac{2}{3} & \frac{1}{3} \end{pmatrix} \end{matrix}$$



The states and components of the chain can be divided as:

- Absorbing components: $\{0\}$ ja $\{3, 4\}$
- Transient components: $\{1, 2\}$


... Example 6

A MC starting from 0 stays fixed to its initial state. Hence $\pi_{\{0\}} = [1, 0, 0, 0, 0]$ is an invariant distribution.

... Example 6

A MC starting from 0 stays fixed to its initial state. Hence $\pi_{\{0\}} = [1, 0, 0, 0, 0]$ is an invariant distribution.

A MC starting from the absorbing component $\{3, 4\}$ is like




This has invariant distribution $[2/3, 1/3]$.

... Example 6

A MC starting from 0 stays fixed to its initial state. Hence $\pi_{\{0\}} = [1, 0, 0, 0, 0]$ is an invariant distribution.

A MC starting from the absorbing component $\{3, 4\}$ is like




This has invariant distribution $[2/3, 1/3]$.

Also $\pi_{\{3,4\}} = [0, 0, 0, 2/3, 1/3]$ is an invariant distribution.

... Example 6

A MC starting from 0 stays fixed to its initial state. Hence $\pi_{\{0\}} = [1, 0, 0, 0, 0]$ is an invariant distribution.

A MC starting from the absorbing component $\{3, 4\}$ is like




This has invariant distribution $[2/3, 1/3]$.

Also $\pi_{\{3,4\}} = [0, 0, 0, 2/3, 1/3]$ is an invariant distribution.

Example 6

$$P = \begin{matrix} & \begin{matrix} 0 & 1 & 2 & 3 & 4 \end{matrix} \\ \begin{pmatrix} 1 \\ \frac{1}{3} \\ 0 \\ 0 \\ 0 \end{pmatrix} & \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{3} & \frac{1}{3} & 0 \\ 0 & \frac{2}{3} & 0 & 0 & \frac{1}{3} \\ 0 & 0 & 0 & \frac{2}{3} & \frac{1}{3} \\ 0 & 0 & 0 & \frac{2}{3} & \frac{1}{3} \end{pmatrix} \end{matrix}$$




The limiting distribution depends on the initial state:

$$\delta_0 P^t \rightarrow \pi_{\{0\}} \quad \text{but} \quad \delta_3 P^t \rightarrow \pi_{\{3,4\}}$$

A chain starting at 1 gets absorbed to $\{0\}$ or $\{3,4\}$.

Example 6

$$P = \begin{matrix} & \begin{matrix} 0 & 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ \frac{1}{3} \\ 0 \\ 0 \\ 0 \end{matrix} & \begin{pmatrix} 0 & 1 & 2 & 3 & 4 \\ 1 & 0 & 0 & 0 & 0 \\ \frac{1}{3} & 0 & \frac{1}{3} & \frac{1}{3} & 0 \\ 0 & \frac{2}{3} & 0 & 0 & \frac{1}{3} \\ 0 & 0 & 0 & \frac{2}{3} & \frac{1}{3} \\ 0 & 0 & 0 & \frac{2}{3} & \frac{1}{3} \end{pmatrix} \end{matrix}$$



The limiting distribution depends on the initial state:

$$\delta_0 P^t \rightarrow \pi_{\{0\}} \quad \text{but} \quad \delta_3 P^t \rightarrow \pi_{\{3,4\}}$$

A chain starting at 1 gets absorbed to $\{0\}$ or $\{3,4\}$. We analyze this more next time.



Pierre Brémaud.

Markov Chains.

Springer, 1999.



Richard Durrett.

Essentials of Stochastic Processes.

Springer, second edition, 2012.



Vidyadhar G. Kulkarni.

Introduction to Modeling and Analysis of Stochastic Systems.

Springer, second edition, 2011.



David A. Levin, Yuval Peres, and Elizabeth L. Wilmer.

Markov Chains and Mixing Times.

American Mathematical Society,

<http://pages.uoregon.edu/dlevin/MARKOV/>, 2008.



James R. Norris.

Markov Chains.

Cambridge University Press, 1997.

References

Images used in the slides (in order of appearance)

1. Image courtesy of think4photop at FreeDigitalPhotos.net