

MS-C2111 Stochastic processes



Lecture 4

Passage times and hitting probabilities

Jukka Kohonen
Aalto University

Contents

Passage times

Hitting probabilities

Gambler's ruin

Example: HR management

Kalvonvääntäjät Oyj is a management consulting company with three types of employees: 1 = junior, 2 = senior, 3 = partner

Example: HR management

Kalvonvääntäjät Oyj is a management consulting company with three types of employees: 1 = junior, 2 = senior, 3 = partner

During each month:

Example: HR management

Kalvonvääntäjät Oyj is a management consulting company with three types of employees: 1 = junior, 2 = senior, 3 = partner

During each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020

Example: HR management

Kalvonvääntäjät Oyj is a management consulting company with three types of employees: 1 = junior, 2 = senior, 3 = partner

During each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008

Example: HR management

Kalvonvääntäjät Oyj is a management consulting company with three types of employees: 1 = junior, 2 = senior, 3 = partner

During each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008
- Partner exits w.pr. 0.010

Example: HR management

Kalvonvääntäjät Oyj is a management consulting company with three types of employees: 1 = junior, 2 = senior, 3 = partner

During each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008
- Partner exits w.pr. 0.010

What is the expected duration that a newly recruited employee remains in the company?

Example: HR management

Kalvonväentäjät Oyj is a management consulting company with three types of employees: **1** = junior, **2** = senior, **3** = partner

During each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008
- Partner exits w.pr. 0.010

What is the expected duration that a newly recruited employee remains in the company?

How long is a freshly promoted partner expected to serve in the company?

Passage time

Passage time of (X_0, X_1, \dots) into A

$$T_A = \min\{t \geq 0 : X_t \in A\} \quad (\min \emptyset = \infty)$$

Passage time

Passage time of (X_0, X_1, \dots) into A

$$T_A = \min\{t \geq 0 : X_t \in A\} \quad (\min \emptyset = \infty)$$

Random variable with state space $T_A \in$

Passage time

Passage time of (X_0, X_1, \dots) into A

$$T_A = \min\{t \geq 0 : X_t \in A\} \quad (\min \emptyset = \infty)$$

Random variable with state space $T_A \in \{0, 1, 2, \dots, \infty\}$

Passage time

Passage time of (X_0, X_1, \dots) into A

$$T_A = \min\{t \geq 0 : X_t \in A\} \quad (\min \emptyset = \infty)$$

Random variable with state space $T_A \in \{0, 1, 2, \dots, \infty\}$

$$x + \infty =$$

Passage time

Passage time of (X_0, X_1, \dots) into A

$$T_A = \min\{t \geq 0 : X_t \in A\} \quad (\min \emptyset = \infty)$$

Random variable with state space $T_A \in \{0, 1, 2, \dots, \infty\}$

$$x + \infty = \infty$$

Passage time

Passage time of (X_0, X_1, \dots) into A

$$T_A = \min\{t \geq 0 : X_t \in A\} \quad (\min \emptyset = \infty)$$

Random variable with state space $T_A \in \{0, 1, 2, \dots, \infty\}$

$$x + \infty = \infty$$

$$x \cdot \infty =$$

Passage time

Passage time of (X_0, X_1, \dots) into A

$$T_A = \min\{t \geq 0 : X_t \in A\} \quad (\min \emptyset = \infty)$$

Random variable with state space $T_A \in \{0, 1, 2, \dots, \infty\}$

$$x + \infty = \infty$$

$$x \cdot \infty = \begin{cases} 0, & x = 0 \\ \infty, & x > 0 \end{cases}$$

Passage time

Passage time of (X_0, X_1, \dots) into A

$$T_A = \min\{t \geq 0 : X_t \in A\} \quad (\min \emptyset = \infty)$$

Random variable with state space $T_A \in \{0, 1, 2, \dots, \infty\}$

$$x + \infty = \infty$$

$$x \cdot \infty = \begin{cases} 0, & x = 0 \\ \infty, & x > 0 \end{cases}$$

Expected passage time into A

$$k_A(x) = \mathbb{E}(T_A | X_0 = x)$$

Passage time

Passage time of (X_0, X_1, \dots) into A

$$T_A = \min\{t \geq 0 : X_t \in A\} \quad (\min \emptyset = \infty)$$

Random variable with state space $T_A \in \{0, 1, 2, \dots, \infty\}$

$$x + \infty = \infty$$

$$x \cdot \infty = \begin{cases} 0, & x = 0 \\ \infty, & x > 0 \end{cases}$$

Expected passage time into A

$$k_A(x) = \mathbb{E}(T_A | X_0 = x)$$

Deterministic number in $k_A(x) \in$

Passage time

Passage time of (X_0, X_1, \dots) into A

$$T_A = \min\{t \geq 0 : X_t \in A\} \quad (\min \emptyset = \infty)$$

Random variable with state space $T_A \in \{0, 1, 2, \dots, \infty\}$

$$x + \infty = \infty$$

$$x \cdot \infty = \begin{cases} 0, & x = 0 \\ \infty, & x > 0 \end{cases}$$

Expected passage time into A

$$k_A(x) = \mathbb{E}(T_A | X_0 = x)$$

Deterministic number in $k_A(x) \in \mathbb{R}_+ \cup \{\infty\}$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A | X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A \mid X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y)f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A \mid X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y)f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A \mid X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y)f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A \implies T_A = 0$ surely

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A \mid X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y)f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A \implies T_A = 0$ surely $\implies k_A(x) =$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A \mid X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y)f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A \implies T_A = 0$ surely $\implies k_A(x) = 0$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A \mid X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y)f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A \implies T_A = 0$ surely $\implies k_A(x) = 0$

(ii) $x \notin A$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A | X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y)f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A \implies T_A = 0$ surely $\implies k_A(x) = 0$

(ii) $x \notin A \implies$

$$k_A(x) = \mathbb{E}(T_A | X_0 = x)$$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A | X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A \implies T_A = 0$ surely $\implies k_A(x) = 0$

(ii) $x \notin A \implies$

$$k_A(x) = \mathbb{E}(T_A | X_0 = x) = \sum_y P(x, y) \mathbb{E}(T_A | X_1 = y, X_0 = x)$$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A | X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A \implies T_A = 0$ surely $\implies k_A(x) = 0$

(ii) $x \notin A \implies$

$$k_A(x) = \mathbb{E}(T_A | X_0 = x) = \sum_y P(x, y) \mathbb{E}(T_A | X_1 = y, X_0 = x)$$

where $\mathbb{E}(T_A | X_1 = y, X_0 = x)$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A | X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A \implies T_A = 0$ surely $\implies k_A(x) = 0$

(ii) $x \notin A \implies$

$$k_A(x) = \mathbb{E}(T_A | X_0 = x) = \sum_y P(x, y) \mathbb{E}(T_A | X_1 = y, X_0 = x)$$

where $\mathbb{E}(T_A | X_1 = y, X_0 = x) = 1 + \mathbb{E}(T_A | X_0 = y)$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A | X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A \implies T_A = 0$ surely $\implies k_A(x) = 0$

(ii) $x \notin A \implies$

$$k_A(x) = \mathbb{E}(T_A | X_0 = x) = \sum_y P(x, y) \mathbb{E}(T_A | X_1 = y, X_0 = x)$$

where $\mathbb{E}(T_A | X_1 = y, X_0 = x) = 1 + \mathbb{E}(T_A | X_0 = y) = 1 + k_A(y)$

Computing the expected passage times

Theorem: The expected passage times $k_A(x) = \mathbb{E}(T_A | X_0 = x)$ form the smallest nonnegative solution to

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Proof (that k_A is a solution)

(i) $x \in A \implies T_A = 0$ surely $\implies k_A(x) = 0$

(ii) $x \notin A \implies$

$$k_A(x) = \mathbb{E}(T_A | X_0 = x) = \sum_y P(x, y) \mathbb{E}(T_A | X_1 = y, X_0 = x)$$

$$\text{where } \mathbb{E}(T_A | X_1 = y, X_0 = x) = 1 + \mathbb{E}(T_A | X_0 = y) = 1 + k_A(y)$$

$$\begin{aligned} k_A(x) &= \sum_y P(x, y)(1 + k_A(y)) = \sum_y P(x, y) + \sum_y P(x, y)k_A(y) \\ &= 1 + \sum_{y \in A^c} P(x, y)k_A(y). \end{aligned}$$

Computing the expected passage times

How to find the smallest nonnegative solution k_A for

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Computing the expected passage times

How to find the smallest nonnegative solution k_A for

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Poisson equation

$$Df(x) = -1, \quad x \in B,$$

$$f(x) = 0, \quad x \in \partial B,$$

$$D = P - I, \quad B = A^c, \quad \partial B = A.$$

Computing the expected passage times

How to find the smallest nonnegative solution k_A for

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Poisson equation

Recursive solution method: $f_0(x) = 0$,

$$Df(x) = -1, \quad x \in B,$$

$$f(x) = 0, \quad x \in \partial B,$$

$$D = P - I, \quad B = A^c, \quad \partial B = A.$$

Computing the expected passage times

How to find the smallest nonnegative solution k_A for

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Poisson equation

$$\begin{aligned} Df(x) &= -1, & x \in B, \\ f(x) &= 0, & x \in \partial B, \end{aligned}$$

Recursive solution method: $f_0(x) = 0$,

$$f_{n+1}(x) = \begin{cases} 1 + \sum_{y \notin A} P(x, y) f_n(y), & x \notin A \\ 0, & x \in A \end{cases}$$

$$D = P - I, \quad B = A^c, \quad \partial B = A.$$

Computing the expected passage times

How to find the smallest nonnegative solution k_A for

$$f(x) = 1 + \sum_{y \notin A} P(x, y) f(y), \quad x \notin A,$$

$$f(x) = 0, \quad x \in A.$$

Poisson equation

$$\begin{aligned} Df(x) &= -1, & x \in B, \\ f(x) &= 0, & x \in \partial B, \end{aligned}$$

Recursive solution method: $f_0(x) = 0$,

$$f_{n+1}(x) = \begin{cases} 1 + \sum_{y \notin A} P(x, y) f_n(y), & x \notin A \\ 0, & x \in A \end{cases}$$

$$D = P - I, \quad B = A^c, \quad \partial B = A.$$

$$f_n(x) \rightarrow k_A(x) \text{ when } n \rightarrow \infty$$

Example: HR management

1 = junior, 2 = senior, 3 = partner, during each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008
- Partner exits w.pr. 0.010

Q: How long does a freshly recruited employee stay in company?

Example: HR management

1 = junior, 2 = senior, 3 = partner, during each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008
- Partner exits w.pr. 0.010

Q: How long does a freshly recruited employee stay in company?

X_t = Employee status after t months

Example: HR management

1 = junior, 2 = senior, 3 = partner, during each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008
- Partner exits w.pr. 0.010

Q: How long does a freshly recruited employee stay in company?

X_t = Employee status after t months

Markov chain?

Example: HR management

1 = junior, 2 = senior, 3 = partner, during each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008
- Partner exits w.pr. 0.010

Q: How long does a freshly recruited employee stay in company?

X_t = Employee status after t months

Markov chain? **YES** for $S = \{0, 1, 2, 3\}$ with extra state **0** = exit

Example: HR management

1 = junior, 2 = senior, 3 = partner, during each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008
- Partner exits w.pr. 0.010

Q: How long does a freshly recruited employee stay in company?

X_t = Employee status after t months

Markov chain? **YES** for $S = \{0, 1, 2, 3\}$ with extra state 0 = exit

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix}$$

Example: HR management

1 = junior, 2 = senior, 3 = partner, during each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008
- Partner exits w.pr. 0.010

Q: How long does a freshly recruited employee stay in company?

X_t = Employee status after t months

Markov chain? **YES** for $S = \{0, 1, 2, 3\}$ with extra state 0 = exit

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix}$$

A: Expected passage time $k_A(1)$ from state 1 into $A =$

Example: HR management

1 = junior, 2 = senior, 3 = partner, during each month:

- Junior promoted to senior w.pr. 0.030, exits w.pr. 0.020
- Senior promoted to partner w.pr. 0.010, exits w.pr. 0.008
- Partner exits w.pr. 0.010

Q: How long does a freshly recruited employee stay in company?

X_t = Employee status after t months

Markov chain? **YES** for $S = \{0, 1, 2, 3\}$ with extra state 0 = exit

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix}$$

A: Expected passage time $k_A(1)$ from state 1 into $A = \{0\}$

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x, y)f(y)$, $x = 1, 2, 3$

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x,y)f(y)$, $x = 1, 2, 3$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix}$$

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x, y)f(y)$, $x = 1, 2, 3$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix}$$

$$f(1) = 1 + 0.950 f(1) + 0.030 f(2)$$

$$f(2) = 1 + 0.982 f(2) + 0.010 f(3)$$

$$f(3) = 1 + 0.990 f(3).$$

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x, y)f(y)$, $x = 1, 2, 3$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(1) &= 1 + 0.950 f(1) + 0.030 f(2) \\ f(2) &= 1 + 0.982 f(2) + 0.010 f(3) \\ f(3) &= 1 + 0.990 f(3). \end{aligned}$$
$$f(3) = \frac{1}{1 - 0.990} = 100$$

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x, y)f(y)$, $x = 1, 2, 3$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(1) &= 1 + 0.950 f(1) + 0.030 f(2) \\ f(2) &= 1 + 0.982 f(2) + 0.010 f(3) \\ f(3) &= 1 + 0.990 f(3). \end{aligned}$$

$$f(3) = \frac{1}{1 - 0.990} = 100$$

$$f(2) = \frac{1 + 0.010 f(3)}{1 - 0.982} = 111.11$$

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x, y)f(y)$, $x = 1, 2, 3$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(1) &= 1 + 0.950 f(1) + 0.030 f(2) \\ f(2) &= 1 + 0.982 f(2) + 0.010 f(3) \\ f(3) &= 1 + 0.990 f(3). \end{aligned}$$

$$f(3) = \frac{1}{1 - 0.990} = 100$$

$$f(2) = \frac{1 + 0.010 f(3)}{1 - 0.982} = 111.11$$

$$f(1) = \frac{1 + 0.030 f(2)}{1 - 0.950} = 86.67$$

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x, y)f(y)$, $x = 1, 2, 3$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(1) &= 1 + 0.950 f(1) + 0.030 f(2) \\ f(2) &= 1 + 0.982 f(2) + 0.010 f(3) \\ f(3) &= 1 + 0.990 f(3). \end{aligned}$$

$$f(3) = \frac{1}{1 - 0.990} = 100$$

$$f(2) = \frac{1 + 0.010 f(3)}{1 - 0.982} = 111.11$$

$$f(1) = \frac{1 + 0.030 f(2)}{1 - 0.950} = 86.67$$

Minimal solution?

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x, y)f(y)$, $x = 1, 2, 3$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(1) &= 1 + 0.950 f(1) + 0.030 f(2) \\ f(2) &= 1 + 0.982 f(2) + 0.010 f(3) \\ f(3) &= 1 + 0.990 f(3). \end{aligned}$$

$$f(3) = \frac{1}{1 - 0.990} = 100$$

$$f(2) = \frac{1 + 0.010 f(3)}{1 - 0.982} = 111.11$$

$$f(1) = \frac{1 + 0.030 f(2)}{1 - 0.950} = 86.67$$

Minimal solution? YES because this is the only nonnegative solution.

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x, y)f(y)$, $x = 1, 2, 3$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(1) &= 1 + 0.950 f(1) + 0.030 f(2) \\ f(2) &= 1 + 0.982 f(2) + 0.010 f(3) \\ f(3) &= 1 + 0.990 f(3). \end{aligned}$$

$$f(3) = \frac{1}{1 - 0.990} = 100$$

$$f(2) = \frac{1 + 0.010 f(3)}{1 - 0.982} = 111.11$$

$$f(1) = \frac{1 + 0.030 f(2)}{1 - 0.950} = 86.67$$

Minimal solution? YES because this is the only nonnegative solution.

$$\implies (k_A(0), k_A(1), k_A(2), k_A(3)) = (0, 86.67, 111.11, 100)$$

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x, y)f(y)$, $x = 1, 2, 3$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(1) &= 1 + 0.950 f(1) + 0.030 f(2) \\ f(2) &= 1 + 0.982 f(2) + 0.010 f(3) \\ f(3) &= 1 + 0.990 f(3). \end{aligned}$$

$$f(3) = \frac{1}{1 - 0.990} = 100$$

$$f(2) = \frac{1 + 0.010 f(3)}{1 - 0.982} = 111.11$$

$$f(1) = \frac{1 + 0.030 f(2)}{1 - 0.950} = 86.67$$

Minimal solution? YES because this is the only nonnegative solution.

$$\implies (k_A(0), k_A(1), k_A(2), k_A(3)) = (0, 86.67, 111.11, 100)$$

Freshly recruited employee is expected to stay for 86.67 mo ≈ 7.2 y

Example: HR management

Let us solve $f(0) = 0$ and $f(x) = 1 + \sum_{y=1}^3 P(x, y)f(y)$, $x = 1, 2, 3$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(1) &= 1 + 0.950 f(1) + 0.030 f(2) \\ f(2) &= 1 + 0.982 f(2) + 0.010 f(3) \\ f(3) &= 1 + 0.990 f(3). \end{aligned}$$
$$f(3) = \frac{1}{1 - 0.990} = 100$$

$$f(2) = \frac{1 + 0.010 f(3)}{1 - 0.982} = 111.11$$

$$f(1) = \frac{1 + 0.030 f(2)}{1 - 0.950} = 86.67$$

Minimal solution? YES because this is the only nonnegative solution.

$$\implies (k_A(0), k_A(1), k_A(2), k_A(3)) = (0, 86.67, 111.11, 100)$$

Freshly recruited employee is expected to stay for 86.67 mo ≈ 7.2 y

Freshly promoted partner is expected to stay for 100 mo ≈ 8.3 y

Contents

Passage times

Hitting probabilities

Gambler's ruin

Example: HR management

What is the probability that a freshly recruited junior eventually gets promoted to a partner?

Hitting probabilities

The probability that a process starting at x visits a set A is denoted by

$$h_A(x) = \mathbb{P}(X_t \in A \text{ for some } t \geq 0 \mid X_0 = x).$$

Hitting probabilities

The probability that a process starting at x visits a set A is denoted by

$$h_A(x) = \mathbb{P}(X_t \in A \text{ for some } t \geq 0 \mid X_0 = x).$$

In practice, to solve one hitting probability, we need to solve them all.

Computing hitting probabilities

Theorem

The hitting probabilities $h_A = (h_A(x) : x \in S)$ form the smallest nonnegative solution to

$$\begin{aligned} f(x) &= \sum_{y \in S} P(x, y) f(y), & x \notin A, \\ f(x) &= 1, & x \in A, \end{aligned}$$

Computing hitting probabilities

Theorem

The hitting probabilities $h_A = (h_A(x) : x \in S)$ form the smallest nonnegative solution to

$$\begin{aligned} f(x) &= \sum_{y \in S} P(x, y) f(y), & x \notin A, \\ f(x) &= 1, & x \in A, \end{aligned}$$

Poisson equation

$$\begin{aligned} Df(x) &= 0, & x \in B, \\ f(x) &= 1, & x \in \partial B, \end{aligned}$$

Proof: Lecture notes.

$$D = P - I, \quad B = A^c, \quad \partial B = A.$$

Example: HR management

The probability that a freshly recruited junior eventually gets promoted to a partner is $h_A(1)$, $A = \{3\}$.

Example: HR management

The probability that a freshly recruited junior eventually gets promoted to a partner is $h_A(1)$, $A = \{3\}$.

Let us solve $f(3) = 1$ and $f(x) = \sum_{y=0}^3 P(x, y)f(y)$, $x = 0, 1, 2$

Example: HR management

The probability that a freshly recruited junior eventually gets promoted to a partner is $h_A(1)$, $A = \{3\}$.

Let us solve $f(3) = 1$ and $f(x) = \sum_{y=0}^3 P(x, y)f(y)$, $x = 0, 1, 2$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix}$$

$$f(0) = f(0),$$

$$f(1) = 0.020 f(0) + 0.950 f(1) + 0.030 f(2),$$

$$f(2) = 0.008 f(0) + 0.982 f(2) + 0.010 f(3),$$

$$f(3) = 1.$$

Example: HR management

The probability that a freshly recruited junior eventually gets promoted to a partner is $h_A(1)$, $A = \{3\}$.

Let us solve $f(3) = 1$ and $f(x) = \sum_{y=0}^3 P(x, y)f(y)$, $x = 0, 1, 2$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix}$$

$$f(0) = f(0),$$

$$f(1) = 0.020 f(0) + 0.950 f(1) + 0.030 f(2),$$

$$f(2) = 0.008 f(0) + 0.982 f(2) + 0.010 f(3),$$

$$f(3) = 1.$$

$$0 \not\rightarrow 3 \implies f(0) = 0.$$

Example: HR management

The probability that a freshly recruited junior eventually gets promoted to a partner is $h_A(1)$, $A = \{3\}$.

Let us solve $f(3) = 1$ and $f(x) = \sum_{y=0}^3 P(x, y)f(y)$, $x = 0, 1, 2$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(0) &= f(0), \\ f(1) &= 0.020 f(0) + 0.950 f(1) + 0.030 f(2), \\ f(2) &= 0.008 f(0) + 0.982 f(2) + 0.010 f(3), \\ f(3) &= 1. \end{aligned}$$

$$0 \not\rightarrow 3 \implies f(0) = 0.$$

$$\implies f = [0, 0.333, 0.556, 1] \text{ is a solution.}$$

Example: HR management

The probability that a freshly recruited junior eventually gets promoted to a partner is $h_A(1)$, $A = \{3\}$.

Let us solve $f(3) = 1$ and $f(x) = \sum_{y=0}^3 P(x, y)f(y)$, $x = 0, 1, 2$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(0) &= f(0), \\ f(1) &= 0.020 f(0) + 0.950 f(1) + 0.030 f(2), \\ f(2) &= 0.008 f(0) + 0.982 f(2) + 0.010 f(3), \\ f(3) &= 1. \end{aligned}$$

$$0 \not\rightarrow 3 \implies f(0) = 0.$$

$$\implies f = [0, 0.333, 0.556, 1] \text{ is a solution.}$$

Smallest nonnegative?

Example: HR management

The probability that a freshly recruited junior eventually gets promoted to a partner is $h_A(1)$, $A = \{3\}$.

Let us solve $f(3) = 1$ and $f(x) = \sum_{y=0}^3 P(x, y)f(y)$, $x = 0, 1, 2$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(0) &= f(0), \\ f(1) &= 0.020 f(0) + 0.950 f(1) + 0.030 f(2), \\ f(2) &= 0.008 f(0) + 0.982 f(2) + 0.010 f(3), \\ f(3) &= 1. \end{aligned}$$

$$0 \not\rightarrow 3 \implies f(0) = 0.$$

$$\implies f = [0, 0.333, 0.556, 1] \text{ is a solution.}$$

Smallest nonnegative? Yes.

Example: HR management

The probability that a freshly recruited junior eventually gets promoted to a partner is $h_A(1)$, $A = \{3\}$.

Let us solve $f(3) = 1$ and $f(x) = \sum_{y=0}^3 P(x, y)f(y)$, $x = 0, 1, 2$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(0) &= f(0), \\ f(1) &= 0.020 f(0) + 0.950 f(1) + 0.030 f(2), \\ f(2) &= 0.008 f(0) + 0.982 f(2) + 0.010 f(3), \\ f(3) &= 1. \end{aligned}$$

$$0 \not\rightarrow 3 \implies f(0) = 0.$$

$$\implies f = [0, 0.333, 0.556, 1] \text{ is a solution.}$$

Smallest nonnegative? Yes.

$$\implies [h_A(0), h_A(1), h_A(2), h_A(3)] = [0, 0.333, 0.556, 1]$$

Example: HR management

The probability that a freshly recruited junior eventually gets promoted to a partner is $h_A(1)$, $A = \{3\}$.

Let us solve $f(3) = 1$ and $f(x) = \sum_{y=0}^3 P(x, y)f(y)$, $x = 0, 1, 2$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(0) &= f(0), \\ f(1) &= 0.020 f(0) + 0.950 f(1) + 0.030 f(2), \\ f(2) &= 0.008 f(0) + 0.982 f(2) + 0.010 f(3), \\ f(3) &= 1. \end{aligned}$$

$$0 \not\rightarrow 3 \implies f(0) = 0.$$

$$\implies f = [0, 0.333, 0.556, 1] \text{ is a solution.}$$

Smallest nonnegative? Yes.

$$\implies [h_A(0), h_A(1), h_A(2), h_A(3)] = [0, 0.333, 0.556, 1]$$

\implies New junior becomes a partner with probability

$$f(1) = h_A(1) = 0.333$$

Example: HR management

The probability that a freshly recruited junior eventually gets promoted to a partner is $h_A(1)$, $A = \{3\}$.

Let us solve $f(3) = 1$ and $f(x) = \sum_{y=0}^3 P(x, y)f(y)$, $x = 0, 1, 2$

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.020 & 0.950 & 0.030 & 0 \\ 0.008 & 0 & 0.982 & 0.010 \\ 0.010 & 0 & 0 & 0.990 \end{bmatrix} \quad \begin{aligned} f(0) &= f(0), \\ f(1) &= 0.020 f(0) + 0.950 f(1) + 0.030 f(2), \\ f(2) &= 0.008 f(0) + 0.982 f(2) + 0.010 f(3), \\ f(3) &= 1. \end{aligned}$$

$$0 \not\rightarrow 3 \implies f(0) = 0.$$

$$\implies f = [0, 0.333, 0.556, 1] \text{ is a solution.}$$

Smallest nonnegative? Yes.

$$\implies [h_A(0), h_A(1), h_A(2), h_A(3)] = [0, 0.333, 0.556, 1]$$

\implies New junior becomes a partner with probability

$$f(1) = h_A(1) = 0.333$$

$$\sum_x f(x) \neq 1 \text{ (why?)}$$

Contents

Passage times

Hitting probabilities

Gambler's ruin

Gambler's ruin

X_t = Wealth of a gambler after t rounds

- Win 1 EUR w.pr. q
- Lose 1 EUR otherwise

Gambler's ruin

X_t = Wealth of a gambler after t rounds

- Win 1 EUR w.pr. q
- Lose 1 EUR otherwise


Game stops when the gambler's wealth hits M (gambler's target) or 0 (gambler's money is all gone).

Gambler's ruin

X_t = Wealth of a gambler after t rounds

- Win 1 EUR w.pr. q
- Lose 1 EUR otherwise

Game stops when the gambler's wealth hits M (gambler's target) or 0 (gambler's money is all gone).




Gambler's ruin

X_t = Wealth of a gambler after t rounds

- Win 1 EUR w.pr. q
- Lose 1 EUR otherwise

Game stops when the gambler's wealth hits M (gambler's target) or 0 (gambler's money is all gone).



Success probability


$$h(x) = \mathbb{P}(X_t = M \text{ for some } t \geq 0 \mid X_0 = x).$$

Gambler's ruin

X_t = Wealth of a gambler after t rounds

- Win 1 EUR w.pr. q
- Lose 1 EUR otherwise

Game stops when the gambler's wealth hits M (gambler's target) or 0 (gambler's money is all gone).



Success probability

$$h(x) = \mathbb{P}(X_t = M \text{ for some } t \geq 0 \mid X_0 = x).$$

Ruin probability = $1 - h(x)$.

Gambler's ruin

The success probability $h(x) = h_A(x)$ with $A = \{M\}$.

$\implies h(x)$ is the minimal nonnegative solution of

$$h(0) = h(0),$$

$$h(x) = (1 - q)h(x - 1) + qh(x + 1), \quad 0 < x < M,$$

$$h(M) = 1.$$

Gambler's ruin

The success probability $h(x) = h_A(x)$ with $A = \{M\}$.

$\implies h(x)$ is the minimal nonnegative solution of

$$h(0) = h(0),$$

$$h(x) = (1 - q)h(x - 1) + qh(x + 1), \quad 0 < x < M,$$

$$h(M) = 1.$$

Clearly $h(0) = 0$.

Gambler's ruin

The success probability $h(x) = h_A(x)$ with $A = \{M\}$.

$\implies h(x)$ is the minimal nonnegative solution of

$$h(0) = 0,$$

$$h(x) = (1 - q)h(x - 1) + qh(x + 1), \quad 0 < x < M,$$

$$h(M) = 1.$$

Clearly $h(0) = 0$.

We need to find the minimal nonnegative solution to

$$h(x) = (1 - q)h(x - 1) + qh(x + 1), \quad 0 < x < M$$

with boundary conditions $h(0) = 0$ and $h(M) = 1$.

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (yrite): $h(x) = z^x$ for some z

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (yrite): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (yrite): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

$$\implies qz^2 - z + (1 - q) = 0$$

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (yrite): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

$$\implies qz^2 - z + (1 - q) = 0$$

which has two distinct roots

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (yrite): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

$$\implies qz^2 - z + (1 - q) = 0$$

which has two distinct roots $\alpha = \frac{1-q}{q}$ and $\beta = 1$.

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (yrite): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

$$\implies qz^2 - z + (1 - q) = 0$$

which has two distinct roots $\alpha = \frac{1-q}{q}$ and $\beta = 1$.

\implies All solutions are of the form

$$h(x) = c\alpha^x + d\beta^x$$

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (write): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

$$\implies qz^2 - z + (1 - q) = 0$$

which has two distinct roots $\alpha = \frac{1-q}{q}$ and $\beta = 1$.

\implies All solutions are of the form

$$h(x) = c\alpha^x + d\beta^x$$

The boundary conditions $h(0) = 0$ and $h(M) = 1$ now become

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (yrite): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

$$\implies qz^2 - z + (1 - q) = 0$$

which has two distinct roots $\alpha = \frac{1-q}{q}$ and $\beta = 1$.

\implies All solutions are of the form

$$h(x) = c\alpha^x + d\beta^x$$

The boundary conditions $h(0) = 0$ and $h(M) = 1$ now become

$$c + d = 0,$$

$$c\alpha^M + d = 1,$$

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (yrite): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

$$\implies qz^2 - z + (1 - q) = 0$$

which has two distinct roots $\alpha = \frac{1-q}{q}$ and $\beta = 1$.

\implies All solutions are of the form

$$h(x) = c\alpha^x + d\beta^x$$

The boundary conditions $h(0) = 0$ and $h(M) = 1$ now become

$$c + d = 0,$$

$$c\alpha^M + d = 1,$$

from which we solve $d =$

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (write): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

$$\implies qz^2 - z + (1 - q) = 0$$

which has two distinct roots $\alpha = \frac{1-q}{q}$ and $\beta = 1$.

\implies All solutions are of the form

$$h(x) = c\alpha^x + d\beta^x$$

The boundary conditions $h(0) = 0$ and $h(M) = 1$ now become

$$c + d = 0,$$

$$c\alpha^M + d = 1,$$

from which we solve $d = -c$ and $c =$

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (yrite): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

$$\implies qz^2 - z + (1 - q) = 0$$

which has two distinct roots $\alpha = \frac{1-q}{q}$ and $\beta = 1$.

\implies All solutions are of the form

$$h(x) = c\alpha^x + d\beta^x$$

The boundary conditions $h(0) = 0$ and $h(M) = 1$ now become

$$c + d = 0,$$

$$c\alpha^M + d = 1,$$

from which we solve $d = -c$ and $c = 1/(\alpha^M - 1)$,

Asymmetric case: $q \neq \frac{1}{2}$

Ansatz (yrite): $h(x) = z^x$ for some z

$$\implies z^x = (1 - q)z^{x-1} + qz^{x+1}$$

$$\implies qz^2 - z + (1 - q) = 0$$

which has two distinct roots $\alpha = \frac{1-q}{q}$ and $\beta = 1$.

\implies All solutions are of the form

$$h(x) = c\alpha^x + d\beta^x$$

The boundary conditions $h(0) = 0$ and $h(M) = 1$ now become

$$c + d = 0,$$

$$c\alpha^M + d = 1,$$

from which we solve $d = -c$ and $c = 1/(\alpha^M - 1)$, and obtain

$$h(x) = \frac{\alpha^x - 1}{\alpha^M - 1}.$$

Symmetric case: $q = \frac{1}{2}$

Let us study how

$$h(x) = \frac{\alpha^x - 1}{\alpha^M - 1}, \quad \alpha = \frac{1 - q}{q}$$

behaves as $q \rightarrow \frac{1}{2}$.

Symmetric case: $q = \frac{1}{2}$

Let us study how

$$h(x) = \frac{\alpha^x - 1}{\alpha^M - 1}, \quad \alpha = \frac{1 - q}{q}$$

behaves as $q \rightarrow \frac{1}{2}$.

Now

$$\lim_{\alpha \rightarrow 1} h(x) =$$

Symmetric case: $q = \frac{1}{2}$

Let us study how

$$h(x) = \frac{\alpha^x - 1}{\alpha^M - 1}, \quad \alpha = \frac{1 - q}{q}$$

behaves as $q \rightarrow \frac{1}{2}$.

Now

$$\lim_{\alpha \rightarrow 1} h(x) = \lim_{\alpha \rightarrow 1} \frac{\alpha^x - 1}{\alpha^M - 1} =$$

Symmetric case: $q = \frac{1}{2}$

Let us study how

$$h(x) = \frac{\alpha^x - 1}{\alpha^M - 1}, \quad \alpha = \frac{1 - q}{q}$$

behaves as $q \rightarrow \frac{1}{2}$.

Now

$$\lim_{\alpha \rightarrow 1} h(x) = \lim_{\alpha \rightarrow 1} \frac{\alpha^x - 1}{\alpha^M - 1} = \lim_{\alpha \rightarrow 1} \frac{x\alpha^{x-1}}{M\alpha^{M-1}} =$$

Symmetric case: $q = \frac{1}{2}$

Let us study how

$$h(x) = \frac{\alpha^x - 1}{\alpha^M - 1}, \quad \alpha = \frac{1-q}{q}$$

behaves as $q \rightarrow \frac{1}{2}$.

Now

$$\lim_{\alpha \rightarrow 1} h(x) = \lim_{\alpha \rightarrow 1} \frac{\alpha^x - 1}{\alpha^M - 1} = \lim_{\alpha \rightarrow 1} \frac{x\alpha^{x-1}}{M\alpha^{M-1}} = \frac{x}{M}$$


by l'Hôpital's rule.

Gambler's ruin — Main theorem

Theorem

The probability that the random walk started at x eventually hits M equals

$$h(x) = \begin{cases} \frac{\left(\frac{1-q}{q}\right)^x - 1}{\left(\frac{1-q}{q}\right)^M - 1}, & q \neq \frac{1}{2}, \\ \frac{x}{M}, & q = \frac{1}{2}. \end{cases}$$




Gambler's ruin — Main theorem

Theorem

The probability that the random walk started at x eventually hits M equals

$$h(x) = \begin{cases} \frac{\left(\frac{1-q}{q}\right)^x - 1}{\left(\frac{1-q}{q}\right)^M - 1}, & q \neq \frac{1}{2}, \\ \frac{x}{M}, & q = \frac{1}{2}. \end{cases}$$



Main message: When $q \leq \frac{1}{2}$, the probability of ever reaching a state M from an initial state x tends to zero as $M \rightarrow \infty$.

Example: Roulette

In a game of roulette where a bet of 1 EUR is placed on the ball falling into one of 18 red pockets out of 37 pockets, the probability of winning 1 EUR is $q = \frac{18}{37}$ and the probability of losing 1 EUR is $1 - q$. If a gambler targets to double his initial wealth x , then the probability $h(x)$ of successfully ending the game is obtained by applying the previous theorem with $M = 2x$.

Example: Roulette

In a game of roulette where a bet of 1 EUR is placed on the ball falling into one of 18 red pockets out of 37 pockets, the probability of winning 1 EUR is $q = \frac{18}{37}$ and the probability of losing 1 EUR is $1 - q$. If a gambler targets to double his initial wealth x , then the probability $h(x)$ of successfully ending the game is obtained by applying the previous theorem with $M = 2x$.

Initial wealth (EUR)	1	5	10	20	50
Success probability	0.4865	0.4328	0.3680	0.2533	0.0628

Table: Probability of successfully doubling the initial wealth in a game of roulette by betting 1 EUR on red.

Aineistolähteet

Esityksessä käytetyt kuvat (esiintymisjärjestyksessä)

1. Image courtesy of think4photop at FreeDigitalPhotos.net