Email training, N4 October 2-8

Problem 4.1. Let a, b are solutions of equation $x^2 + 2x - 13 = 0$. Construct an equation which roots are a + 1, b + 1.

Solution 4.1. Since a, b are solutions of the equation $x^2 + 2x - 13 = 0$ then according to Viet's theorem one has

$$\begin{cases} a+b=-2\\ ab=-13 \end{cases}.$$

If a+1, b+1 and c+1 are solutions of the equation $x^2+px+r=0$ then, according to Viet's theorem one has

$$\begin{cases} p = -(a+1+b+1) = -(-2+2) = 0 \\ r = (a+1)(b+1) = ab+a+b+1 = -14 \end{cases}$$

So one gets the equation $x^2 - 14 = 0$.

Solution 2. Note that if the roots of the equation P(x) = 0 are a, b then the roots of the equation P(x-1) = 0 will be a+1, b+1. It remains to conclude that

$$P(x-1) = (x-1)^2 + 2(x-1) - 13 =$$

$$x^2 - 2x - 1 + 2x - 2 - 13 =$$

$$x^2 - 14.$$

Answer: $x^2 - 14$.

Problem 4.2. Factorize

$$a^3 + a - 2.$$

Solution 4.2.

$$a^3+a-2 = a^3-1+a-1 = (a-1)(a^2+a+1)+(a-1) = (a-1)(a^2+a+2).$$

Remark. The value of the expression $a^3 + a - 2$ at a = 1 is equal to 0 which means that it must have factor a - 1 which is 0 at a = 1 (Bezaut).

Problem 4.3. Solve equation in integers

$$x! + 13 = y^2$$
.

Solution 4.3. Note that for the numbers bigger than 4 their factorial is divisible by 10, so their last digit is 0. Therefore for $x \geq 5$ the last digit of the left side of the equation is 3 so it can't be perfect square. So $x \leq$. By verifying each case x = 1, 2, 3, 4 we conclude that the equation has no solution in integers.

Problem 4.4. Let numbers x_1, x_2, \ldots, x_n are given and each of them is equal either +1 or -1. Prove that if

$$x_1x_2 + x_2x_3 + \ldots + x_nx_1 = 0$$

then n is divisible by 4.

Solution 4.4. There are n terms in the sum, and half of them are +1 and half of them are equal -1, so n is even. Now lets prove that there are even number of -1's. From this we will conclude the statement of the problem. To obtian -1 we need consecutive written numbers +1 and -1. Lets write numbers x_1, x_2, \ldots, x_n around the circle and by starting from x_1 go around the circle. Note that we move from +1 to -1 as many times as we move from -1 to +1. So there are even number of terms -1 in the sum $x_1x_2 + x_2x_3 + \ldots + x_nx_1$.

Problem 4.5. Chess king has started from some cell and by passing over each cell exactly ones came back to original position. Prove that the king has done even number of diagonal moves.

Solution 4.5. The king has made 64 moves. Note that when king makes diagonal move then it doesn't change it's cell color, otherwise he changes it. Since he came back to the same cell, it means that he made even number of non-diagonal moves. Since he made in total 64 moves, therefore the number of diagonal moves is even as well.

Problem 4.6. Let k is given and numbers from 1 to 100 are written on the board. Ali erases from the board arbitrary k numbers. Is it true that Bob may choose k numbers written on the board, which sum is equal to 100. Consider cases when a) k = 8, b) k = 9.

Solution 4.6. a) Let's group numbers from 1 to 24 to the pairs such that their sum is equal 25.

$$(1;24), (2;23), (3;22), \ldots, (11;14), (12;13):$$

In total there are 12 pairs. If we remove any 8 numbers then there will be at lease 12-8=4 pairs where no any number is removed. If we take all numbers from these 4 groups then we get 8 numbers which sum is equal $4 \cdot 25 = 100$.

b) Lets remove numbers from 1 to 9. If we choose any 9 numbers from the remaining numbers then their sum will be at least 10 + 11 +

12 + 13 + 14 + 15 + 16 + 17 + 18 = 126 > 100. So we can't choose 9 numbers which sum is equal 100.

Answer: a) possible, b) not possible.