Human expression changes depend on the endpoint

Eight libraries from one donor · four-day treatment comparison

Panel	OSK CPM	OSK median ratio	O4YRSK CPM	O4YRSK median ratio
Fibroblast markers	-1.508	-1.940	-1.500	-1.851
Cell cycle	+1.064	+0.656	+0.564	+0.226
Stress	+0.258	-0.113	+0.561	+0.274
Reprogramming*	+0.628	+0.527	+0.427	+0.354

OSK stress reverses direction under the normalization sensitivity.

Mean log-expression differences. *POU5F1 and SOX2 excluded. No young or recovery arm.

A mouse neural dataset retains seven animal identities


3,857 cells · 31,057 RNA genes · three multiplexed libraries

Cohort-2 group	Animals	Cells
Old treated	2	732
Old untreated	3	1,254
Young untreated	2	1,871

The deposited subset contains no withdrawal samples.
Both young animals occur in lane 3; 70 cells have unresolved cluster labels.

Admission complete. Animal-level expression testing is the next step.

Balanced donors improve repeat detection in the simulation



1,200 cells

Assumed prevalence: 1%
Sensitivity: 80%
Specificity: 100%
Donor correlation: 0.05

Synthetic only

Probability of flags (%). With no true states and 99.9% specificity, six balanced donors still produce a 29.9% chance of flags in two donors under independent errors.

The next analysis and the remaining measurement

Next computation	Animal-level neural expression and composition, with lane sensitivity and a cohort-1 replication audit.
Pending source inputs	Reconcile clock, sample and guide metadata when the requested artifacts arrive.
Biological gap	Matched late human identity, calibrated unusual-state detection, withdrawal stability and trajectory continuity.

All four readouts remain insufficient_signal. Experiment replacement: INCONCLUSIVE.