
Cell-cycle scores fall in all nine donor pairs

Mouse neural cultures after 18 hours of Dox/OSKM exposure



Every donor retains the negative direction under median-ratio and rank scoring.

Mean log2(1 + CPM) score differences. The age contrast depends on the scoring method.

Cody Mitchell / FRG October 18, 2026 Public-data reanalysis

Microglial cell-cycle means agree across cohorts

Individual animal omissions expose the limit of that agreement


Mouse cohort	Full CPM mean	Omit one animal
Cohort 2	-1.014	-1.829 to -0.199
Cohort 1	-0.290	-1.120 to +0.539

Cohort 1 can reverse direction when one animal is removed.

Two treated and two control animals per primary comparison. Omission ranges are descriptive.

CD24 supplies most of the human bulk score change

Seven fixed markers after excluding POU5F1 and SOX2



CD24 contributes 80–90% across the two original normalizations.

One donor, eight libraries. This score does not establish a calibrated pluripotent state.

Cody Mitchell / FRG October 18, 2026 Public-data reanalysis

Human co-detection fractions depend on UMI depth

Day 7: at least two of five endogenous markers in the same barcode


Detection definition	GM00731	GM23815
At least 1 UMI per marker	852/6915 12.32%	2851/12209 23.35%
At least 2 UMIs per marker	321/6915 4.64%	1392/12209 11.40%
1,000-UMI thinning, seed 181026	51/6774 0.75%	311/11552 2.69%

57,317 observed barcodes across two donors and four collection times.

All 22 sensitivity definitions are reported. These are marker observations without an age comparison.

Composition can oppose within-type changes

Conditional stress-panel summary in the two admitted mouse cell types



Symmetric partition (rounded): -0.017309 + 0.012498 = -0.004811.

Common support retains 58.8–68.1% of each animal’s recovered cells. Algebraic attribution is descriptive.

Existing sequenced samples are the next data target

Seven omitted mouse samples

The published sample table lists them. The shared objects lack the complete cell-to-animal assignments. Five carry a withdrawal label whose exact endpoint timing still needs confirmation.

The decisive human measurement remains open

Recover compatible existing data first. Then evaluate matched late identity and age, unusual-state detection and stability after factor withdrawal.

All four CX004 readouts remain insufficient_signal. The conditional funding plan specifies the remaining observations.