

Bioinformatics

CS300

Chap 2

Computational Manipulation of DNA

Fall 2017



Oliver Bonham-Carter



Consider this....

- What is the difference between a gene and an allele?
- Answer in the context of cystic fibrosis and the *CFTR* gene



Hint: *Think Mendelian Genetics*





ALLEGHENY
COLLEGE

Cystic Fibrosis



Cystic Fibrosis

Cystic Fibrosis causes the mucous to become thick and sticky which may make the body prone to infections and can even block the airways.

ePainAssist.com

- Inherited medical condition of the secretory glands (producers of mucous and sweat)



ALLEGHENY
COLLEGE

Cystic Fibrosis




- Clubbed fingers: occurs in heart and lung diseases that reduce the amount of oxygen in the blood



Cystic Fibrosis

- Autosomal recessive type condition: one faulty gene is inherited from both parents (together) in order for the offspring to get this condition
- Mendelian Genetic
- Impossible to know that someone is sure to get a condition.






The Cystic Fibrosis Gene

- Cystic Fibrosis Transmembrane conductance: **CFTR**
- Gene product is a bad regulator which fails to move water after displacing chloride ions in epithelial (thin tissue) cells
- Water follows chloride ions by osmosis.
- **What if water regulation were not possible in the cells and organs?**



Cystic Fibrosis



- Restricted flow in airways from mucous build-ups.
- Suffocation



ALLEGHENY
COLLEGE

A Build-Up of Anything is Bad



- What if the the garbage collection crews in Paris went on strike (as they did in 2016)?



ALLEGHENY
COLLEGE

The Cystic Fibrosis Gene

- Cystic Fibrosis Transmembrane conductance: **CFTR**
- Gene product is a bad regulator which fails to move water after displacing chloride ions in epithelial (thin tissue) cells
- Water follows chloride ions by osmosis.
- <https://www.youtube.com/watch?v=EuLVCYrurok>
- **What if water regulation were not possible in the cells and organs?**



Three Bad Proteins From the Four





Figure 2.2 The wild-type allele (A) of the CFTR gene produces a chloride transport protein localized in the membrane; three different common CF alleles illustrated here result in variant proteins that are folded incorrectly (ΔF508; B), truncated (G542X; C), or unable to transport chloride (G551D; D).



The Cystic Fibrosis Gene

- Gene codes for four different proteins: only one working type to move chloride ions and enable water displacement.

Mucous build-up





Open Reading Frames: Explaining Disorders?

- **Pam Can See The Man and Dog**
- **Frame shift by one letter!**
- **P amC anS eeT heM ana ndD eg**
- **Frame shift by two letters!**
- **Pa mCa nSe eTh eMa nan dDo g**
- **Frame shift by three letters!**
- **Pam Can See The Man and Dog**

Notice how the code changes depending on where you start reading?



Open Reading Frames

Note: RF means *reading frame, where you start reading the words.*

Original: CAATGGCGAATCGACGTGTATAAA

RF1 - 5' - CAA TGG CGA ATC GAC GTG TAT AAA – 3'

RF2 - 5' - C AAT GGC GAA TCG ACG TGT ATA AA – 3'

RF 3 - 5' - CA ATG GCG AAT CGA CGT GTA TAA A – 3'

3' - CAA TGG CGA ATC GAC GTG TAT AAA - 5' – RF 4


3' - C AAT GGC GAA TCG ACG TGT ATA AA - 5' – RF 5

3' - CA ATG GCG AAT CGA CGT GTA TAA A - 5' – RF 6



Sequence is Carrier?


- How do we determine if a sequence carries the CF allele?
- Get DNA sample
- Translate DNA to protein: Compare this seq to seq of a “working protein”
- Is difference found between both proteins?





Analyze the Protein


- Translating DNA to find defects





The Unnamed Sequence


- Unlabeled strands of DNA are assumed to be in the 5' to 3', (left to right) direction.
- A new sequence is given to us for analysis.
- What are the steps to place this sequence into a format for use with bioinformatics tools?





DNA Manipulation Algorithm

- A series of steps when handling DNA






The Algorithm in Words

- Input a DNA sequence, including whether or it template or non-template strand and orientation
- Choose the appropriate operation:
 - If it is the template strand and oriented 3' -> 5', simply output the same sequence
 - If it is the template strand and oriented 5' -> 3', **inverse** the sequence (traverse the string from right to left and add each character to output string)
 - If it is the non-template strand and oriented 3' -> 5', generate the **inverse complement** sequence ((i.) traverse the string from right to left and (ii) for each character add the complement to the output string)
 - If it is the non-template strand and oriented 5' -> 3', generate the **complement** ((i.) traverse the string from left to right and (ii) for each character add the complement to the output string)
- Output the completed sequence, including 5' and 3' end labels



Template vs nonTemplate

- Input:
 - DNA sequence – AGCAT
 - Strand – template (used to make mRNA) or non-template (the compliment of this strand that looks like mRNA)
 - Orientation – 3' -> 5' or 5' -> 3'
- Output:
 - Template strand in 3' -> 5' orientation ready for transcription





Codon Table to Translate the Protein Product

- DNA triplets read in groups of three called codons and represent an amino acid

		Standard genetic code						
1st base	2nd base						3rd base	
	T		C		A			
T	TTT	(Phe/F) Phenylalanine	TCT	(Ser/S) Serine	TAT	(Tyr/Y) Tyrosine	TGT	(Cys/C) Cysteine
	TTC		TCC		TAC		TGC	
	TTA		TCA		TAA ^[B]	Stop (Ochre)	TGA ^[B]	Stop (Opal)
	TTG		TCG		TAG ^[B]	Stop (Amber)	TGG	(Trp/W) Tryptophan
C	CTT	(Leu/L) Leucine	CCT	(Pro/P) Proline	CAT	(His/H) Histidine	CGT	(Arg/R) Arginine
	CTC		CCC		CAC		CGC	
	CTA		CCA		CAA	(Gln/Q) Glutamine	CGA	
	CTG		CCG		CAG		CGG	
A	ATT	(Ile/I) Isoleucine	ACT	(Thr/T) Threonine	AAT	(Asn/N) Asparagine	AGT	(Ser/S) Serine
	ATC		ACC		AAC		AGC	
	ATA		ACA		AAA	(Lys/K) Lysine	AGA	(Arg/R) Arginine
	ATG ^[A]		(Met/M) Methionine		AAG		AGG	
G	GTT	(Val/V) Valine	GCT	(Ala/A) Alanine	GAT	(Asp/D) Aspartic acid	GGT	(Gly/G) Glycine
	GTC		GCC		GAC		GGC	
	GTA		GCA		GAA	(Glu/E) Glutamic acid	GGA	
	GTG		GCG		GAG		GGG	



ALLEGHENY
COLLEGE

Python Programming

- Biopython
- Translation functions
 - DNA → RNA
 - RNA → DNA
 - RNA → Protein
- Gives a protein sequence to compare to the wild type protein sequence

Follow along in class and save your notes in a text file!!

