

Introduction

Marčenko-Pastur's theorem

Large covariance matrices

Spiked models

Statistical Test for Single-Source Detection

The setup

Asymptotic behaviour of the GLRT

Fluctuations of the test statistics

Power of the test

The GLRT: Summary

Direction of Arrival Estimation

Applications to the MIMO channel

Conclusion

The hypothesis testing problem

Statistical Setup

let

$$\vec{y}(k) = \begin{cases} \sigma \vec{w}(k) & \text{under } H_0 \\ \vec{h}_s(k) + \sigma \vec{w}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

The $\vec{y}(k)$'s are n observations all either drawn under H_0 or H_1 . Here,

The hypothesis testing problem

Statistical Setup

let

$$\vec{y}(k) = \begin{cases} \sigma \vec{w}(k) & \text{under } H_0 \\ \vec{h}_s(k) + \sigma \vec{w}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

The $\vec{y}(k)$'s are n observations all either drawn under H_0 or H_1 . Here,

- ▶ $\vec{w}(k)$ is a $N \times 1$ complex gaussian white noise process:

$$\vec{w}(k) \sim \mathcal{CN}(0, \mathbf{I}_N)$$

The hypothesis testing problem

Statistical Setup

let

$$\vec{y}(k) = \begin{cases} \sigma \vec{w}(k) & \text{under } H_0 \\ \vec{h} s(k) + \sigma \vec{w}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

The $\vec{y}(k)$'s are n observations all either drawn under H_0 or H_1 . Here,

- ▶ $\vec{w}(k)$ is a $N \times 1$ complex gaussian white noise process:

$$\vec{w}(k) \sim \mathcal{CN}(0, \mathbf{I}_N)$$

- ▶ \vec{h} is a $N \times 1$ **deterministic** and **unknown** vector and typically represents the propagation channel

The hypothesis testing problem

Statistical Setup

let

$$\vec{y}(k) = \begin{cases} \sigma \vec{w}(k) & \text{under } H_0 \\ \vec{h} s(k) + \sigma \vec{w}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

The $\vec{y}(k)$'s are n observations all either drawn under H_0 or H_1 . Here,

- ▶ $\vec{w}(k)$ is a $N \times 1$ complex gaussian white noise process:

$$\vec{w}(k) \sim \mathcal{CN}(0, \mathbf{I}_N)$$

- ▶ \vec{h} is a $N \times 1$ **deterministic** and **unknown** vector and typically represents the propagation channel
- ▶ $s(k)$ represent the signal; it is a scalar complex gaussian i.i.d. process

The hypothesis testing problem

Statistical Setup

let

$$\vec{y}(k) = \begin{cases} \sigma \vec{w}(k) & \text{under } H_0 \\ \vec{h} s(k) + \sigma \vec{w}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

The $\vec{y}(k)$'s are n observations all either drawn under H_0 or H_1 . Here,

- ▶ $\vec{w}(k)$ is a $N \times 1$ complex gaussian white noise process:

$$\vec{w}(k) \sim \mathcal{CN}(0, \mathbf{I}_N)$$

- ▶ \vec{h} is a $N \times 1$ **deterministic** and **unknown** vector and typically represents the propagation channel
- ▶ $s(k)$ represent the signal; it is a scalar complex gaussian i.i.d. process

Objective

Given n observations $(\vec{y}(k), 1 \leq k \leq n)$, and the associated **sample covariance matrix**

$$\hat{\mathbf{R}}_n = \frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^* \quad \text{where} \quad \mathbf{Y}_n = [\vec{y}(1), \dots, \vec{y}(n)] \quad \text{is } N \times n ,$$

The hypothesis testing problem

Statistical Setup

let

$$\vec{y}(k) = \begin{cases} \sigma \vec{w}(k) & \text{under } H_0 \\ \vec{h}s(k) + \sigma \vec{w}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

The $\vec{y}(k)$'s are n observations all either drawn under H_0 or H_1 . Here,

- ▶ $\vec{w}(k)$ is a $N \times 1$ complex gaussian white noise process:

$$\vec{w}(k) \sim \mathcal{CN}(0, \mathbf{I}_N)$$

- ▶ \vec{h} is a $N \times 1$ **deterministic** and **unknown** vector and typically represents the propagation channel
- ▶ $s(k)$ represent the signal; it is a scalar complex gaussian i.i.d. process

Objective

Given n observations $(\vec{y}(k), 1 \leq k \leq n)$, and the associated **sample covariance matrix**

$$\hat{\mathbf{R}}_n = \frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^* \quad \text{where} \quad \mathbf{Y}_n = [\vec{y}(1), \dots, \vec{y}(n)] \quad \text{is } N \times n ,$$

the aim is to decide H_0 (no signal) or H_1 (single-source detection) in the case where

$$\frac{N}{n} \rightarrow c \in (0, 1) \quad \text{i.e.}$$

The hypothesis testing problem

Statistical Setup

let

$$\vec{y}(k) = \begin{cases} \sigma \vec{w}(k) & \text{under } H_0 \\ \vec{h}s(k) + \sigma \vec{w}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

The $\vec{y}(k)$'s are n observations all either drawn under H_0 or H_1 . Here,

- ▶ $\vec{w}(k)$ is a $N \times 1$ complex gaussian white noise process:

$$\vec{w}(k) \sim \mathcal{CN}(0, \mathbf{I}_N)$$

- ▶ \vec{h} is a $N \times 1$ **deterministic** and **unknown** vector and typically represents the propagation channel
- ▶ $s(k)$ represent the signal; it is a scalar complex gaussian i.i.d. process

Objective

Given n observations $(\vec{y}(k), 1 \leq k \leq n)$, and the associated **sample covariance matrix**

$$\hat{\mathbf{R}}_n = \frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^* \quad \text{where} \quad \mathbf{Y}_n = [\vec{y}(1), \dots, \vec{y}(n)] \quad \text{is } N \times n ,$$

the aim is to decide H_0 (no signal) or H_1 (single-source detection) in the case where

$$\frac{N}{n} \rightarrow c \in (0, 1) \quad \text{i.e.} \quad \boxed{\text{Dimension } N \text{ of observations } \propto \text{size } n \text{ of sample}}$$

Neyman-Pearson procedure

Likelihood functions

Neyman-Pearson procedure

Likelihood functions

Notice that \mathbf{Y}_n is a $N \times n$ matrix whose columns are i.i.d. vectors with covariance matrix defined by

$$\Sigma_N = \begin{cases} \sigma^2 \mathbf{I}_N & \text{under } H_0, \\ \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N & \text{under } H_1 \end{cases}$$

Neyman-Pearson procedure

Likelihood functions

Notice that \mathbf{Y}_n is a $N \times n$ matrix whose columns are i.i.d. vectors with covariance matrix defined by

$$\mathbf{\Sigma}_N = \begin{cases} \sigma^2 \mathbf{I}_N & \text{under } H_0, \\ \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N & \text{under } H_1 \end{cases}$$

hence the likelihood functions write

$$p_0(\mathbf{Y}_N; \sigma^2) = \frac{1}{(\pi\sigma^2)^{Nn}} \exp\left(-\frac{n}{\sigma^2} \text{Trace } \hat{\mathbf{R}}_N\right)$$

$$p_1(\mathbf{Y}_N; \vec{\mathbf{h}}; \sigma^2) = \frac{1}{\left[\pi^N \det(\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)\right]^n} \exp\left(-\frac{n}{\sigma^2} \text{Trace } \hat{\mathbf{R}}_N (\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)^{-1}\right)$$

Neyman-Pearson procedure

Likelihood functions

Notice that \mathbf{Y}_n is a $N \times n$ matrix whose columns are i.i.d. vectors with covariance matrix defined by

$$\boldsymbol{\Sigma}_N = \begin{cases} \sigma^2 \mathbf{I}_N & \text{under } H_0, \\ \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N & \text{under } H_1 \end{cases}$$

hence the likelihood functions write

$$p_0(\mathbf{Y}_N; \sigma^2) = \frac{1}{(\pi\sigma^2)^{Nn}} \exp\left(-\frac{n}{\sigma^2} \text{Trace } \hat{\mathbf{R}}_N\right)$$

$$p_1(\mathbf{Y}_N; \vec{\mathbf{h}}; \sigma^2) = \frac{1}{\left[\pi^N \det(\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)\right]^n} \exp\left(-\frac{n}{\sigma^2} \text{Trace } \hat{\mathbf{R}}_N (\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)^{-1}\right)$$

Neyman-Pearson

In case where σ^2 and $\vec{\mathbf{h}}$ are **known**, the
Likelihood Ratio Statistics

$$\frac{p_1(\mathbf{Y}_N; \vec{\mathbf{h}}; \sigma^2)}{p_0(\mathbf{Y}_N; \sigma^2)}$$

provides a **uniformly most powerful test**:

Neyman-Pearson procedure

Likelihood functions

Notice that \mathbf{Y}_n is a $N \times n$ matrix whose columns are i.i.d. vectors with covariance matrix defined by

$$\Sigma_N = \begin{cases} \sigma^2 \mathbf{I}_N & \text{under } H_0, \\ \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N & \text{under } H_1 \end{cases}$$

hence the likelihood functions write

$$p_0(\mathbf{Y}_N; \sigma^2) = \frac{1}{(\pi\sigma^2)^{Nn}} \exp\left(-\frac{n}{\sigma^2} \text{Trace } \hat{\mathbf{R}}_N\right)$$

$$p_1(\mathbf{Y}_N; \vec{\mathbf{h}}; \sigma^2) = \frac{1}{\left[\pi^N \det(\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)\right]^n} \exp\left(-\frac{n}{\sigma^2} \text{Trace } \hat{\mathbf{R}}_N (\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)^{-1}\right)$$

Neyman-Pearson

- Fix a given level $\alpha \in (0, 1)$

In case where σ^2 and $\vec{\mathbf{h}}$ are **known**, the **Likelihood Ratio Statistics**

$$\frac{p_1(\mathbf{Y}_N; \vec{\mathbf{h}}; \sigma^2)}{p_0(\mathbf{Y}_N; \sigma^2)}$$

provides a **uniformly most powerful test**:

Neyman-Pearson procedure

Likelihood functions

Notice that \mathbf{Y}_n is a $N \times n$ matrix whose columns are i.i.d. vectors with covariance matrix defined by

$$\Sigma_N = \begin{cases} \sigma^2 \mathbf{I}_N & \text{under } H_0, \\ \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N & \text{under } H_1 \end{cases}$$

hence the likelihood functions write

$$p_0(\mathbf{Y}_N; \sigma^2) = \frac{1}{(\pi\sigma^2)^{Nn}} \exp\left(-\frac{n}{\sigma^2} \text{Trace } \hat{\mathbf{R}}_N\right)$$

$$p_1(\mathbf{Y}_N; \vec{\mathbf{h}}; \sigma^2) = \frac{1}{\left[\pi^N \det(\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)\right]^n} \exp\left(-\frac{n}{\sigma^2} \text{Trace } \hat{\mathbf{R}}_N (\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)^{-1}\right)$$

Neyman-Pearson

In case where σ^2 and $\vec{\mathbf{h}}$ are **known**, the **Likelihood Ratio Statistics**

$$\frac{p_1(\mathbf{Y}_N; \vec{\mathbf{h}}; \sigma^2)}{p_0(\mathbf{Y}_N; \sigma^2)}$$

provides a **uniformly most powerful test**:

- Fix a given level $\alpha \in (0, 1)$
- The condition over the **Probability of False Alarm** $\mathbb{P}(H_1 | H_0) \leq \alpha$ sets the threshold

Neyman-Pearson procedure

Likelihood functions

Notice that \mathbf{Y}_n is a $N \times n$ matrix whose columns are i.i.d. vectors with covariance matrix defined by

$$\Sigma_N = \begin{cases} \sigma^2 \mathbf{I}_N & \text{under } H_0, \\ \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N & \text{under } H_1 \end{cases}$$

hence the likelihood functions write

$$p_0(\mathbf{Y}_N; \sigma^2) = \frac{1}{(\pi\sigma^2)^{Nn}} \exp\left(-\frac{n}{\sigma^2} \text{Trace } \hat{\mathbf{R}}_N\right)$$

$$p_1(\mathbf{Y}_N; \vec{\mathbf{h}}; \sigma^2) = \frac{1}{\left[\pi^N \det(\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)\right]^n} \exp\left(-\frac{n}{\sigma^2} \text{Trace } \hat{\mathbf{R}}_N (\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)^{-1}\right)$$

Neyman-Pearson

In case where σ^2 and $\vec{\mathbf{h}}$ are **known**, the **Likelihood Ratio Statistics**

$$\frac{p_1(\mathbf{Y}_N; \vec{\mathbf{h}}; \sigma^2)}{p_0(\mathbf{Y}_N; \sigma^2)}$$

provides a **uniformly most powerful test**:

- ▶ Fix a given level $\alpha \in (0, 1)$
- ▶ The condition over the **Probability of False Alarm** $\mathbb{P}(H_1 | H_0) \leq \alpha$ sets the threshold
- ▶ the **maximum achievable power**

$$1 - \mathbb{P}(H_0 | H_1)$$

is guaranteed by Neyman-Pearson.

The GLRT

The Generalized Likelihood Ratio Test

In the case where $\vec{\mathbf{h}}$ and σ^2 are unknown, we use instead:

$$L_n = \frac{\sup_{\sigma^2, \vec{\mathbf{h}}} p_1(\mathbf{Y}_n, \sigma^2, \vec{\mathbf{h}})}{\sup_{\sigma^2} p_0(\mathbf{Y}_n, \sigma^2)}$$

which is no longer uniformly most powerful.

The GLRT

The Generalized Likelihood Ratio Test

In the case where $\vec{\mathbf{h}}$ and σ^2 are unknown, we use instead:

$$L_n = \frac{\sup_{\sigma^2, \vec{\mathbf{h}}} p_1(\mathbf{Y}_n, \sigma^2, \vec{\mathbf{h}})}{\sup_{\sigma^2} p_0(\mathbf{Y}_n, \sigma^2)}$$

which is no longer uniformly most powerful.

Expression of the GLRT

The GLRT statistics writes

$$L_n = \frac{\left(1 - \frac{1}{N}\right)^{(1-N)n}}{\left(\frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}\right)^n \left(1 - \frac{1}{N} \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}\right)^{(N-1)n}}$$

and is a **deterministic function** of $T_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$

Introduction

Marčenko-Pastur's theorem

Large covariance matrices

Spiked models

Statistical Test for Single-Source Detection

The setup

Asymptotic behaviour of the GLRT

Fluctuations of the test statistics

Power of the test

The GLRT: Summary

Direction of Arrival Estimation

Applications to the MIMO channel

Conclusion

Limit of the test statistics T_n I

Under H_0

Recall $T_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$.

Limit of the test statistics T_n I

Under H_0

Recall $T_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$. We have:

$$\lambda_{\max}(\hat{\mathbf{R}}_n) \xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2(1 + \sqrt{c})^2$$

Limit of the test statistics T_n I

Under H_0

Recall $T_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$. We have:

$$\begin{aligned} \lambda_{\max}(\hat{\mathbf{R}}_n) &\xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2(1 + \sqrt{c})^2 \\ \frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n = \frac{1}{Nn} \sum_{i,j} |Y_{ij}|^2 &\xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2 \end{aligned}$$

Limit of the test statistics T_n I

Under H_0

Recall $T_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$. We have:

$$\begin{aligned}\lambda_{\max}(\hat{\mathbf{R}}_n) &\xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2(1 + \sqrt{c})^2 \\ \frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n = \frac{1}{Nn} \sum_{i,j} |Y_{ij}|^2 &\xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2\end{aligned}$$

hence

$$T_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n} \xrightarrow[N, n \rightarrow \infty]{a.s.} (1 + \sqrt{c})^2$$

Limit of the test statistics T_n II

Under H_1

Denote by

$$\mathbf{snr} = \frac{\|\vec{\mathbf{h}}\|^2}{\sigma^2}$$

the **Signal-to-Noise (SNR)** ratio.

Limit of the test statistics T_n II

Under H_1

Denote by

$$\mathbf{snr} = \frac{\|\vec{\mathbf{h}}\|^2}{\sigma^2}$$

the **Signal-to-Noise (SNR)** ratio.

► if $\mathbf{snr} > \sqrt{c}$ then

$$T_n \xrightarrow[N, n \rightarrow \infty]{a.s.} (1 + \mathbf{snr}) \left(1 + \frac{c}{\mathbf{snr}}\right)$$

Limit of the test statistics T_n II

Under H_1

Denote by

$$\mathbf{snr} = \frac{\|\vec{\mathbf{h}}\|^2}{\sigma^2}$$

the **Signal-to-Noise (SNR)** ratio.

► if $\mathbf{snr} > \sqrt{c}$ then

$$T_n \xrightarrow[N, n \rightarrow \infty]{a.s.} (1 + \mathbf{snr}) \left(1 + \frac{c}{\mathbf{snr}}\right) > (1 + \sqrt{c})^2$$

Limit of the test statistics T_n II

Under H_1

Denote by

$$\mathbf{snr} = \frac{\|\vec{\mathbf{h}}\|^2}{\sigma^2}$$

the **Signal-to-Noise (SNR)** ratio.

► if $\mathbf{snr} > \sqrt{c}$ then

$$T_n \xrightarrow[N, n \rightarrow \infty]{a.s.} (1 + \mathbf{snr}) \left(1 + \frac{c}{\mathbf{snr}}\right) > (1 + \sqrt{c})^2$$

► if $\mathbf{snr} \leq \sqrt{c}$ then

$$T_n \xrightarrow[N, n \rightarrow \infty]{a.s.} (1 + \sqrt{c})^2$$

Limit of the test statistics T_n III

Remarks

- Condition $\boxed{\text{snr} > \sqrt{c}}$ is **automatically fulfilled** in the standard regime where

$$N \text{ fixed} \quad \text{and} \quad n \rightarrow \infty \quad \text{as} \quad c = \lim_{n \rightarrow \infty} \frac{N}{n} = 0 .$$

Limit of the test statistics T_n III

Remarks

- ▶ Condition $\boxed{\text{snr} > \sqrt{c}}$ is **automatically fulfilled** in the standard regime where

$$N \text{ fixed and } n \rightarrow \infty \text{ as } c = \lim_{n \rightarrow \infty} \frac{N}{n} = 0 .$$

- ▶ In the case $N, n \rightarrow \infty$, recall that the support of Marčenko-Pastur distribution is

$$[(1 - \sqrt{c})^2, (1 + \sqrt{c})^2] ,$$

Limit of the test statistics T_n III

Remarks

- ▶ Condition $\boxed{\text{snr} > \sqrt{c}}$ is **automatically fulfilled** in the standard regime where

$$N \text{ fixed and } n \rightarrow \infty \text{ as } c = \lim_{n \rightarrow \infty} \frac{N}{n} = 0 .$$

- ▶ In the case $N, n \rightarrow \infty$, recall that the support of Marčenko-Pastur distribution is

$$[(1 - \sqrt{c})^2, (1 + \sqrt{c})^2] ,$$

i.e.

$\boxed{\text{The higher } \sqrt{c}, \text{ the larger the support}}$

Limit of the test statistics T_n III

Remarks

- ▶ Condition $\boxed{\text{snr} > \sqrt{c}}$ is **automatically fulfilled** in the standard regime where

$$N \text{ fixed and } n \rightarrow \infty \text{ as } c = \lim_{n \rightarrow \infty} \frac{N}{n} = 0 .$$

- ▶ In the case $N, n \rightarrow \infty$, recall that the support of Marčenko-Pastur distribution is

$$[(1 - \sqrt{c})^2, (1 + \sqrt{c})^2] ,$$

i.e.

$\boxed{\text{The higher } \sqrt{c}, \text{ the larger the support}}$

One can interpret \sqrt{c} as a level of the **asymptotic noise induced by the data dimension** (=asymptotic data noise).

Limit of the test statistics T_n III

Remarks

- ▶ Condition $\text{snr} > \sqrt{c}$ is **automatically fulfilled** in the standard regime where

$$N \text{ fixed and } n \rightarrow \infty \text{ as } c = \lim_{n \rightarrow \infty} \frac{N}{n} = 0 .$$

- ▶ In the case $N, n \rightarrow \infty$, recall that the support of Marčenko-Pastur distribution is

$$[(1 - \sqrt{c})^2, (1 + \sqrt{c})^2] ,$$

i.e.

The higher \sqrt{c} , the larger the support

One can interpret \sqrt{c} as a level of the **asymptotic noise induced by the data dimension** (=asymptotic data noise).

Hence the rule of thumb

Detection occurs if **snr** higher than **asymptotic data noise**.

Simulations




Figure : Influence of **asymptotic data noise** as \sqrt{c} increases

Simulations




Figure : Influence of **asymptotic data noise** as \sqrt{c} increases

Simulations




Figure : Influence of **asymptotic data noise** as \sqrt{c} increases

Simulations




Figure : Influence of **asymptotic data noise** as \sqrt{c} increases

Simulations




Figure : Influence of **asymptotic data noise** as \sqrt{c} increases

Elements of proof I

- We are interested in the largest eigenvalue of the matrix model

$$\frac{\frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^*}{\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)}$$

Elements of proof I

- We are interested in the largest eigenvalue of the matrix model

$$\frac{\frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^*}{\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)}$$

asymptotically equivalent to

$$\frac{1}{n} \frac{\mathbf{Y}_n \mathbf{Y}_n^*}{\sigma^2} \quad \text{as} \quad \frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n) \xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2$$

Elements of proof I

- We are interested in the largest eigenvalue of the matrix model

$$\frac{\frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^*}{\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)}$$

asymptotically equivalent to

$$\frac{1}{n} \frac{\mathbf{Y}_n \mathbf{Y}_n^*}{\sigma^2} \quad \text{as} \quad \frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n) \xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2$$

- Notice that

$$\mathbf{Y}_n = [\vec{\mathbf{y}}_1, \dots, \vec{\mathbf{y}}_n] \quad \text{with} \quad \vec{\mathbf{y}}_i \sim \mathcal{CN}(0, \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)$$

Elements of proof I

- We are interested in the largest eigenvalue of the matrix model

$$\frac{\frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^*}{\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)}$$

asymptotically equivalent to

$$\frac{1}{n} \frac{\mathbf{Y}_n \mathbf{Y}_n^*}{\sigma^2} \quad \text{as} \quad \frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n) \xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2$$

- Notice that

$$\mathbf{Y}_n = [\vec{\mathbf{y}}_1, \dots, \vec{\mathbf{y}}_n] \quad \text{with} \quad \vec{\mathbf{y}}_i \sim \mathcal{CN}(0, \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)$$

Hence

$$\mathbf{Y}_N = \left(\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N \right)^{1/2} \mathbf{X}_N$$

Elements of proof I

- We are interested in the largest eigenvalue of the matrix model

$$\frac{\frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^*}{\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)}$$

asymptotically equivalent to

$$\frac{1}{n} \frac{\mathbf{Y}_n \mathbf{Y}_n^*}{\sigma^2} \quad \text{as} \quad \frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n) \xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2$$

- Notice that

$$\mathbf{Y}_n = [\vec{\mathbf{y}}_1, \dots, \vec{\mathbf{y}}_n] \quad \text{with} \quad \vec{\mathbf{y}}_i \sim \mathcal{CN}(0, \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)$$

Hence

$$\mathbf{Y}_N = \left(\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N \right)^{1/2} \mathbf{X}_N \quad \Rightarrow \quad \frac{\mathbf{Y}_N}{\sigma} = \left(\mathbf{I}_N + \frac{\vec{\mathbf{h}}\vec{\mathbf{h}}^*}{\sigma^2} \right)^{1/2} \mathbf{X}_N$$

Elements of proof I

- We are interested in the largest eigenvalue of the matrix model

$$\frac{\frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^*}{\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)}$$

asymptotically equivalent to

$$\frac{1}{n} \frac{\mathbf{Y}_n \mathbf{Y}_n^*}{\sigma^2} \quad \text{as} \quad \frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n) \xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2$$

- Notice that

$$\mathbf{Y}_n = [\vec{\mathbf{y}}_1, \dots, \vec{\mathbf{y}}_n] \quad \text{with} \quad \vec{\mathbf{y}}_i \sim \mathcal{CN}(0, \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)$$

Hence

$$\begin{aligned} \mathbf{Y}_N = \left(\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N \right)^{1/2} \mathbf{X}_N &\Rightarrow \frac{\mathbf{Y}_N}{\sigma} = \left(\mathbf{I}_N + \frac{\vec{\mathbf{h}}\vec{\mathbf{h}}^*}{\sigma^2} \right)^{1/2} \mathbf{X}_N \\ &= \left(\mathbf{I}_N + \frac{\|\vec{\mathbf{h}}\|^2}{\sigma^2} \vec{\mathbf{u}}\vec{\mathbf{u}}^* \right)^{1/2} \mathbf{X}_N \end{aligned}$$

with \mathbf{X}_N a $N \times n$ matrix having i.i.d. entries $\mathcal{CN}(0, 1)$ and $\vec{\mathbf{u}} = \frac{\vec{\mathbf{h}}}{\|\vec{\mathbf{h}}\|}$

Elements of proof I

- We are interested in the largest eigenvalue of the matrix model

$$\frac{\frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^*}{\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)}$$

asymptotically equivalent to

$$\frac{1}{n} \frac{\mathbf{Y}_n \mathbf{Y}_n^*}{\sigma^2} \quad \text{as} \quad \frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n) \xrightarrow[N, n \rightarrow \infty]{a.s.} \sigma^2$$

- Notice that

$$\mathbf{Y}_n = [\vec{\mathbf{y}}_1, \dots, \vec{\mathbf{y}}_n] \quad \text{with} \quad \vec{\mathbf{y}}_i \sim \mathcal{CN}(0, \vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N)$$

Hence

$$\begin{aligned} \mathbf{Y}_N &= \left(\vec{\mathbf{h}}\vec{\mathbf{h}}^* + \sigma^2 \mathbf{I}_N \right)^{1/2} \mathbf{X}_N \quad \Rightarrow \quad \frac{\mathbf{Y}_N}{\sigma} = \left(\mathbf{I}_N + \frac{\vec{\mathbf{h}}\vec{\mathbf{h}}^*}{\sigma^2} \right)^{1/2} \mathbf{X}_N \\ &= \left(\mathbf{I}_N + \frac{\|\vec{\mathbf{h}}\|^2}{\sigma^2} \vec{\mathbf{u}}\vec{\mathbf{u}}^* \right)^{1/2} \mathbf{X}_N \end{aligned}$$

with \mathbf{X}_N a $N \times n$ matrix having i.i.d. entries $\mathcal{CN}(0, 1)$ and $\vec{\mathbf{u}} = \frac{\vec{\mathbf{h}}}{\|\vec{\mathbf{h}}\|}$

Conclusion

Spectrum of $\frac{1}{n} \mathbf{Y}_n \mathbf{Y}_n^*$ follows a spiked model with **rank-one perturbation**

Elements of proof II

We can now conclude:

Elements of proof II

We can now conclude:

► If $\boxed{\text{snr} > \sqrt{c}}$ then

$$\frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)} \xrightarrow[N, n \rightarrow \infty]{(H_1)} (1 + \text{snr}) \left(1 + \frac{c}{\text{snr}}\right) > (1 + \sqrt{c})^2$$

and the test statistics discriminates between the hypotheses H_0 and H_1 .

Elements of proof II

We can now conclude:

- If $\boxed{\text{snr} > \sqrt{c}}$ then

$$\frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)} \xrightarrow[N, n \rightarrow \infty]{(H_1)} (1 + \text{snr}) \left(1 + \frac{c}{\text{snr}}\right) > (1 + \sqrt{c})^2$$

and the test statistics discriminates between the hypotheses H_0 and H_1 .

- If $\boxed{\text{snr} \leq \sqrt{c}}$ then

$$\frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)} \xrightarrow[N, n \rightarrow \infty]{(H_1)} (1 + \sqrt{c})^2$$

Same limit as under H_0 . The test statistics does not discriminate between the two hypotheses.

Introduction

Marčenko-Pastur's theorem

Large covariance matrices

Spiked models

Statistical Test for Single-Source Detection

The setup

Asymptotic behaviour of the GLRT

Fluctuations of the test statistics

Power of the test

The GLRT: Summary

Direction of Arrival Estimation

Applications to the MIMO channel

Conclusion

Fluctuations of the GLRT under H_0 - I

- The exact distribution of the statistics

$$L_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$$

is needed to set the threshold of the test for a given confidence level $\alpha \in (0, 1)$:

$$\mathbb{P}_{H_0} (L_N > \mathbf{t}_\alpha) = \alpha ,$$

Fluctuations of the GLRT under H_0 - I

- The exact distribution of the statistics

$$L_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$$

is needed to set the threshold of the test for a given confidence level $\alpha \in (0, 1)$:

$$\mathbb{P}_{H_0} (L_N > \mathbf{t}_\alpha) = \alpha ,$$

but **hard to obtain**.

Fluctuations of the GLRT under H_0 - I

- The exact distribution of the statistics

$$L_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$$

is needed to set the threshold of the test for a given confidence level $\alpha \in (0, 1)$:

$$\mathbb{P}_{H_0} (L_N > \mathbf{t}_\alpha) = \alpha ,$$

but **hard to obtain**.

- We rather study the asymptotic fluctuations of L_n under the regime

$$N, n \rightarrow \infty , \quad \frac{N}{n} \rightarrow c \in (0, 1) .$$

Fluctuations of the GLRT under H_0 - I

- ▶ The exact distribution of the statistics

$$L_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$$

is needed to set the threshold of the test for a given confidence level $\alpha \in (0, 1)$:

$$\mathbb{P}_{H_0}(L_N > t_\alpha) = \alpha ,$$

but **hard to obtain**.

- ▶ We rather study the asymptotic fluctuations of L_n under the regime

$$N, n \rightarrow \infty , \quad \frac{N}{n} \rightarrow c \in (0, 1) .$$

- ▶ L_N is the ratio of **two random variables**. We need to understand

Fluctuations of the GLRT under H_0 - I

- ▶ The exact distribution of the statistics

$$L_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$$

is needed to set the threshold of the test for a given confidence level $\alpha \in (0, 1)$:

$$\mathbb{P}_{H_0}(L_N > \mathbf{t}_\alpha) = \alpha ,$$

but **hard to obtain**.

- ▶ We rather study the asymptotic fluctuations of L_n under the regime

$$N, n \rightarrow \infty , \quad \frac{N}{n} \rightarrow c \in (0, 1) .$$

- ▶ L_N is the ratio of **two random variables**. We need to understand
 - the fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_0 ,

Fluctuations of the GLRT under H_0 - I

- ▶ The exact distribution of the statistics

$$L_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$$

is needed to set the threshold of the test for a given confidence level $\alpha \in (0, 1)$:

$$\mathbb{P}_{H_0}(L_N > \mathbf{t}_\alpha) = \alpha ,$$

but **hard to obtain**.

- ▶ We rather study the asymptotic fluctuations of L_n under the regime

$$N, n \rightarrow \infty , \quad \frac{N}{n} \rightarrow c \in (0, 1) .$$

- ▶ L_N is the ratio of **two random variables**. We need to understand
 - the fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_0 ,
 - the fluctuations of $\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n$ under H_0 .

Fluctuations of the GLRT under H_0 - II

Fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$: Tracy-Widom distribution **at rate** $N^{2/3}$

Fluctuations of the GLRT under H_0 - II

Fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$: Tracy-Widom distribution **at rate $N^{2/3}$**

$$\frac{N^{2/3}}{\Theta_N} \left\{ \lambda_{\max}(\hat{\mathbf{R}}_n) - \sigma^2(1 + \sqrt{c_n})^2 \right\} \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathbb{P}_{\text{TW}}$$

Fluctuations of the GLRT under H_0 - II

Fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$: Tracy-Widom distribution **at rate $N^{2/3}$**

$$\frac{N^{2/3}}{\Theta_N} \left\{ \lambda_{\max}(\hat{\mathbf{R}}_n) - \sigma^2(1 + \sqrt{c_n})^2 \right\} \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathbb{P}_{\text{TW}}$$

where

$$c_n = \frac{N}{n} \quad \text{and} \quad \Theta_N = \sigma^2(1 + \sqrt{c_n}) \left(\frac{1}{\sqrt{c_n}} + 1 \right)^{1/3}$$

Fluctuations of the GLRT under H_0 - II

Fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$: Tracy-Widom distribution **at rate $N^{2/3}$**

$$\frac{N^{2/3}}{\Theta_N} \left\{ \lambda_{\max}(\hat{\mathbf{R}}_n) - \sigma^2(1 + \sqrt{c_n})^2 \right\} \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathbb{P}_{\text{TW}}$$

where

$$c_n = \frac{N}{n} \quad \text{and} \quad \Theta_N = \sigma^2(1 + \sqrt{c_n}) \left(\frac{1}{\sqrt{c_n}} + 1 \right)^{1/3}$$

Otherwise stated,

$$\lambda_{\max}(\hat{\mathbf{R}}_n) = \sigma^2(1 + \sqrt{c_n})^2 + \frac{\Theta_N}{N^{2/3}} \mathbf{X}_{\text{TW}} + \varepsilon_n$$

where \mathbf{X}_{TW} is a random variable with Tracy-Widom distribution.

Details on Tracy-Widom distribution

Tracy-Widom distribution is defined by

- ▶ its cumulative distribution function

$$F_{TW}(x) = \exp \left\{ - \int_x^\infty (u - x)^2 q^2(u) du \right\}$$

- ▶ where

$$q''(x) = xq(x) + 2q^3(x) \quad \text{and} \quad q(x) \sim \text{Ai}(x) \text{ as } x \rightarrow \infty .$$

$x \mapsto \text{Ai}(x)$ being the Airy function.

Details on Tracy-Widom distribution

Tracy-Widom distribution is defined by

- ▶ its cumulative distribution function

$$F_{TW}(x) = \exp \left\{ - \int_x^\infty (u - x)^2 q^2(u) du \right\}$$

- ▶ where

$$q''(x) = xq(x) + 2q^3(x) \quad \text{and} \quad q(x) \sim \text{Ai}(x) \text{ as } x \rightarrow \infty .$$

$x \mapsto \text{Ai}(x)$ being the Airy function.

Don't bother .. just download it

- ▶ For simulations, cf. R Package 'RMTstat', by Johnstone et al.
- ▶ Also, Folkmar Bornemann (TU München) has developed fast matlab code

Tracy-Widom curve




Figure : Tracy-Widom density

Tracy-Widom curve




Figure : Fluctuations of the largest eigenvalue $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_0

Fluctuations of the GLRT under H_0 - III

Fluctuations of $\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)$: Gaussian distributions **at rate N**

$$N \left\{ \frac{1}{N} \sum_{i=1}^N \lambda_i(\hat{\mathbf{R}}_n) - \sigma^2 \right\} \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, \Gamma) ,$$

Fluctuations of the GLRT under H_0 - III

Fluctuations of $\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n)$: Gaussian distributions **at rate N**

$$N \left\{ \frac{1}{N} \sum_{i=1}^N \lambda_i(\hat{\mathbf{R}}_n) - \sigma^2 \right\} \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, \Gamma) ,$$

Otherwise stated:

$$\frac{1}{N} \text{Trace}(\hat{\mathbf{R}}_n) = \frac{1}{N} \sum_{i=1}^N \lambda_i(\hat{\mathbf{R}}_n) = \sigma^2 + \frac{\sqrt{\Gamma}}{N} \mathbf{Z} + \varepsilon_n$$

where \mathbf{Z} is a random variable with distribution $\mathcal{N}(0, 1)$.

Fluctuations of the GLRT under H_0 - IV

Conclusion

- Fluctuations of $L_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$ are driven by $\lambda_{\max}(\hat{\mathbf{R}}_n)$:

$$\frac{N^{2/3}}{\tilde{\Theta}_N} \{L_N - (1 + \sqrt{c_n})^2\} \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathbb{P}_{\text{TW}} \quad \text{with} \quad \tilde{\Theta}_N = (1 + \sqrt{c_n}) \left(\frac{1}{\sqrt{c_n}} + 1 \right)^{1/3}$$

Fluctuations of the GLRT under H_0 - IV

Conclusion

- Fluctuations of $L_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$ are driven by $\lambda_{\max}(\hat{\mathbf{R}}_n)$:

$$\frac{N^{2/3}}{\tilde{\Theta}_N} \{L_N - (1 + \sqrt{c_n})^2\} \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathbb{P}_{\text{TW}} \quad \text{with} \quad \tilde{\Theta}_N = (1 + \sqrt{c_n}) \left(\frac{1}{\sqrt{c_n}} + 1 \right)^{1/3}$$

- In order to set the threshold α , we choose t_α^n as

$$t_\alpha^n = (1 + \sqrt{c_n})^2 + \frac{\tilde{\Theta}_N}{N^{2/3}} t_\alpha^{\text{Tracy-Widom}}$$

where $t_\alpha^{\text{Tracy-Widom}}$ is the corresponding quantile for a Tracy-Widom random variable:

$$\mathbb{P}\{\mathbf{X}_{\text{TW}} > t_\alpha^{\text{Tracy-Widom}}\} \leq \alpha.$$

Introduction

Marčenko-Pastur's theorem

Large covariance matrices

Spiked models

Statistical Test for Single-Source Detection

The setup

Asymptotic behaviour of the GLRT

Fluctuations of the test statistics

Power of the test

The GLRT: Summary

Direction of Arrival Estimation

Applications to the MIMO channel

Conclusion

Power of the GLRT I

Type II error and Power of the test

Power of the GLRT I

Type II error and Power of the test

Given a level of confidence $\alpha \in (0, 1)$, the **type I error** defines the associate quantile t_α

$$\mathbb{P}_{H_0} (L_N > t_\alpha^n) \leq \alpha .$$

Power of the GLRT I

Type II error and Power of the test

Given a level of confidence $\alpha \in (0, 1)$, the **type I error** defines the associate quantile t_α

$$\mathbb{P}_{H_0} (L_N > t_\alpha^n) \leq \alpha .$$

The **type II error** is defined as

$$\mathbb{P}_{H_1} (L_N < t_\alpha^n) ,$$

Power of the GLRT I

Type II error and Power of the test

Given a level of confidence $\alpha \in (0, 1)$, the **type I error** defines the associate quantile t_α

$$\mathbb{P}_{H_0} (L_N > t_\alpha^n) \leq \alpha .$$

The **type II error** is defined as

$$\mathbb{P}_{H_1} (L_N < t_\alpha^n) ,$$

and the associated **power of the test** is defined as

$$\mathbb{P}_{H_1} (L_N \geq t_\alpha^n) = 1 - \mathbb{P}_{H_1} (L_N < t_\alpha^n) .$$

Power of the GLRT I

Type II error and Power of the test

Given a level of confidence $\alpha \in (0, 1)$, the **type I error** defines the associate quantile t_α

$$\mathbb{P}_{H_0} (L_N > t_\alpha^n) \leq \alpha .$$

The **type II error** is defined as

$$\mathbb{P}_{H_1} (L_N < t_\alpha^n) ,$$

and the associated **power of the test** is defined as

$$\mathbb{P}_{H_1} (L_N \geq t_\alpha^n) = 1 - \mathbb{P}_{H_1} (L_N < t_\alpha^n) .$$

No optimality

Contrary to Neyman-Pearson procedure, there is **no theoretical guarantee** that the GLRT is a uniformly most powerful test.

- It is therefore important to be able to compute the power of the GLRT

Power of the GLRT II

- For fixed level of confidence α

$$\mathbb{P}_{H_0} (L_N > t_{\alpha}^n) \leq \alpha ,$$

the type II error **exponentially decreases to 0**.

Power of the GLRT II

- For fixed level of confidence α

$$\mathbb{P}_{H_0} (L_N > \mathbf{t}_\alpha^n) \leq \alpha ,$$

the type II error **exponentially decreases to 0**.

- Indeed, we want to evaluate

$$\begin{aligned} & \mathbb{P}_{H_1} (L_N \leq \mathbf{t}_\alpha^n) \\ &= \mathbb{P}_{H_1} \left(L_N \leq (1 + \sqrt{c})^2 + \frac{\tilde{\Theta}_N}{N^{2/3}} \mathbf{t}_\alpha^{\text{Tracy-Widom}} \right) \\ &= \mathbb{P}_{H_1} \left(L_N \leq (1 + \sqrt{c})^2 + \mathcal{O} \left(\frac{1}{N^{2/3}} \right) \right) \end{aligned}$$

Power of the GLRT II

- For fixed level of confidence α

$$\mathbb{P}_{H_0} (L_N > \mathbf{t}_\alpha^n) \leq \alpha ,$$

the type II error **exponentially decreases to 0**.

- Indeed, we want to evaluate

$$\begin{aligned} & \mathbb{P}_{H_1} (L_N \leq \mathbf{t}_\alpha^n) \\ &= \mathbb{P}_{H_1} \left(L_N \leq (1 + \sqrt{c})^2 + \frac{\tilde{\Theta}_N}{N^{2/3}} \mathbf{t}_\alpha^{\text{Tracy-Widom}} \right) \\ &= \mathbb{P}_{H_1} \left(L_N \leq (1 + \sqrt{c})^2 + \mathcal{O} \left(\frac{1}{N^{2/3}} \right) \right) \end{aligned}$$

but

$$L_N \xrightarrow[N, n \rightarrow \infty]{H_1} (1 + \mathbf{snr}) \left(1 + \frac{c}{\mathbf{snr}} \right)$$

Power of the GLRT II

- For fixed level of confidence α

$$\mathbb{P}_{H_0} (L_N > \mathbf{t}_\alpha^n) \leq \alpha ,$$

the type II error **exponentially decreases to 0**.

- Indeed, we want to evaluate

$$\begin{aligned} & \mathbb{P}_{H_1} (L_N \leq \mathbf{t}_\alpha^n) \\ &= \mathbb{P}_{H_1} \left(L_N \leq (1 + \sqrt{c})^2 + \frac{\tilde{\Theta}_N}{N^{2/3}} \mathbf{t}_\alpha^{\text{Tracy-Widom}} \right) \\ &= \mathbb{P}_{H_1} \left(L_N \leq (1 + \sqrt{c})^2 + \mathcal{O} \left(\frac{1}{N^{2/3}} \right) \right) \end{aligned}$$

but

$$L_N \xrightarrow[N, n \rightarrow \infty]{H_1} (1 + \mathbf{snr}) \left(1 + \frac{c}{\mathbf{snr}} \right) > (1 + \sqrt{c})^2 + \mathcal{O} \left(\frac{1}{N^{2/3}} \right)$$

Power of the GLRT III: How to compute type II error?

Power of the GLRT III: How to compute type II error?

Asymptotically, of course ..

Power of the GLRT III: How to compute type II error?

Asymptotically, of course .. **but how exactly?**

Power of the GLRT III: How to compute type II error?

Asymptotically, of course .. **but how exactly?**

Easy but risky: The fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_1

Power of the GLRT III: How to compute type II error?

Asymptotically, of course .. **but how exactly?**

Easy but risky: The fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_1

Theorem

$$\sqrt{N} \left(\lambda_{\max}(\hat{\mathbf{R}}_n) - (1 + \text{snr}) \left(1 + \frac{c_n}{\text{snr}} \right) \right) \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, \Gamma)$$

Power of the GLRT III: How to compute type II error?

Asymptotically, of course .. **but how exactly?**

Easy but risky: The fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_1

Theorem

$$\sqrt{N} \left(\lambda_{\max}(\hat{\mathbf{R}}_n) - (1 + \text{snr}) \left(1 + \frac{c_n}{\text{snr}} \right) \right) \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, \Gamma)$$

Then evaluate $\mathbb{P}_{H_1} (L_N \leq t_\alpha^n)$ by Gaussian quantiles.

Power of the GLRT III: How to compute type II error?

Asymptotically, of course .. **but how exactly?**

Easy but risky: The fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_1

Theorem

$$\sqrt{N} \left(\lambda_{\max}(\hat{\mathbf{R}}_n) - (1 + \text{snr}) \left(1 + \frac{c_n}{\text{snr}} \right) \right) \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, \Gamma)$$

Then evaluate $\mathbb{P}_{H_1} (L_N \leq t_\alpha^n)$ by Gaussian quantiles. Well ..

Power of the GLRT III: How to compute type II error?

Asymptotically, of course .. **but how exactly?**

Easy but risky: The fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_1

Theorem

$$\sqrt{N} \left(\lambda_{\max}(\hat{\mathbf{R}}_n) - (1 + \text{snr}) \left(1 + \frac{c_n}{\text{snr}} \right) \right) \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, \Gamma)$$

Then evaluate $\mathbb{P}_{H_1} (L_N \leq t_{\alpha}^n)$ by Gaussian quantiles. Well .. **not a very good idea** ..

Power of the GLRT III: How to compute type II error?

Asymptotically, of course .. **but how exactly?**

Easy but risky: The fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_1

Theorem

$$\sqrt{N} \left(\lambda_{\max}(\hat{\mathbf{R}}_n) - (1 + \text{snr}) \left(1 + \frac{c_n}{\text{snr}} \right) \right) \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, \Gamma)$$

Then evaluate $\mathbb{P}_{H_1} (L_N \leq t_\alpha^n)$ by Gaussian quantiles. Well .. **not a very good idea ..**

Be serious, compute the large deviations!

► one can define its error exponent \mathcal{E} as:

$$\mathcal{E} = \lim_{N, n \rightarrow \infty} -\frac{1}{n} \log \mathbb{P}_{H_1} (L_N < t_\alpha^n) .$$

Power of the GLRT III: How to compute type II error?

Asymptotically, of course .. **but how exactly?**

Easy but risky: The fluctuations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_1

Theorem

$$\sqrt{N} \left(\lambda_{\max}(\hat{\mathbf{R}}_n) - (1 + \text{snr}) \left(1 + \frac{c_n}{\text{snr}} \right) \right) \xrightarrow[N, n \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, \Gamma)$$

Then evaluate $\mathbb{P}_{H_1}(L_N \leq t_\alpha^n)$ by Gaussian quantiles. Well .. **not a very good idea** ..

Be serious, compute the large deviations!

- ▶ one can define its error exponent \mathcal{E} as:

$$\mathcal{E} = \lim_{N, n \rightarrow \infty} -\frac{1}{n} \log \mathbb{P}_{H_1}(L_N < t_\alpha^n) .$$

- ▶ Hence, the type II error writes:

$$\mathbb{P}_{H_1}(L_N < t(\alpha)) \approx_{N, n \rightarrow \infty} e^{-n\mathcal{E}}$$

Power of the GLRT IV: The error exponent

Theorem

Power of the GLRT IV: The error exponent

Theorem

- ▶ The type II error writes:

$$\mathbb{P}_{H_1}(L_N < \mathbf{t}_\alpha^n) \approx_{N, n \rightarrow \infty} e^{-n\mathcal{E}}$$

Power of the GLRT IV: The error exponent

Theorem

- The type II error writes:

$$\mathbb{P}_{H_1}(L_N < \mathbf{t}_\alpha^n) \approx_{N, n \rightarrow \infty} e^{-n\mathcal{E}}$$

in the sense that

$$\mathcal{E} = \lim_{N, n \rightarrow \infty} -\frac{1}{n} \log \mathbb{P}_{H_1}(L_N < \mathbf{t}_\alpha) .$$

Power of the GLRT IV: The error exponent

Theorem

- ▶ The type II error writes:

$$\mathbb{P}_{H_1}(L_N < \mathbf{t}_\alpha^n) \approx_{N, n \rightarrow \infty} e^{-n\mathcal{E}}$$

in the sense that

$$\mathcal{E} = \lim_{N, n \rightarrow \infty} -\frac{1}{n} \log \mathbb{P}_{H_1}(L_N < \mathbf{t}_\alpha) .$$

- ▶ The error exponent \mathcal{E} is fully explicit

$$\mathcal{E} = \frac{\lambda^+ - \lambda_{spk}^\infty}{1 + \mathbf{snr}} - (1 - c) \log \left(\frac{\lambda^+}{\lambda_{spk}^\infty} - 2c \left[F^+(\lambda^+) - F^+(\lambda_{spk}^\infty) \right] \right)$$

Power of the GLRT IV: The error exponent

Theorem

- ▶ The type II error writes:

$$\mathbb{P}_{H_1}(L_N < \mathbf{t}_\alpha^n) \approx_{N, n \rightarrow \infty} e^{-n\mathcal{E}}$$

in the sense that

$$\mathcal{E} = \lim_{N, n \rightarrow \infty} -\frac{1}{n} \log \mathbb{P}_{H_1}(L_N < \mathbf{t}_\alpha) .$$

- ▶ The error exponent \mathcal{E} is fully explicit

$$\mathcal{E} = \frac{\lambda^+ - \lambda_{spk}^\infty}{1 + \mathbf{snr}} - (1 - c) \log \left(\frac{\lambda^+}{\lambda_{spk}^\infty} - 2c \left[F^+(\lambda^+) - F^+(\lambda_{spk}^\infty) \right] \right)$$

Elements of proof

- ▶ Proof essentially based on the large deviations of $\lambda_{\max}(\hat{\mathbf{R}}_n)$ under H_1

Power of the GLRT V

The error exponent curve

Instead of letting the type I error fixed, it is of interest to let it go exponentially to zero:

$$\mathbb{P}_{H_0} (L_N > t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathbf{a}}$$

Power of the GLRT V

The error exponent curve

Instead of letting the type I error fixed, it is of interest to let it go exponentially to zero:

$$\mathbb{P}_{H_0} (L_N > t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathbf{a}}$$

and to compute the corresponding type II error (or its error component $\mathcal{E}(\mathbf{a})$)

$$\mathbb{P}_{H_1} (L_N < t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathcal{E}(\mathbf{a})}$$

Power of the GLRT V

The error exponent curve

Instead of letting the type I error fixed, it is of interest to let it go exponentially to zero:

$$\mathbb{P}_{H_0} (L_N > t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathbf{a}}$$

and to compute the corresponding type II error (or its error component $\mathcal{E}(\mathbf{a})$)

$$\mathbb{P}_{H_1} (L_N < t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathcal{E}(\mathbf{a})}$$

► The set $(\mathbf{a}, \mathcal{E}(\mathbf{a}))$ such that

$$\mathbb{P}_{H_0} (L_N > t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathbf{a}}$$

$$\mathbb{P}_{H_1} (L_N < t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathcal{E}(\mathbf{a})}$$

is called the **error exponent curve** and can be fully described in terms of large deviation rate functions.

Power of the GLRT V

The error exponent curve

Instead of letting the type I error fixed, it is of interest to let it go exponentially to zero:

$$\mathbb{P}_{H_0} (L_N > t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathbf{a}}$$

and to compute the corresponding type II error (or its error component $\mathcal{E}(\mathbf{a})$)

$$\mathbb{P}_{H_1} (L_N < t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathcal{E}(\mathbf{a})}$$

► The set $(\mathbf{a}, \mathcal{E}(\mathbf{a}))$ such that

$$\mathbb{P}_{H_0} (L_N > t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathbf{a}}$$

$$\mathbb{P}_{H_1} (L_N < t(\mathbf{a})) \approx_{N, n \rightarrow \infty} e^{-n\mathcal{E}(\mathbf{a})}$$

is called the **error exponent curve** and can be fully described in terms of large deviation rate functions.

Interest

If we want to discriminate between two tests, we can compare their error exponent curves.

Introduction

Marčenko-Pastur's theorem

Large covariance matrices

Spiked models

Statistical Test for Single-Source Detection

The setup

Asymptotic behaviour of the GLRT

Fluctuations of the test statistics

Power of the test

The GLRT: Summary

Direction of Arrival Estimation

Applications to the MIMO channel

Conclusion

Summary

- Consider the following hypothesis

$$\vec{\mathbf{y}}(k) = \begin{cases} \sigma \vec{\mathbf{w}}(k) & \text{under } H_0 \\ \vec{\mathbf{h}}_s(k) + \sigma \vec{\mathbf{w}}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

Summary

- Consider the following hypothesis

$$\vec{\mathbf{y}}(k) = \begin{cases} \sigma \vec{\mathbf{w}}(k) & \text{under } H_0 \\ \vec{\mathbf{h}}_s(k) + \sigma \vec{\mathbf{w}}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

then the GLRT amounts to study

$$T_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$$

Summary

- Consider the following hypothesis

$$\vec{\mathbf{y}}(k) = \begin{cases} \sigma \vec{\mathbf{w}}(k) & \text{under } H_0 \\ \vec{\mathbf{h}} s(k) + \sigma \vec{\mathbf{w}}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

then the GLRT amounts to study

$$T_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$$

- The test statistics T_n discriminates between H_0 and H_1 if $\text{snr} = \frac{\|\vec{\mathbf{h}}\|^2}{\sigma^2} > \sqrt{c}$

Summary

- ▶ Consider the following hypothesis

$$\vec{\mathbf{y}}(k) = \begin{cases} \sigma \vec{\mathbf{w}}(k) & \text{under } H_0 \\ \vec{\mathbf{h}} s(k) + \sigma \vec{\mathbf{w}}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

then the GLRT amounts to study

$$T_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$$

- ▶ The test statistics T_n discriminates between H_0 and H_1 if $\text{snr} = \frac{\|\vec{\mathbf{h}}\|^2}{\sigma^2} > \sqrt{c}$
- ▶ The threshold can be asymptotically determined by Tracy-Widom quantiles.

Summary

- ▶ Consider the following hypothesis

$$\vec{\mathbf{y}}(k) = \begin{cases} \sigma \vec{\mathbf{w}}(k) & \text{under } H_0 \\ \vec{\mathbf{h}} s(k) + \sigma \vec{\mathbf{w}}(k) & \text{under } H_1 \end{cases} \quad \text{for } k = 1 : n$$

then the GLRT amounts to study

$$T_n = \frac{\lambda_{\max}(\hat{\mathbf{R}}_n)}{\frac{1}{N} \text{Trace } \hat{\mathbf{R}}_n}$$

- ▶ The test statistics T_n discriminates between H_0 and H_1 if $\text{snr} = \frac{\|\vec{\mathbf{h}}\|^2}{\sigma^2} > \sqrt{c}$
- ▶ The threshold can be asymptotically determined by Tracy-Widom quantiles.
- ▶ The type II error (equivalently power of the test) can be analyzed via the error exponent of the test

$$\mathcal{E} = \lim_{N, n \rightarrow \infty} -\frac{1}{n} \log \mathbb{P}_{H_1}(L_N < t_\alpha) ,$$

which relies on the study of large deviations of λ_{\max} under H_1 .