

Introduction

Marčenko-Pastur's theorem

Large covariance matrices

Spiked models

Statistical Test for Single-Source Detection

Direction of Arrival Estimation

Position of the problem

MUSIC Algorithm

Spiked model eigenvectors

Summary

Applications to the MIMO channel

Conclusion

Spiked model eigenvectors I

- Consider the following $N \times n$ spiked model:

$$\begin{aligned}\tilde{\mathbf{X}}_N &= (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{1/2} \mathbf{X}_N \quad \text{with} \quad \|\vec{\mathbf{u}}\| = 1, \\ &= \mathbf{\Pi}^{1/2} \mathbf{X}_N\end{aligned}$$

Spiked model eigenvectors I

- Consider the following $N \times n$ spiked model:

$$\begin{aligned}\tilde{\mathbf{X}}_N &= (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{1/2} \mathbf{X}_N \quad \text{with} \quad \|\vec{\mathbf{u}}\| = 1, \\ &= \mathbf{\Pi}^{1/2} \mathbf{X}_N\end{aligned}$$

where \mathbf{X}_N has i.i.d. $0/\sigma^2$ entries.

Spiked model eigenvectors I

- ▶ Consider the following $N \times n$ spiked model:

$$\begin{aligned}\tilde{\mathbf{X}}_N &= (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{1/2} \mathbf{X}_N \quad \text{with} \quad \|\vec{\mathbf{u}}\| = 1, \\ &= \mathbf{\Pi}^{1/2} \mathbf{X}_N\end{aligned}$$

where \mathbf{X}_N has i.i.d. $0/\sigma^2$ entries.

- ▶ Let $\vec{\mathbf{v}}_{\max}$ be the eigenvector associated to λ_{\max} , the largest eigenvalue of the covariance matrix associated to $\tilde{\mathbf{X}}_N$:

$$\left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* \right) \vec{\mathbf{v}}_{\max} = \lambda_{\max} \vec{\mathbf{v}}_{\max}$$

Spiked model eigenvectors I

- ▶ Consider the following $N \times n$ spiked model:

$$\begin{aligned}\tilde{\mathbf{X}}_N &= (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{1/2} \mathbf{X}_N \quad \text{with} \quad \|\vec{\mathbf{u}}\| = 1, \\ &= \mathbf{\Pi}^{1/2} \mathbf{X}_N\end{aligned}$$

where \mathbf{X}_N has i.i.d. $0/\sigma^2$ entries.

- ▶ Let $\vec{\mathbf{v}}_{\max}$ be the eigenvector associated to λ_{\max} , the largest eigenvalue of the covariance matrix associated to $\tilde{\mathbf{X}}_N$:

$$\left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* \right) \vec{\mathbf{v}}_{\max} = \lambda_{\max} \vec{\mathbf{v}}_{\max}$$

Question

- ▶ What is the behavior of $\vec{\mathbf{v}}_{\max}$ as $N, n \rightarrow \infty$ in the regime where

$$\frac{N}{n} \rightarrow c \in (0, \infty)?$$

Spiked model eigenvectors I

- ▶ Consider the following $N \times n$ spiked model:

$$\begin{aligned}\tilde{\mathbf{X}}_N &= (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{1/2} \mathbf{X}_N \quad \text{with} \quad \|\vec{\mathbf{u}}\| = 1, \\ &= \mathbf{\Pi}^{1/2} \mathbf{X}_N\end{aligned}$$

where \mathbf{X}_N has i.i.d. $0/\sigma^2$ entries.

- ▶ Let $\vec{\mathbf{v}}_{\max}$ be the eigenvector associated to λ_{\max} , the largest eigenvalue of the covariance matrix associated to $\tilde{\mathbf{X}}_N$:

$$\left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* \right) \vec{\mathbf{v}}_{\max} = \lambda_{\max} \vec{\mathbf{v}}_{\max}$$

Question

- ▶ What is the behavior of $\vec{\mathbf{v}}_{\max}$ as $N, n \rightarrow \infty$ in the regime where

$$\frac{N}{n} \rightarrow c \in (0, \infty)?$$

Reminder

Behaviour of largest eigenvalue λ_{\max} well-understood:

Spiked model eigenvectors I

- ▶ Consider the following $N \times n$ spiked model:

$$\begin{aligned}\tilde{\mathbf{X}}_N &= (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{1/2} \mathbf{X}_N \quad \text{with} \quad \|\vec{\mathbf{u}}\| = 1, \\ &= \mathbf{\Pi}^{1/2} \mathbf{X}_N\end{aligned}$$

where \mathbf{X}_N has i.i.d. $0/\sigma^2$ entries.

- ▶ Let $\vec{\mathbf{v}}_{\max}$ be the eigenvector associated to λ_{\max} , the largest eigenvalue of the covariance matrix associated to $\tilde{\mathbf{X}}_N$:

$$\left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* \right) \vec{\mathbf{v}}_{\max} = \lambda_{\max} \vec{\mathbf{v}}_{\max}$$

Question

- ▶ What is the behavior of $\vec{\mathbf{v}}_{\max}$ as $N, n \rightarrow \infty$ in the regime where

$$\frac{N}{n} \rightarrow c \in (0, \infty)?$$

Reminder

Behaviour of largest eigenvalue λ_{\max} well-understood:

- ▶ if $\boxed{\theta \leq \sqrt{c}}$ then $\lambda_{\max} \rightarrow \sigma^2(1 + \sqrt{c})^2$, **the right edge of MP bulk.**

Spiked model eigenvectors I

- ▶ Consider the following $N \times n$ spiked model:

$$\begin{aligned}\tilde{\mathbf{X}}_N &= (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{1/2} \mathbf{X}_N \quad \text{with} \quad \|\vec{\mathbf{u}}\| = 1, \\ &= \mathbf{\Pi}^{1/2} \mathbf{X}_N\end{aligned}$$

where \mathbf{X}_N has i.i.d. $0/\sigma^2$ entries.

- ▶ Let $\vec{\mathbf{v}}_{\max}$ be the eigenvector associated to λ_{\max} , the largest eigenvalue of the covariance matrix associated to $\tilde{\mathbf{X}}_N$:

$$\left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* \right) \vec{\mathbf{v}}_{\max} = \lambda_{\max} \vec{\mathbf{v}}_{\max}$$

Question

- ▶ What is the behavior of $\vec{\mathbf{v}}_{\max}$ as $N, n \rightarrow \infty$ in the regime where

$$\frac{N}{n} \rightarrow c \in (0, \infty)?$$

Reminder

Behaviour of largest eigenvalue λ_{\max} well-understood:

- ▶ if $\theta \leq \sqrt{c}$ then $\lambda_{\max} \rightarrow \sigma^2(1 + \sqrt{c})^2$, **the right edge of MP bulk.**
- ▶ if $\theta > \sqrt{c}$ then $\lambda_{\max} \rightarrow \sigma^2(1 + \theta)(1 + c/\theta)$, i.e. λ_{\max} **separates from the bulk.**

Spiked model eigenvectors II

Preliminary observations

Spiked model eigenvectors II

Preliminary observations

1. Let N finite, $n \rightarrow \infty$, then

$$\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* = \mathbf{\Pi}^{1/2} \left(\frac{1}{n} \mathbf{X}_N \mathbf{X}_N^* \right) \mathbf{\Pi}^{1/2} \xrightarrow[n \rightarrow \infty]{} \mathbf{\Pi}$$

Spiked model eigenvectors II

Preliminary observations

1. Let N finite, $n \rightarrow \infty$, then

$$\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* = \mathbf{\Pi}^{1/2} \left(\frac{1}{n} \mathbf{X}_N \mathbf{X}_N^* \right) \mathbf{\Pi}^{1/2} \xrightarrow{n \rightarrow \infty} \mathbf{\Pi}$$

Largest eigenvalue of $\mathbf{\Pi}$ is $1 + \theta$; associated eigenvector is $\vec{\mathbf{u}}$:

$$\mathbf{\Pi} \vec{\mathbf{u}} = (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*) \vec{\mathbf{u}} = (1 + \theta) \vec{\mathbf{u}} .$$

Spiked model eigenvectors II

Preliminary observations

1. Let N finite, $n \rightarrow \infty$, then

$$\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* = \mathbf{\Pi}^{1/2} \left(\frac{1}{n} \mathbf{X}_N \mathbf{X}_N^* \right) \mathbf{\Pi}^{1/2} \xrightarrow{n \rightarrow \infty} \mathbf{\Pi}$$

Largest eigenvalue of $\mathbf{\Pi}$ is $1 + \theta$; associated eigenvector is $\vec{\mathbf{u}}$:

$$\mathbf{\Pi} \vec{\mathbf{u}} = (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*) \vec{\mathbf{u}} = (1 + \theta) \vec{\mathbf{u}} .$$

As a consequence:

$$\vec{\mathbf{v}}_{\max} \xrightarrow{n \rightarrow \infty} \vec{\mathbf{u}} .$$

Spiked model eigenvectors II

Preliminary observations

1. Let N finite, $n \rightarrow \infty$, then

$$\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* = \mathbf{\Pi}^{1/2} \left(\frac{1}{n} \mathbf{X}_N \mathbf{X}_N^* \right) \mathbf{\Pi}^{1/2} \xrightarrow{n \rightarrow \infty} \mathbf{\Pi}$$

Largest eigenvalue of $\mathbf{\Pi}$ is $1 + \theta$; associated eigenvector is $\vec{\mathbf{u}}$:

$$\mathbf{\Pi} \vec{\mathbf{u}} = (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*) \vec{\mathbf{u}} = (1 + \theta) \vec{\mathbf{u}} .$$

As a consequence:

$$\vec{\mathbf{v}}_{\max} \xrightarrow{n \rightarrow \infty} \vec{\mathbf{u}} .$$

2. If

$$N, n \rightarrow \infty , \quad \frac{N}{n} \rightarrow c ,$$

then $\boxed{\dim(\vec{\mathbf{v}}_{\max}) = N \nearrow \infty}$.

Spiked model eigenvectors II

Preliminary observations

1. Let N finite, $n \rightarrow \infty$, then

$$\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* = \mathbf{\Pi}^{1/2} \left(\frac{1}{n} \mathbf{X}_N \mathbf{X}_N^* \right) \mathbf{\Pi}^{1/2} \xrightarrow{n \rightarrow \infty} \mathbf{\Pi}$$

Largest eigenvalue of $\mathbf{\Pi}$ is $1 + \theta$; associated eigenvector is $\vec{\mathbf{u}}$:

$$\mathbf{\Pi} \vec{\mathbf{u}} = (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*) \vec{\mathbf{u}} = (1 + \theta) \vec{\mathbf{u}} .$$

As a consequence:

$$\vec{\mathbf{v}}_{\max} \xrightarrow{n \rightarrow \infty} \vec{\mathbf{u}} .$$

2. If

$$N, n \rightarrow \infty , \quad \frac{N}{n} \rightarrow c ,$$

then $\boxed{\dim(\vec{\mathbf{v}}_{\max}) = N \nearrow \infty}$. We therefore consider the projection

$$\vec{\mathbf{v}}_{\max} \vec{\mathbf{v}}_{\max}^*$$

on $\vec{\mathbf{v}}_{\max}$ of a generic deterministic vector $\vec{\mathbf{a}}_N$

Spiked model eigenvectors II

Preliminary observations

1. Let N finite, $n \rightarrow \infty$, then

$$\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* = \mathbf{\Pi}^{1/2} \left(\frac{1}{n} \mathbf{X}_N \mathbf{X}_N^* \right) \mathbf{\Pi}^{1/2} \xrightarrow{n \rightarrow \infty} \mathbf{\Pi}$$

Largest eigenvalue of $\mathbf{\Pi}$ is $1 + \theta$; associated eigenvector is $\vec{\mathbf{u}}$:

$$\mathbf{\Pi} \vec{\mathbf{u}} = (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*) \vec{\mathbf{u}} = (1 + \theta) \vec{\mathbf{u}} .$$

As a consequence:

$$\vec{\mathbf{v}}_{\max} \xrightarrow{n \rightarrow \infty} \vec{\mathbf{u}} .$$

2. If

$$N, n \rightarrow \infty , \quad \frac{N}{n} \rightarrow c ,$$

then $\boxed{\dim(\vec{\mathbf{v}}_{\max}) = N \nearrow \infty}$. We therefore consider the projection

$$\vec{\mathbf{v}}_{\max} \vec{\mathbf{v}}_{\max}^*$$

on $\vec{\mathbf{v}}_{\max}$ of a generic deterministic vector $\vec{\mathbf{a}}_N$, i.e.

$$\boxed{\vec{\mathbf{a}}_N^* \vec{\mathbf{v}}_{\max} \vec{\mathbf{v}}_{\max}^* \vec{\mathbf{a}}_N}$$

Spiked model eigenvectors III

Theorem

Let \vec{a}_N be a deterministic vector with norm 1, then

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N - \left(1 - \frac{c}{\theta^2}\right) \left(1 + \frac{c}{\theta}\right)^{-1} \vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N \xrightarrow[N, n \rightarrow \infty]{a.s.} 0 .$$

Spiked model eigenvectors III

Theorem

Let \vec{a}_N be a deterministic vector with norm 1, then

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N - \left(1 - \frac{c}{\theta^2}\right) \left(1 + \frac{c}{\theta}\right)^{-1} \vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N \xrightarrow[N, n \rightarrow \infty]{a.s.} 0 .$$

Remarks

► If N finite, $n \rightarrow \infty$, then

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N - \vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N \xrightarrow[N, n \rightarrow \infty]{a.s.} 0 .$$

Spiked model eigenvectors III

Theorem

Let \vec{a}_N be a deterministic vector with norm 1, then

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N - \left(1 - \frac{c}{\theta^2}\right) \left(1 + \frac{c}{\theta}\right)^{-1} \vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N \xrightarrow[N, n \rightarrow \infty]{a.s.} 0 .$$

Remarks

- If N finite, $n \rightarrow \infty$, then

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N - \vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N \xrightarrow[N, n \rightarrow \infty]{a.s.} 0 .$$

- The large dimension $\frac{N}{n} \rightarrow c$ induces a correction factor:

$$\boxed{\kappa(c) = \left(1 - \frac{c}{\theta^2}\right) \left(1 + \frac{c}{\theta}\right)^{-1}}$$

Spiked model eigenvectors III

Theorem

Let \vec{a}_N be a deterministic vector with norm 1, then

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N - \left(1 - \frac{c}{\theta^2}\right) \left(1 + \frac{c}{\theta}\right)^{-1} \vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N \xrightarrow[N, n \rightarrow \infty]{a.s.} 0 .$$

Remarks

- If N finite, $n \rightarrow \infty$, then

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N - \vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N \xrightarrow[N, n \rightarrow \infty]{a.s.} 0 .$$

- The large dimension $\frac{N}{n} \rightarrow c$ induces a correction factor:

$$\boxed{\kappa(c) = \left(1 - \frac{c}{\theta^2}\right) \left(1 + \frac{c}{\theta}\right)^{-1}}$$

- Of course $\kappa(c) \rightarrow 1$ if $c \rightarrow 0$.

Proof I

Reminder from complex analysis

- We need a simple result from complex analysis:

$$\frac{1}{2i\pi} \oint_{\mathcal{C}^-} \frac{dz}{z} = 1$$

if \mathcal{C}^- is a contour (take a circle of radius 1) enclosing counterclockwise 0.

Proof I

Reminder from complex analysis

- We need a simple result from complex analysis:

$$\frac{1}{2i\pi} \oint_{\mathcal{C}^-} \frac{dz}{z} = 1$$

if \mathcal{C}^- is a contour (take a circle of radius 1) enclosing counterclockwise 0.

- In particular, if \mathcal{C}^+ is a contour enclosing λ **clockwise**, then:

$$\boxed{\frac{1}{2i\pi} \oint_{\mathcal{C}^+} \frac{dz}{\lambda - z} = 1}$$

Proof I

Reminder from complex analysis

- ▶ We need a simple result from complex analysis:

$$\frac{1}{2i\pi} \oint_{\mathcal{C}^-} \frac{dz}{z} = 1$$

if \mathcal{C}^- is a contour (take a circle of radius 1) enclosing counterclockwise 0.

- ▶ In particular, if \mathcal{C}^+ is a contour enclosing λ **clockwise**, then:

$$\boxed{\frac{1}{2i\pi} \oint_{\mathcal{C}^+} \frac{dz}{\lambda - z} = 1}$$

- ▶ If \mathcal{C}^+ does not enclose λ , then the integral **equals zero**.

Proof II

Our objective

Proof II

Our objective

To express \vec{v}_{\max} with the help of the **resolvent** $\tilde{\mathbf{Q}}_n(z) = \left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* - z \mathbf{I}_N \right)^{-1}$

Proof II

Our objective

To express \vec{v}_{\max} with the help of the **resolvent** $\tilde{\mathbf{Q}}_n(z) = \left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* - z \mathbf{I}_N \right)^{-1}$

By the spectral theorem,

$$\begin{aligned} \frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* &= \mathbf{O}_N \begin{pmatrix} \lambda_{\max} & & \\ & \ddots & \\ & & \lambda_N \end{pmatrix} \mathbf{O}_N^* \\ &= [\vec{v}_{\max} \ \mathbf{O}_{N-1}] \begin{pmatrix} \lambda_{\max} & & \\ & \ddots & \\ & & \lambda_N \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \end{aligned}$$

Proof II

Our objective

To express \vec{v}_{\max} with the help of the **resolvent** $\tilde{\mathbf{Q}}_n(z) = \left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* - z \mathbf{I}_N \right)^{-1}$

By the spectral theorem,

$$\begin{aligned} \frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* &= \mathbf{O}_N \begin{pmatrix} \lambda_{\max} & & \\ & \ddots & \\ & & \lambda_N \end{pmatrix} \mathbf{O}_N^* \\ &= [\vec{v}_{\max} \ \mathbf{O}_{N-1}] \begin{pmatrix} \lambda_{\max} & & \\ & \ddots & \\ & & \lambda_N \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \end{aligned}$$

In particular,

$$\left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* - z \mathbf{I}_N \right)^{-1} = [\vec{v}_{\max} \ \mathbf{O}_{N-1}] \begin{pmatrix} \frac{1}{\lambda_{\max} - z} & & \\ & \ddots & \\ & & \frac{1}{\lambda_N - z} \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix}$$

Recall that

- if $\theta > \sqrt{c}$, λ_{\max} separates from the bulk

Proof II

Our objective

To express \vec{v}_{\max} with the help of the **resolvent** $\tilde{\mathbf{Q}}_n(z) = \left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* - z \mathbf{I}_N \right)^{-1}$

By the spectral theorem,

$$\begin{aligned} \frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* &= \mathbf{O}_N \begin{pmatrix} \lambda_{\max} & & \\ & \ddots & \\ & & \lambda_N \end{pmatrix} \mathbf{O}_N^* \\ &= [\vec{v}_{\max} \ \mathbf{O}_{N-1}] \begin{pmatrix} \lambda_{\max} & & \\ & \ddots & \\ & & \lambda_N \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \end{aligned}$$

In particular,

$$\left(\frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^* - z \mathbf{I}_N \right)^{-1} = [\vec{v}_{\max} \ \mathbf{O}_{N-1}] \begin{pmatrix} \frac{1}{\lambda_{\max} - z} & & \\ & \ddots & \\ & & \frac{1}{\lambda_N - z} \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix}$$

Recall that

► if $\theta > \sqrt{c}$, λ_{\max} separates from the bulk
and consider a contour \mathcal{C}^+ **exclusively enclosing** the eigenvalue λ_{\max} .

Proof III

We have

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz$$

Proof III

We have

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz$$

Indeed,

$$\frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz$$

Proof III

We have

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz$$

Indeed,

$$\begin{aligned} & \frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz \\ &= \frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* [\vec{v}_{\max} \mathbf{O}_{N-1}] \begin{pmatrix} \frac{1}{\lambda_{\max} - z} & & \\ & \ddots & \\ & & \frac{1}{\lambda_N - z} \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \vec{a}_N dz \end{aligned}$$

Proof III

We have

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz$$

Indeed,

$$\begin{aligned} & \frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz \\ &= \frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* [\vec{v}_{\max} \mathbf{O}_{N-1}] \begin{pmatrix} \frac{1}{\lambda_{\max}-z} & & \\ & \ddots & \\ & & \frac{1}{\lambda_N-z} \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \vec{a}_N dz \\ &= \vec{a}_N^* [\vec{v}_{\max} \mathbf{O}_{N-1}] \begin{pmatrix} \frac{1}{2i\pi} \oint \frac{1}{\lambda_{\max}-z} dz & & \\ & \ddots & \\ & & \frac{1}{2i\pi} \oint \frac{1}{\lambda_N-z} dz \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \vec{a}_N \end{aligned}$$

Proof III

We have

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz$$

Indeed,

$$\begin{aligned} & \frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz \\ &= \frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* [\vec{v}_{\max} \mathbf{O}_{N-1}] \begin{pmatrix} \frac{1}{\lambda_{\max}-z} & & \\ & \ddots & \\ & & \frac{1}{\lambda_N-z} \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \vec{a}_N dz \\ &= \vec{a}_N^* [\vec{v}_{\max} \mathbf{O}_{N-1}] \begin{pmatrix} \frac{1}{2i\pi} \oint \frac{1}{\lambda_{\max}-z} dz & & \\ & \ddots & \\ & & \frac{1}{2i\pi} \oint \frac{1}{\lambda_N-z} dz \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \vec{a}_N \\ &= \vec{a}_N^* [\vec{v}_{\max} \mathbf{O}_{N-1}] \begin{pmatrix} 1 & & \\ & \ddots & \\ & & 0 \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \vec{a}_N \end{aligned}$$

Proof III

We have

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz$$

Indeed,

$$\begin{aligned} & \frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz \\ &= \frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* [\vec{v}_{\max} \mathbf{O}_{N-1}] \begin{pmatrix} \frac{1}{\lambda_{\max}-z} & & \\ & \ddots & \\ & & \frac{1}{\lambda_N-z} \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \vec{a}_N dz \\ &= \vec{a}_N^* [\vec{v}_{\max} \mathbf{O}_{N-1}] \begin{pmatrix} \frac{1}{2i\pi} \oint \frac{1}{\lambda_{\max}-z} dz & & \\ & \ddots & \\ & & \frac{1}{2i\pi} \oint \frac{1}{\lambda_N-z} dz \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \vec{a}_N \\ &= \vec{a}_N^* [\vec{v}_{\max} \mathbf{O}_{N-1}] \begin{pmatrix} 1 & & \\ & \ddots & \\ & & 0 \end{pmatrix} \begin{bmatrix} \vec{v}_{\max}^* \\ \mathbf{O}_{N-1}^* \end{bmatrix} \vec{a}_N \\ &= \vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N . \end{aligned}$$

Proof IV

Recall

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* \tilde{\mathbf{Q}}_n(z) \vec{a}_N dz$$

and temporarily forget about the integral.

Proof IV

Recall

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* \tilde{\mathbf{Q}}_n(z) \vec{a}_N dz$$

and temporarily forget about the integral.

► Our objective now is:

to find a new formulation of $\vec{a}_N^* \tilde{\mathbf{Q}}_n(z) \vec{a}_N$ and clearly separate the contribution from the perturbation (\vec{u} and θ) and the resolvent $\mathbf{Q}_n(z)$ from the non-perturbed model.

Proof IV

Recall

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* \tilde{\mathbf{Q}}_n(z) \vec{a}_N dz$$

and temporarily forget about the integral.

► Our objective now is:

to find a new formulation of $\vec{a}_N^* \tilde{\mathbf{Q}}_n(z) \vec{a}_N$ and clearly separate the contribution from the perturbation (\vec{u} and θ) and the resolvent $\mathbf{Q}_n(z)$ from the non-perturbed model.

Introduce the notations

$$\mathbf{Z}_N = \frac{1}{n} \mathbf{X}_N \mathbf{X}_N^* \quad \text{and} \quad \tilde{\mathbf{Z}}_N = \frac{1}{n} \tilde{\mathbf{X}}_N \tilde{\mathbf{X}}_N^*$$

and recall the formula for the inverse of a rank-one perturbation:

$$(\mathbf{A} + \vec{u} \vec{u}^*)^{-1} = \mathbf{A}^{-1} - \frac{\mathbf{A}^{-1} \vec{u} \vec{u}^* \mathbf{A}^{-1}}{1 + \vec{u} \mathbf{A} \vec{u}^*},$$

In particular

$$\mathbf{\Pi}^{-1} = (\mathbf{I}_N + \theta \vec{u} \vec{u}^*)^{-1} = \mathbf{I}_N - \frac{\theta}{1 + \theta} \vec{u} \vec{u}^*$$

Proof V

$$\tilde{\mathbf{Q}}_n(z) = \left(\mathbf{\Pi}^{1/2} \tilde{\mathbf{Z}}_N \mathbf{\Pi}^{1/2} - z \mathbf{I}_N \right)^{-1}$$

Proof V

$$\begin{aligned}\tilde{\mathbf{Q}}_n(z) &= \left(\mathbf{\Pi}^{1/2} \tilde{\mathbf{Z}}_N \mathbf{\Pi}^{1/2} - z \mathbf{I}_N \right)^{-1} \\ &= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \mathbf{\Pi}^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2}\end{aligned}$$

Proof V

$$\begin{aligned}\tilde{\mathbf{Q}}_n(z) &= \left(\mathbf{\Pi}^{1/2} \tilde{\mathbf{Z}}_N \mathbf{\Pi}^{1/2} - z \mathbf{I}_N \right)^{-1} \\ &= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \mathbf{\Pi}^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\ &= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2}\end{aligned}$$

Proof V

$$\begin{aligned}\tilde{\mathbf{Q}}_n(z) &= \left(\mathbf{\Pi}^{1/2} \tilde{\mathbf{Z}}_N \mathbf{\Pi}^{1/2} - z \mathbf{I}_N \right)^{-1} \\ &= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \mathbf{\Pi}^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\ &= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\ &= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \left(\mathbf{I}_N - \frac{\theta}{1 + \theta} \vec{\mathbf{u}} \vec{\mathbf{u}}^* \right) \right)^{-1} \mathbf{\Pi}^{-1/2}\end{aligned}$$

Proof V

$$\begin{aligned}
\tilde{\mathbf{Q}}_n(z) &= \left(\mathbf{\Pi}^{1/2} \tilde{\mathbf{Z}}_N \mathbf{\Pi}^{1/2} - z \mathbf{I}_N \right)^{-1} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \mathbf{\Pi}^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \left(\mathbf{I}_N - \frac{\theta}{1 + \theta} \vec{\mathbf{u}} \vec{\mathbf{u}}^* \right) \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} (\mathbf{Z}_N - z \mathbf{I}_N + \xi \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \mathbf{\Pi}^{-1/2} \quad \text{where } \xi = z \frac{\theta}{1 + \theta}
\end{aligned}$$

Proof V

$$\begin{aligned}
\tilde{\mathbf{Q}}_n(z) &= \left(\mathbf{\Pi}^{1/2} \tilde{\mathbf{Z}}_N \mathbf{\Pi}^{1/2} - z \mathbf{I}_N \right)^{-1} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \mathbf{\Pi}^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \left(\mathbf{I}_N - \frac{\theta}{1 + \theta} \vec{\mathbf{u}} \vec{\mathbf{u}}^* \right) \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \mathbf{I}_N + \xi \vec{\mathbf{u}} \vec{\mathbf{u}}^* \right)^{-1} \mathbf{\Pi}^{-1/2} \quad \text{where } \xi = z \frac{\theta}{1 + \theta} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Q}_n - \frac{\mathbf{Q}_n \xi \vec{\mathbf{u}} \vec{\mathbf{u}}^* \mathbf{Q}_n}{1 + \xi \vec{\mathbf{u}}^* \mathbf{Q}_n \vec{\mathbf{u}}} \right) \mathbf{\Pi}^{-1/2}
\end{aligned}$$

Proof V

$$\begin{aligned}
\tilde{\mathbf{Q}}_n(z) &= \left(\mathbf{\Pi}^{1/2} \tilde{\mathbf{Z}}_N \mathbf{\Pi}^{1/2} - z \mathbf{I}_N \right)^{-1} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \mathbf{\Pi}^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \left(\mathbf{I}_N - \frac{\theta}{1 + \theta} \vec{\mathbf{u}} \vec{\mathbf{u}}^* \right) \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} (\mathbf{Z}_N - z \mathbf{I}_N + \xi \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \mathbf{\Pi}^{-1/2} \quad \text{where } \xi = z \frac{\theta}{1 + \theta} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Q}_n - \frac{\mathbf{Q}_n \xi \vec{\mathbf{u}} \vec{\mathbf{u}}^* \mathbf{Q}_n}{1 + \xi \vec{\mathbf{u}}^* \mathbf{Q}_n \vec{\mathbf{u}}} \right) \mathbf{\Pi}^{-1/2}
\end{aligned}$$

Hence

$$\vec{\mathbf{a}}_N^* \tilde{\mathbf{Q}}_n(z) \vec{\mathbf{a}}_N = \vec{\mathbf{a}}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n(z) \mathbf{\Pi}^{1/2} \vec{\mathbf{a}}_N - \xi \frac{\vec{\mathbf{a}}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n \vec{\mathbf{u}} \vec{\mathbf{u}}^* \mathbf{Q}_n \mathbf{\Pi}^{1/2} \vec{\mathbf{a}}_N}{1 + \xi \vec{\mathbf{u}}^* \mathbf{Q}_n \vec{\mathbf{u}}}$$

Proof V

$$\begin{aligned}
\tilde{\mathbf{Q}}_n(z) &= \left(\mathbf{\Pi}^{1/2} \tilde{\mathbf{Z}}_N \mathbf{\Pi}^{1/2} - z \mathbf{I}_N \right)^{-1} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \mathbf{\Pi}^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \left(\mathbf{I}_N - \frac{\theta}{1 + \theta} \vec{\mathbf{u}} \vec{\mathbf{u}}^* \right) \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} (\mathbf{Z}_N - z \mathbf{I}_N + \xi \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \mathbf{\Pi}^{-1/2} \quad \text{where } \xi = z \frac{\theta}{1 + \theta} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Q}_n - \frac{\mathbf{Q}_n \xi \vec{\mathbf{u}} \vec{\mathbf{u}}^* \mathbf{Q}_n}{1 + \xi \vec{\mathbf{u}}^* \mathbf{Q}_n \vec{\mathbf{u}}} \right) \mathbf{\Pi}^{-1/2}
\end{aligned}$$

Hence

$$\vec{\mathbf{a}}_N^* \tilde{\mathbf{Q}}_n(z) \vec{\mathbf{a}}_N = \vec{\mathbf{a}}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n(z) \mathbf{\Pi}^{1/2} \vec{\mathbf{a}}_N - \xi \frac{\vec{\mathbf{a}}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n \vec{\mathbf{u}} \vec{\mathbf{u}}^* \mathbf{Q}_n \mathbf{\Pi}^{1/2} \vec{\mathbf{a}}_N}{1 + \xi \vec{\mathbf{u}}^* \mathbf{Q}_n \vec{\mathbf{u}}}$$

Not so ugly!

Proof V

$$\begin{aligned}
\tilde{\mathbf{Q}}_n(z) &= \left(\mathbf{\Pi}^{1/2} \tilde{\mathbf{Z}}_N \mathbf{\Pi}^{1/2} - z \mathbf{I}_N \right)^{-1} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \mathbf{\Pi}^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z (\mathbf{I}_N + \theta \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Z}_N - z \left(\mathbf{I}_N - \frac{\theta}{1 + \theta} \vec{\mathbf{u}} \vec{\mathbf{u}}^* \right) \right)^{-1} \mathbf{\Pi}^{-1/2} \\
&= \mathbf{\Pi}^{-1/2} (\mathbf{Z}_N - z \mathbf{I}_N + \xi \vec{\mathbf{u}} \vec{\mathbf{u}}^*)^{-1} \mathbf{\Pi}^{-1/2} \quad \text{where } \xi = z \frac{\theta}{1 + \theta} \\
&= \mathbf{\Pi}^{-1/2} \left(\mathbf{Q}_n - \frac{\mathbf{Q}_n \xi \vec{\mathbf{u}} \vec{\mathbf{u}}^* \mathbf{Q}_n}{1 + \xi \vec{\mathbf{u}}^* \mathbf{Q}_n \vec{\mathbf{u}}} \right) \mathbf{\Pi}^{-1/2}
\end{aligned}$$

Hence

$$\vec{\mathbf{a}}_N^* \tilde{\mathbf{Q}}_n(z) \vec{\mathbf{a}}_N = \vec{\mathbf{a}}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n(z) \mathbf{\Pi}^{1/2} \vec{\mathbf{a}}_N - \xi \frac{\vec{\mathbf{a}}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n \vec{\mathbf{u}} \vec{\mathbf{u}}^* \mathbf{Q}_n \mathbf{\Pi}^{1/2} \vec{\mathbf{a}}_N}{1 + \xi \vec{\mathbf{u}}^* \mathbf{Q}_n \vec{\mathbf{u}}}$$

Not so ugly! And we have separated the contribution of the perturbation from the non-perturbated model.

Proof VI

Recall

$$\vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N = \vec{a}_N^* \Pi^{1/2} Q_n(z) \Pi^{1/2} \vec{a}_N - \xi \frac{\vec{a}_N^* \Pi^{1/2} Q_n \vec{u} \vec{u}^* Q_n \Pi^{1/2} \vec{a}_N}{1 + \xi \vec{u}^* Q_n \vec{u}}$$

Proof VI

Recall

$$\vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N = \vec{a}_N^* \Pi^{1/2} Q_n(z) \Pi^{1/2} \vec{a}_N - \xi \frac{\vec{a}_N^* \Pi^{1/2} Q_n \vec{u} \vec{u}^* Q_n \Pi^{1/2} \vec{a}_N}{1 + \xi \vec{u}^* Q_n \vec{u}}$$

and integrate the first term

$$\frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* \Pi^{1/2} Q_n(z) \Pi^{1/2} \vec{a}_N = ??$$

Proof VI

Recall

$$\vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N = \vec{a}_N^* \Pi^{1/2} Q_n(z) \Pi^{1/2} \vec{a}_N - \xi \frac{\vec{a}_N^* \Pi^{1/2} Q_n \vec{u} \vec{u}^* Q_n \Pi^{1/2} \vec{a}_N}{1 + \xi \vec{u}^* Q_n \vec{u}}$$

and integrate the first term

$$\frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \Pi^{1/2} Q_n(z) \Pi^{1/2} \vec{a}_N = \mathbf{0}$$

Why?

Proof VI

Recall

$$\vec{a}_N^* \tilde{\mathbf{Q}}_n(z) \vec{a}_N = \vec{a}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n(z) \mathbf{\Pi}^{1/2} \vec{a}_N - \xi \frac{\vec{a}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n \vec{u} \vec{u}^* \mathbf{Q}_n \mathbf{\Pi}^{1/2} \vec{a}_N}{1 + \xi \vec{u}^* \mathbf{Q}_n \vec{u}}$$

and integrate the first term

$$\frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n(z) \mathbf{\Pi}^{1/2} \vec{a}_N = 0$$

Why? Because

1. the contour only encloses $\lambda_{\max}(\tilde{\mathbf{Z}}_n)$ which is away from the bulk,
2. but all the eigenvalues of \mathbf{Z}_n are **in the bulk**. Hence:

$$\frac{1}{2i\pi} \oint_{C^+} \frac{1}{\lambda_i(\mathbf{Z}_n) - z} dz = 0.$$

Proof VI

Recall

$$\vec{a}_N^* \tilde{\mathbf{Q}}_n(z) \vec{a}_N = \vec{a}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n(z) \mathbf{\Pi}^{1/2} \vec{a}_N - \xi \frac{\vec{a}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n \vec{u} \vec{u}^* \mathbf{Q}_n \mathbf{\Pi}^{1/2} \vec{a}_N}{1 + \xi \vec{u}^* \mathbf{Q}_n \vec{u}}$$

and integrate the first term

$$\frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n(z) \mathbf{\Pi}^{1/2} \vec{a}_N = 0$$

Why? Because

1. the contour only encloses $\lambda_{\max}(\tilde{\mathbf{Z}}_n)$ which is away from the bulk,
2. but all the eigenvalues of \mathbf{Z}_n are **in the bulk**. Hence:

$$\frac{1}{2i\pi} \oint_{C^+} \frac{1}{\lambda_i(\mathbf{Z}_n) - z} dz = 0.$$

Last step is to simplify the remaining expression by systematically use the **isotropic MP law**:

$$\vec{c}^* \mathbf{Q}_n(z) \vec{d} - \vec{c}^* \vec{d} \mathbf{g}_{\check{\text{MP}}}(z) \xrightarrow[N, n \rightarrow \infty]{a.s.} 0$$

Proof VI

Recall

$$\vec{a}_N^* \tilde{\mathbf{Q}}_n(z) \vec{a}_N = \vec{a}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n(z) \mathbf{\Pi}^{1/2} \vec{a}_N - \xi \frac{\vec{a}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n \vec{u} \vec{u}^* \mathbf{Q}_n \mathbf{\Pi}^{1/2} \vec{a}_N}{1 + \xi \vec{u}^* \mathbf{Q}_n \vec{u}}$$

and integrate the first term

$$\frac{1}{2i\pi} \oint_{C^+} \vec{a}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n(z) \mathbf{\Pi}^{1/2} \vec{a}_N = 0$$

Why? Because

1. the contour only encloses $\lambda_{\max}(\tilde{\mathbf{Z}}_n)$ which is away from the bulk,
2. but all the eigenvalues of \mathbf{Z}_n are **in the bulk**. Hence:

$$\frac{1}{2i\pi} \oint_{C^+} \frac{1}{\lambda_i(\mathbf{Z}_n) - z} dz = 0.$$

Last step is to simplify the remaining expression by systematically use the **Isotropic MP law**:

$$\boxed{\vec{c}^* \mathbf{Q}_n(z) \vec{d} - \vec{c}^* \vec{d} \mathbf{g}_{\check{\text{MP}}}(z) \xrightarrow[N, n \rightarrow \infty]{a.s.} 0} \Rightarrow \begin{cases} \vec{a}_N^* \mathbf{\Pi}^{1/2} \mathbf{Q}_n \vec{u} & \approx & \vec{a}_N^* \mathbf{\Pi}^{1/2} \vec{u} \mathbf{g}_{\check{\text{MP}}}(z) \\ \vec{u}^* \mathbf{Q}_n \mathbf{\Pi}^{1/2} \vec{a}_N & \approx & \vec{u}^* \mathbf{\Pi}^{1/2} \vec{a}_N \mathbf{g}_{\check{\text{MP}}}(z) \\ \vec{u}^* \mathbf{Q}_n \vec{u} & \approx & \mathbf{g}_{\check{\text{MP}}}(z) \end{cases}$$

Proof VII

After simplifications,

$$\begin{aligned}
 \vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N &\approx -\frac{1}{2i\pi} \oint_{C^+} |\vec{a}_N^* \mathbf{\Pi}^{1/2} \vec{u}|^2 \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz \\
 &= -\frac{\vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz
 \end{aligned}$$

Proof VII

After simplifications,

$$\begin{aligned}\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N &\approx -\frac{1}{2i\pi} \oint_{C^+} |\vec{a}_N^* \mathbf{\Pi}^{1/2} \vec{u}|^2 \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz \\ &= -\frac{\vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz\end{aligned}$$

It remains to compute the correction factor

$$-\frac{1}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz$$

by residue calculus (not that difficult).

Proof VII

After simplifications,

$$\begin{aligned}\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N &\approx -\frac{1}{2i\pi} \oint_{C^+} |\vec{a}_N^* \mathbf{\Pi}^{1/2} \vec{u}|^2 \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz \\ &= -\frac{\vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz\end{aligned}$$

It remains to compute the correction factor

$$-\frac{1}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz$$

by residue calculus (not that difficult).

A **minor miracle** occurs:

Proof VII

After simplifications,

$$\begin{aligned}\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N &\approx -\frac{1}{2i\pi} \oint_{C^+} |\vec{a}_N^* \mathbf{\Pi}^{1/2} \vec{u}|^2 \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz \\ &= -\frac{\vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz\end{aligned}$$

It remains to compute the correction factor

$$-\frac{1}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz$$

by residue calculus (not that difficult).

A **minor miracle** occurs: **This factor admits a closed form formula!**

Proof VII

After simplifications,

$$\begin{aligned}\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N &\approx -\frac{1}{2i\pi} \oint_{C^+} |\vec{a}_N^* \Pi^{1/2} \vec{u}|^2 \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz \\ &= -\frac{\vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz\end{aligned}$$

It remains to compute the correction factor

$$-\frac{1}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz$$

by residue calculus (not that difficult).

A **minor miracle** occurs: **This factor admits a closed form formula!**

$$-\frac{1}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz = \left(1 - \frac{c}{\theta^2}\right) \left(1 + \frac{c}{\theta}\right)^{-1}$$

Proof VII

After simplifications,

$$\begin{aligned}\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N &\approx -\frac{1}{2i\pi} \oint_{C^+} |\vec{a}_N^* \Pi^{1/2} \vec{u}|^2 \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz \\ &= -\frac{\vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz\end{aligned}$$

It remains to compute the correction factor

$$-\frac{1}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz$$

by residue calculus (not that difficult).

A **minor miracle** occurs: **This factor admits a closed form formula!**

$$-\frac{1}{1 + \theta} \oint_{C^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz = \left(1 - \frac{c}{\theta^2}\right) \left(1 + \frac{c}{\theta}\right)^{-1}$$

Finally:

$$\boxed{\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N - \left(1 - \frac{c}{\theta^2}\right) \left(1 + \frac{c}{\theta}\right)^{-1} \vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N \xrightarrow[N, n \rightarrow \infty]{a.s.} 0 .}$$

Introduction

Marčenko-Pastur's theorem

Large covariance matrices

Spiked models

Statistical Test for Single-Source Detection

Direction of Arrival Estimation

Position of the problem

MUSIC Algorithm

Spiked model eigenvectors

Summary

Applications to the MIMO channel

Conclusion

Summary

1. Expression of \vec{v}_{\max} with the help of the resolvent

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz$$

Summary

1. Expression of \vec{v}_{\max} with the help of the resolvent

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz$$

2. Convenient expression of \vec{v}_{\max} where the contribution of the perturbation is separated from the resolvent of the non-perturbated model ($\check{\text{MP}}$)

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N \approx -\frac{\vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N}{1 + \theta} \oint_{\mathcal{C}^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz$$

Summary

1. Expression of \vec{v}_{\max} with the help of the resolvent

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N = \frac{1}{2i\pi} \oint_{\mathcal{C}^+} \vec{a}_N^* \tilde{Q}_n(z) \vec{a}_N dz$$

2. Convenient expression of \vec{v}_{\max} where the contribution of the perturbation is separated from the resolvent of the non-perturbated model ($\check{\text{MP}}$)

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N \approx - \frac{\vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N}{1 + \theta} \oint_{\mathcal{C}^+} \frac{\mathbf{g}_{\check{\text{MP}}}^2(z)}{\xi^{-1} + \mathbf{g}_{\check{\text{MP}}}(z)} dz$$

3. Residue calculus to find the final form

$$\vec{a}_N^* \vec{v}_{\max} \vec{v}_{\max}^* \vec{a}_N - \left(1 - \frac{c}{\theta^2}\right) \left(1 + \frac{c}{\theta}\right)^{-1} \vec{a}_N^* \vec{u} \vec{u}^* \vec{a}_N \xrightarrow[N, n \rightarrow \infty]{a.s.} 0 .$$