

Pharmacy Drug Policy & Procedure

Policy Name: Vyndaqel/Vyndamax (tafamidis meglumine) Policy#: 2714P

Purpose of the Policy

The purpose of this policy is to establish the criteria for coverage of Vyndaqel or Vyndamax.

Statement of the Policy

Health Alliance Medical Plans will approve the use of Vyndaqel (tafamidis meglumine) or Vyndamax (tafarnidis meglumine) under the specialty pharmacy benefit when the following criteria have been met.

Criteria

1. Coverage Criteria

- 1.1 Diagnosis of transthyretin (ATIR) mediated amyloidosis with cardiomyopathy (ATIR-CM)
- 1.2 One of the following:
 - Documentation that the patient has a pathogenic TIR mutation (e.g., V30M), or
 - Cardiac or non-cardiac tissue biopsy demonstrating histologic confirmation of ATIR amyloid deposits, or
 - ALL of the following:
 - Echocardiogram or cardiac magnetic resonance imaging suggestive of amyloidosis, and
 - Radionuclide imaging (99mTc-DPD, 99mTc-PYP, or 99m Tc-HMDP) showing grade 2 or 3 cardiac uptake, and
 - Absence of monoclonal protein identified in serum, urine immunofixation (JFE), serum free light chain (sFLC) assay
- 1.3 Prescribed by or in consultation with a cardiologist (heart doctor)
- 1.4 Presence of clinical signs and symptoms of cardiomyopathy (e.g., heart failure, dyspnea, edema, hepatomegaly, ascites, angina, etc.)
- 1.5 Documentation of BOTH of the following:
 - One of the following:
 - Patient has New York Heart Association (NYHA) Functional Class I or II heart failure, or
 - Patient has New York Heart Association (NYHA) Functional Class III heart failure, and patient's cardiopulmonary functional status allows patient to ambulate 100 meters or greater in 6 minutes or less
 - Patient has an N-terminal pro-B-type naturetic peptide (NT-proBNP) level greater than or equal to 600 pg/mL
- 1.6 Documentation of previous trial and failure or contraindication to Attruby (acoramidis)

2. Exclusion Criteria

2.1 Concomitant use with any other disedase modifying therapy for ATIR-CM

3. Approval Period

- 3.1 Initial Approval: 12 months
- 3.2 Reapproval: 12 months with documentation that the patient has experienced a positive clinical response to VyndaqeWyndamax (e.g., improved symptoms, quality of life, slowing of disease progression, decreased hospitalizations, etc.)

References

- 1. Vyndaqel (tafamidis meglumine) and Vyndamax (tafamidis) [prescribing information]. New York, NY: Pfizer Labs; October 2023.
- 2. Maurer MS, Schwartz JH, Gundapaneni B, et al. Tafamidis Treatment for Patients with Transthyretin Amyloid Cardiomyopathy. N Engl J Med 2018; 379:1007.
- 3. Bulawa CE, Connelly S, Devit M, et al. Tafamidis, a potent and selective transthyretin kinetic stabilizer that

inhibits the amyloid cascade. Proc Natl Acad Sci US A 2012; 109:9629.

4. Kittleson MM, Maurer MS, Ambardekar AV, et al; American Heart Association Heart Failure and Transplantation Committee of the Council on Clinical Cardiology. Cardiac Amyloidosis: Evolving Diagnosis and Management: A Scientific Statement From the American Heart Association. Circulation. 2020 Jul 7;142(1):e7-e22.

Created Date: 12/04/19 Effective Date: 12/04/19 Posted to Website: 12/04/19 Revision Date: 06/11/25

DISCLAIMER

This Medical Policy has been developed as a guide for determining medical necessity. The process of medical necessity review also entails review of the most recent literature and physician review. Medical Policy is not intended to dictate to providers how to practice medicine. Providers are expected to exercise their medical judgment in providing the most appropriate care. Health Alliance encourages input from providers when developing and implementing medical policies. Benefit determinations are based on applicable contract language in the member's Policy/ Subscription Certificate/ Summary Plan Description. This Medical Policy does not guarantee coverage. There may be a delay between the revision of this policy and the posting on the web. Please contact the Health Alliance Customer Service Department at 1-800-851-3379 for verification of coverage.