

Rinvoq

Pharmacy Drug Policy Checklist

POLICY NAME Tocilizumab Products - Pharmacy benefit POLICY # 1836P

Criteria

Coverage Criteria for Rheumatoid Arthritis (RA) See Rheumatoid Arthritis Immunomodulator Therapies policy Coverage Criteria for Polyarticular Juvenile Idiopathic Arthritis (PJIA)				
				See Polyarticular Juvenile Idiopathic Arthritis Immunomodulator Therapies policy
			Cov	erage Criteria for Systemic Juvenile Idiopathic Arthritis (SJIA)
	Diagnosis of Systemic Juvenile Idiopathic Arthritis (SiIA)			
	Ordered by a Rheumatologist (musculoskeletal doctor)			
	Documentation to support ONE of the following:			
	 Documented trial and failure of one non-steroidal anti-inflammatory drug (NSAID, such as ibuprofen or naproxen) for at least 2 weeks 			
	 Documentation the patient has moderate-to-severe disease including any one of the following systemic manifestations: 			
	• Fever			
	SerositisEarly Macrophage Activation Syndrome (MAS)			
Cov	erage Criteria for Giant Cell Arteritis			
	Diagnosis of Giant Cell Arteritis			
	Ordered by a Rheumatologist (musculoskeletal doctor), Ophthalmologist (eye doctor), or Neuro-Ophthalmologist (doctor of the eyes and nervous system)			
	Documented failure to respond to a minimum 3-month trial of glucocorticoids			

Documented failure to respond to a minimum 3-month trial, intolerance, or contraindication to

Coverage Criteria for Systemic Sclerosis-Associated Interstitial Lung Disease (SSc-ILD)		
	Diagnosis of Systemic Sclerosis-Associated Interstitial Lung Disease (SSc-ILD)	
	Age 18 years or older	
	Ordered by or in consultation with a pulmonologist (lung doctor) or rheumatologist (musculoskeletal doctor)	
	Documented trial and subsequent failure or contraindication to mycophenolate mofetil or cyclophosphamide	
	Only subcutaneous, not IV, Actemra will be used for this indication	
	Medication will not be used in combination with Ofev	
	Medication will not be used in combination with other immunomodulators	
Exclusion Criteria – Any of the following prevents coverage		
	Inadequate response to initial or previous tocilizumab therapy	
	Health Alliance does not cover concurrent therapy with other immunomodulators based upon the possible increased risk for infections and other potential pharmacological interactions	