

Pharmacy Drug Policy Checklist

POLICY NAME Illaris (canakinumab) POLICY # 2388P

Criteria

Coverage Criteria for Periodic Fever Syndromes		
	 1.1 Diagnosis of one of the following Periodic Fever Syndromes Cryopyrin-Associated Periodic Syndromes (CAPS) Familial Mediterranean Fever Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD) Tumor Necrosis Factor (TNF) Receptor-Associated Periodic Syndrome 	
	1.2 Age 4 years or older	
Coverage Criteria for Systemic Juvenile Idiopathic Arthritis (SJIA)		
	2.1 Diagnosis of Systemic Juvenile Idiopathic Arthritis (SJIA)	
	2.2 Age 2 years or older	
	2.3 Ordered by a Rheumatologist (musculoskeletal doctor)	
	 2.4 Documentation to support one of following: Documented trial and failure of one non-steroidal anti-inflammatory drug (NSAID, such as ibuprofen or naproxen) for at least 2 weeks Documentation the patient has moderate-to-severe disease including any one of the following symptoms: ? Fever ? Serositis (tissue inflammation) ? Early Macrophage Activation Syndrome (MAS) 	

3.1 Diagnosis of Adult-Onset Still's Disease (AOSD) that is considered to be moderate to severe
in nature

- 3.2 Documentation the patient has moderate-to-severe disease including any one of the following symptoms:
 - Fever
 - Rash

• Arthritis or arthralgia (joint aches or inflammation)

Coverage Criteria for Adult-Onset Still's Disease (AOSD)

3.3 Ordered by a Rheumatologist (musculoskeletal doctor)

	erosions:	
	 Documented failure to respond, intolerance, or contraindication to non-steroidal anti- inflammatory drugs (NSAIDs, such as ibuprofen or naproxen) 	
	• Documented failure to respond, intolerance, or contraindication to glucocorticoids ? Failure to respond defined as requiring prednisone greater than 10mg daily to remain symptoms free	
	 3.5 For patients with severe disease such as life-threatening organ involvement and/or conditions such as severe hepatic (liver) involvement, cardiac tamponade (serious medical condition in which blood or fluids fill the space between the sac that encases the heart and the heart muscle), and/or disseminated intravascular coagulation (condition in which blood clots form throughout the body, blocking small blood vessels); Documented failure to respond, intolerance, or contraindication to glucocorticoids 	
Coverage Criteria for Gout Flares		
	4.1 Diagnosis of gout flares	
	4.2 Prescribed by or in consultation with a rheumatologist	
	4.3 Age 18 years or older	
	 4.4 Trial and failure contraindication, or intolerance to ALL of the following: Nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen) Colchicine Corticosteroids (e.g., prednisone 	
Exclusion Criteria – Any of the following prevents coverage		
	5.1 Inadequate response to initial or previous canakinumab therapy	
	5.2 Health Alliance does not cover concurrent therapy with other immunomodulators based upon the possible increased risk for infections and other potential pharmacological interactions.	

3.4 For patients with moderate disease that has primarily systemic symptoms with no joint