

<b>POLICY NAME</b>	Pretomanid	<b>POLICY #</b>	2729P
--------------------	------------	-----------------	-------

## Criteria

### Coverage Criteria

- ☐ Treatment of pulmonary tuberculosis (TB) that is resistant to isoniazid, rifamycins, a fluoroquinolone and an alternative injectable antibiotic OR pulmonary TB resistant to isoniazid and rifampin
- ☐ Member is treatment intolerant or non-responsive to standard therapy. [Treatment failure refers to failure of cultures to become negative during the course of treatment, or reappearance of positive cultures after the cultures convert to negative during treatment]
- ☐ Age 18 years or older
- ☐ Prescribed by or in consultation with an Infectious Disease specialist or pulmonologist (lung doctor)
- ☐ Documentation that pretomanid will be used in combination with Sirturo (bedaquiline) and linezolid

### Exclusion Criteria – Any of the following prevents coverage

- ☐ Not to be used in those who have a contraindication to bedaquiline and/or linezolid
- ☐ Drug-sensitive tuberculosis
- ☐ Latent infection due to *Mycobacterium tuberculosis*
- ☐ Extra-pulmonary infection (infection outside of the lungs) due to *Mycobacterium tuberculosis*
- ☐ Multi-drug resistant tuberculosis in patients who are responsive to standard therapy and are not treatment intolerant