

Entropie

Relative Entropie von endlichen Quantenzuständen

Moritz Lanz

24. Juni 2026

Abstract

This is the abstract.

Contents

1	Section 1	1
1.1	Some Definitions	1
1.2	First Inequality	2
2	Section 2	4
2.1	Subsection A	4
	References	5

1 Section 1

1.1 Some Definitions

Question 1.1. Definieren hier die f-Divergenz allgemein für positive lineare Funktionale über von Neumann Algebren. Haben wir dadurch später Vorteile, oder willst du das einfach erstmal so allgemein wie möglich machen? Für die Physik und viele Resultate werden wir schätze ich mal Zustände verwenden.

Definition 1.2. Let $\varphi, \psi : \mathcal{M} \rightarrow \mathbb{C}$ positive linear functionals. We call φ absolutely continuous with respect to ψ and write $\varphi \ll \psi$ if:

$$\psi(A^*A) = 0 \implies \varphi(A^*A) = 0 \quad \forall A \in \mathcal{M}. \quad (1)$$

We call φ and ψ mutually absolutely continuous if:

$$\psi(A^*A) = 0 \iff \varphi(A^*A) = 0. \quad (2)$$

For normal linear functionals we get additional information from mutual continuity. Therefore we first revise some properties.

Theorem 1.3. Let $\mathcal{M}$ be a von Neumann algebra and $\omega \in S(\mathcal{M})$. Then, ω is a normal state if and only if there exists a density operator $\rho \in \mathcal{M}_+$ such that $\omega(A) = \text{Tr}(\rho A)$ for all $A \in \mathcal{M}$.

Definition 1.4. Let $\varphi, \psi : \mathcal{M} \rightarrow \mathbb{C}$ be positive linear functionals over a von Neumann algebra $\mathcal{M}$ such that $\varphi \ll \psi$ *was das nochmal genau???* and $f : (0, \infty) \rightarrow \mathbb{R}$ be a convex function with $f(1) = 0$ (*brauche ich das hier???*). The *f-divergence* of φ and ψ is given by

$$\begin{aligned} D_f(\varphi|\psi) &= \int_1^\infty f''(\gamma) E_\gamma(\varphi|\psi) + \gamma^{-3} f''(\gamma^{-1}) E_\gamma(\psi|\varphi) d\gamma \\ &= \int_1^\infty f''(\gamma) E_\gamma(\varphi|\psi) d\gamma + \int_0^1 \gamma f''(\gamma) E_{\frac{1}{\gamma}}(\psi|\varphi) d\gamma. \end{aligned} \quad (3)$$

Definition 1.5. Let φ and ψ be as above. (sicher? oder doch Zustände?) The hockeystick divergence $E_\gamma(\varphi|\psi)$ is defined as follows:

$$E_\gamma(\varphi|\psi) := (\psi - \gamma\varphi)_+(\mathbb{1}) \quad (4)$$

Some facts about the hockeystick divergence.

Proposition 1.6. Let φ_1, φ_2 be positive linear functionals over a von Neumann algebra, and p_1, p_2 are (pos. lin. functionals or states?)

$$\lim_{t \rightarrow \infty} E_t(\varphi_1|\varphi_2) = \varphi_1(p_2^\perp). \quad (5)$$

$$\lim_{t \rightarrow \infty} E_t(\varphi_1, \varphi_2) = 0 \iff p_1 \leq p_2. \quad (6)$$

1.2 First Inequalities

In [SV16] there are a lot of inequalities for f-divergences in the context of probability measures.

We want to generalize them for states over arbitrary von Neumann algebras (even vNA of type III).

We start with inequalities (84) from [SV16].

Question 1.7. Wann ist $D_f(\varphi|\psi)$ endlich? Evtl wenn ψ normal ist?

Proposition 1.8. Let $f, g : (0, \infty) \rightarrow \mathbb{R}$ convex functions with $f(1) = g(1)$, $f'(1) = g'(1)$ and $f(t) \leq g(t)$ for all $t \in (0, \infty)$. Also let φ, ψ be positive linear functionals over a von Neumann algebra such that $\varphi \ll \psi$ ($D_f(\varphi|\psi), D_g(\varphi|\psi)$ are finite)

$$D_f(\varphi|\psi) \leq D_g(\varphi|\psi). \quad (7)$$

Remark. Alexandrov's theorem states that every convex function defined on an open domain is twice differentiable almost everywhere.

Remark. By assumption $f'(1) = g'(1)$ indirectly require, that f and g are differentiable at $x = 1$. Therefore, we cannot put in e.g. the *total variational distance* (see [SV16] eq. (55)).

Proof. We begin by reformulating the f-divergence given in definition 1.4.

$$\begin{aligned}
D_f(\varphi|\psi) &= \int_1^\infty f''(\gamma) E_\gamma(\varphi|\psi) + \gamma^{-3} f''(\gamma^{-1}) E_\gamma(\psi|\varphi) d\gamma \\
&= \int_1^\infty f''(\gamma) E_\gamma(\varphi|\psi) d\gamma + \int_0^1 \gamma f''(\gamma) E_{\frac{1}{\gamma}}(\psi|\varphi) d\gamma \\
&\stackrel{(a)}{=} \int_1^\infty f''(\gamma) E_\gamma(\varphi|\psi) d\gamma + \int_0^1 \gamma f''(\gamma) \left(\frac{1}{\gamma} E_\gamma(\varphi|\psi) + \psi(\mathbb{1}) - \frac{1}{\gamma} \varphi(\mathbb{1}) \right) d\gamma \\
&= \int_1^\infty f''(\gamma) E_\gamma(\varphi|\psi) d\gamma + \int_0^1 f''(\gamma) E_\gamma(\varphi|\psi) d\gamma + \int_0^1 f''(\gamma) (\gamma \psi(\mathbb{1}) - \varphi(\mathbb{1})) d\gamma \\
&= \int_0^\infty f''(\gamma) E_\gamma(\varphi|\psi) d\gamma + \psi(\mathbb{1}) \int_0^1 \gamma f''(\gamma) d\gamma - \varphi(\mathbb{1}) \int_0^1 f''(\gamma) d\gamma \\
&\stackrel{(b)}{=} [f'(\gamma) E_\gamma(\varphi|\psi)]_0^\infty - \int_0^\infty f'(\gamma) E'_\gamma(\varphi|\psi) d\gamma + \psi(\mathbb{1}) \left([\gamma f'(\gamma)]_0^1 - \int_0^1 f'(\gamma) d\gamma \right) - \varphi(\mathbb{1}) [f'(\gamma)]_0^1 \\
&\stackrel{(c)}{=} [f'(\gamma) E_\gamma(\varphi|\psi)]_0^\infty - \left([f(\gamma) E'_\gamma(\varphi|\psi)]_0^\infty - \int_0^\infty f(\gamma) E''_\gamma(\varphi|\psi) d\gamma \right) \\
&\quad + \psi(\mathbb{1}) \left([\gamma f'(\gamma)]_0^1 - [f(\gamma)]_0^1 \right) - \varphi(\mathbb{1}) [f'(\gamma)]_0^1 \\
&= \lim_{\gamma \rightarrow \infty} f'(\gamma) E_\gamma(\varphi|\psi) - \lim_{\gamma \rightarrow 0} f'(\gamma) E_\gamma(\varphi|\psi) - \left(\lim_{\gamma \rightarrow \infty} f(\gamma) E'_\gamma(\varphi|\psi) - \lim_{\gamma \rightarrow 0} f(\gamma) E'_\gamma(\varphi|\psi) \right) \\
&\quad + \int_0^\infty f(\gamma) E''_\gamma(\varphi|\psi) d\gamma + \psi(\mathbb{1}) \left(f'(1) - \lim_{\gamma \rightarrow 0} \gamma f'(\gamma) - f(1) + \lim_{\gamma \rightarrow 0} f(\gamma) \right) \\
&\quad - \varphi(\mathbb{1}) \left(f'(1) - \lim_{\gamma \rightarrow 0} f'(\gamma) \right) \\
&= \lim_{\gamma \rightarrow \infty} (f'(\gamma) E_\gamma(\varphi|\psi) - f(\gamma) E'_\gamma(\varphi|\psi)) + \int_0^\infty f(\gamma) E''_\gamma(\varphi|\psi) d\gamma \\
&\quad + \lim_{\gamma \rightarrow 0} f(\gamma) (E'_\gamma(\varphi|\psi) + \psi(\mathbb{1})) + \lim_{\gamma \rightarrow 0} f'(\gamma) (\varphi(\mathbb{1}) - \gamma \psi(\mathbb{1}) - E_\gamma(\varphi|\psi)) \\
&\quad + \underbrace{\psi(\mathbb{1}) f'(1) - \psi(\mathbb{1}) f(1) - \varphi(\mathbb{1}) f'(1)}_{=:K}
\end{aligned} \tag{8}$$

(a) state exchange, (b), (c) partial integration K is a constant here, fällt weg bei $D_g - D_f$, da $f(1) = g(1)$ und $f'(1) = g'(1)$.

(a)

Next we want to show under which conditions $D_g(\varphi|\psi) \leq D_f(\varphi|\psi)$ holds.

$$\begin{aligned}
D_g(\varphi|\psi) - D_f(\varphi|\psi) &= \int_0^\infty \underbrace{(g(\gamma) - f(\gamma))}_{\geq 0} \underbrace{E''_\gamma(\varphi|\psi)}_{\geq 0} d\gamma + \lim_{\gamma \rightarrow 0} \underbrace{(g(\gamma) - f(\gamma))}_{\geq 0} \underbrace{(E'_\gamma(\varphi|\psi) + \psi(\mathbb{1}))}_{\geq 0 \text{ (a)}} \\
&\quad + \lim_{\gamma \rightarrow \infty} ((g'(\gamma) - f'(\gamma)) E_\gamma(\varphi|\psi) - (g(\gamma) - f(\gamma)) E'_\gamma(\varphi|\psi)) \\
&\quad + \lim_{\gamma \rightarrow 0} (g'(\gamma) - f'(\gamma)) \underbrace{(\varphi(\mathbb{1}) - \gamma \psi(\mathbb{1}) - E_\gamma(\varphi|\psi))}_{\leq 0 \text{ (b)}}
\end{aligned} \tag{9}$$

(a) follow from lipschitz-continuity of E_γ ; (b) follows from strict positivity of E_γ .

□

2 Section 2

2.1 Subsection A

asdasdasdassdasas

References

- [SV16] Igal Sason and Sergio Verdu. “ f -Divergence Inequalities”. In: *IEEE Transactions on Information Theory* 62.11 (Nov. 2016), pp. 5973–6006. ISSN: 1557-9654. DOI: 10.1109/tit.2016.2603151. URL: <http://dx.doi.org/10.1109/TIT.2016.2603151>.