

Assignment 3 (DB) - Answers

Srijan Kundu

September 26, 2022

Question 1

Let $A = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \end{pmatrix}$ and $B = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$. Find AB and BA .

Answer

Matrix A being an $(1 \times n)$ matrix and matrix B being a $(n \times 1)$ matrix, both the matrix operations AB and BA are valid.

$$AB = \sum_{i=1}^n a_i b_i$$

$$BA = \begin{pmatrix} b_1 a_1 & b_1 a_2 & \cdots & b_1 a_n \\ b_2 a_1 & b_2 a_2 & \cdots & b_2 a_n \\ \vdots & \vdots & \ddots & \vdots \\ b_n a_1 & b_n a_2 & \cdots & b_n a_n \end{pmatrix}$$

Here, AB is a scalar quantity, whereas BA is a $(n \times n)$ matrix.

Question 2

Find examples for the following statements:

1. Suppose that the matrix product AB is defined. Then the product BA need not be defined.
2. Suppose that the matrices A and B are square matrices of order n . Then AB and BA may or may not be equal.

Answer

1. Say $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{pmatrix}$ and $B = \begin{pmatrix} 7 & 8 \\ 9 & 10 \end{pmatrix}$. Then AB is defined as number of columns of A = number of rows of B , and will be of dimension (3×2) . But, BA is not defined, since number of columns of $B \neq$ number of rows of A .

2. Consider $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$. Then $AB = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, whereas $BA = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$.

Question 3

Show that for any square matrix A , $S = \frac{1}{2}(A + A^T)$ is symmetric, $T = \frac{1}{2}(A - A^T)$ is skew-symmetric, and $A = S + T$.

Answer

Say A is a $(n \times n)$ matrix whose elements are $a_{(i,j)}$; $i, j \in \{1, 2, \dots, n\}$.

The $(i, j)^{th}$ element of $S = \frac{1}{2}(A + A^T)$ would be $s_{(i,j)} = \frac{1}{2}(a_{(i,j)} + a_{(j,i)}) = \frac{1}{2}(a_{(j,i)} + a_{(i,j)}) = s_{(j,i)}$; the $(j, i)^{th}$ element of S . Hence, S is symmetric by definition.

The $(i, j)^{th}$ element of $T = \frac{1}{2}(A - A^T)$ would be $t_{(i,j)} = \frac{1}{2}(a_{(i,j)} - a_{(j,i)}) = -\frac{1}{2}(a_{(j,i)} - a_{(i,j)}) = -t_{(j,i)}$; the negative of the $(j, i)^{th}$ element of T . Hence, S is skew-symmetric by definition.

For the last part, $S + T = \{s_{(i,j)} + t_{(i,j)}\} = \{\frac{1}{2}(a_{(i,j)} + a_{(j,i)}) + \frac{1}{2}(a_{(i,j)} - a_{(j,i)})\} = a_{(i,j)} = A$.

Question 4

Show that the product of two lower triangular matrices is a lower triangular matrix. A similar statement holds for upper triangular matrices.

Answer

Lets A and B be two $(n \times n)$ lower triangular matrices. We fix $i < j$. Then

$$[AB]_{(i,j)} = \sum_{k=1}^n A_{(i,k)} \cdot B_{(k,j)}$$

Now, either $i < k$ or $k < j$ holds, because the negation $k \leq i < j \leq k$ is not possible.

So either $A_{(i,k)} = 0$ or $B_{(k,j)} = 0$, since A and B both are lower triangular matrices.

Therefore all the addends are zero; i.e. $[AB]_{(i,j)} = 0 \forall i, j \in \{1, 2, \dots, n\}$ and $i < j$.

Hence, that the product of two lower triangular matrices is a lower triangular matrix.

Question 5

Show that the diagonal entries of a skew-symmetric matrix are zero.

Answer

A $(n \times n)$ matrix A is defined to be a skew-symmetric matrix if $\forall i, j \in \{1, 2, \dots, n\}$, $A[i, j] = -A[j, i]$. For diagonal elements of A , $i = j$; which gives $A[i, i] = 0 \forall i \in \{1, 2, \dots, n\}$.

Hence, the diagonal entries of a skew-symmetric matrix are zero.