
Math for EE



G. V. V. Sharma

About this Book

This book introduces matrices through high school coordinate geometry. This approach makes it easier for beginners to learn Python for scientific computing. All problems in the book are from NCERT mathematics textbooks from Class 9-12. Exercises are from CBSE board exam papers.

The content is sufficient for industry jobs and covers nearly all matrix prerequisites for machine learning. There is no copyright, so readers are free to print and share.

License: https://creativecommons.org/licenses/by-sa/3.0/
and
https://www.gnu.org/licenses/fdl-1.3.en.html

First manual appeared in January 2018
First edition published on July 10, 2024

In this edition, some incorrect solutions were removed. All figures redrawn. More problems added.

CONTENTS

1 CBSE Class 10 Mathematics 2025 (Set 2) 4
1.1 Formulae 4
1.2 Problems 7

i

1. CBSE Class 10 Mathematics 2025 (Set 2)

A. Formulae

1.1.1 Tangent–radius property. If PA is a tangent to a circle with centre O at the point A,

$$OA \perp PA \tag{1.1.1.1}$$

1.1.2 Tangents from an external point. If PA and PB are tangents from P to a circle with centre O,

$$PA = PB, \quad \triangle OAP \cong \triangle OBP, \quad \angle POA = \angle POB \tag{1.1.2.1}$$

Solution: $OA = OB$ (radii), $PA = PB$ and OP is common, so the triangles are congruent by SSS.

1.1.3 Angle sum of a quadrilateral and of a triangle,

$$\sum \text{angles} = 360^\circ \text{ (quadrilateral)}, \quad 180^\circ \text{ (triangle)} \tag{1.1.3.1}$$

1.1.4 Angle in a semicircle. If AB is a diameter and C lies on the circle,

$$\angle ACB = 90^\circ \tag{1.1.4.1}$$

1.1.5 Tangent–chord theorem. The angle between a tangent and a chord through the point of contact equals the angle in the alternate segment. If PC is tangent at C,

$$\angle PCB = \angle CAB \tag{1.1.5.1}$$

1.1.6 Sector of a circle. For radius r, central angle θ (in radians) and arc length L,

$$L = r\theta, \quad A = \frac{1}{2}r^2\theta = \frac{1}{2}rL \tag{1.1.6.1}$$

Solution: The area is the polar integral

$$A = \int_0^\theta \int_0^r \rho \, d\rho \, d\phi = \frac{r^2\theta}{2} \tag{1.1.6.2}$$

and $L = r\theta$, so $A = \frac{1}{2}rL$.

1.1.7 AA similarity. If two angles of one triangle equal two angles of another, the triangles are similar and corresponding sides are in the same ratio. For $\triangle LMN \sim \triangle ABC$,

$$\frac{LM}{AB} = \frac{MN}{BC} = \frac{LN}{AC} \tag{1.1.7.1}$$

1.1.8 Cube of side a,

$$V = a^3, \quad S = 6a^2 \implies \frac{S_1}{S_2} = \left(\frac{V_1}{V_2}\right)^{2/3} \tag{1.1.8.1}$$

Solution: $\frac{a}{b} = \left(\frac{V_1}{V_2}\right)^{1/3}$ and $\frac{S_1}{S_2} = \left(\frac{a}{b}\right)^2$. Used in Q.3.

1.1.9 Cone and sphere. For a cone of radius R and height h, and a sphere of radius r,

$$V_{\text{cone}} = \frac{1}{3}\pi R^2 h, \quad V_{\text{sphere}} = \frac{4}{3}\pi r^3 \tag{1.1.9.1}$$

Volume of solids dropped into a full vessel equals the volume of water displaced. Used in Q.33.

1.1.10 Section formula. The point $\mathbf{R}$ dividing $\mathbf{P}$ and $\mathbf{Q}$ internally in the ratio $m:n$,

$$\mathbf{R} = \frac{m\mathbf{Q} + n\mathbf{P}}{m+n} = \mathbf{P} + \frac{m}{m+n}(\mathbf{Q} - \mathbf{P}) \tag{1.1.10.1}$$

For $m = n$ this is the mid-point $\frac{\mathbf{P}+\mathbf{Q}}{2}$. Used in Q.4, Q.15, Q.31(b).

1.1.11 Squared length and perpendicularity,

$$\|\mathbf{A}\|^2 = \mathbf{A}^\top \mathbf{A}, \quad \mathbf{A} \perp \mathbf{B} \iff \mathbf{A}^\top \mathbf{B} = 0 \tag{1.1.11.1}$$

Used in Q.8, Q.11.

1.1.12 Altitude to the hypotenuse. If $\angle BAC = 90^\circ$ and $AD \perp BC$, then

$$AD^2 = BD \cdot DC \tag{1.1.12.1}$$

Solution: Take $\mathbf{B} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\mathbf{D} = \begin{pmatrix} BD \\ 0 \end{pmatrix}$, $\mathbf{C} = \begin{pmatrix} BD + DC \\ 0 \end{pmatrix}$, $\mathbf{A} = \begin{pmatrix} BD \\ h \end{pmatrix}$. From (1.1.11.1),

$$(\mathbf{B} - \mathbf{A})^\top (\mathbf{C} - \mathbf{A}) = -BD \cdot DC + h^2 = 0 \tag{1.1.12.2}$$

Used in Q.11.

1.1.13 Euclidean algorithm and LCM,

$$\gcd(a, b) = \gcd(b, a \bmod b), \quad \operatorname{lcm}(a, b) = \frac{ab}{\gcd(a, b)} \tag{1.1.13.1}$$

The LCM of three numbers is $\operatorname{lcm}(\operatorname{lcm}(a, b), c)$. Used in Q.9, Q.23.

1.1.14 Numbers in a given ratio. If two numbers are in the ratio $p:q$ with $\gcd(p, q) = 1$ and HCF k,

$$\text{numbers} = pk,\ qk, \quad \operatorname{lcm} = pqk \tag{1.1.14.1}$$

Used in Q.23(b).

1.1.15 Zeroes of a quadratic. For $ax^2 + bx + c$ with zeroes α, β,

$$\alpha + \beta = -\frac{b}{a}, \quad \alpha\beta = \frac{c}{a} \tag{1.1.15.1}$$

Used in Q.6, Q.21(b).

1.1.16 Newton–Raphson method. To solve $f(x) = 0$,

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \tag{1.1.16.1}$$

Solution: The tangent to f at x_n is $y = f(x_n) + f'(x_n)(x - x_n)$. Setting $y = 0$ gives x_{n+1}. Used in Q.6, Q.21(b).

1.1.17 Arithmetic progression with first term a and common difference d,

$$a_n = a + (n-1)\,d, \quad S_n = \frac{n}{2}\left[2a + (n-1)\,d\right] = \frac{n}{2}\left(a + a_n\right) \tag{1.1.17.1}$$

Used in Q.32(a).

1.1.18 Two linear equations (Cramer's rule). For $\begin{pmatrix} a & b \\ c & d \end{pmatrix}\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} e \\ f \end{pmatrix}$ with $\Delta = ad - bc \neq 0$,

$$x = \frac{ed - bf}{\Delta}, \quad y = \frac{af - ec}{\Delta} \tag{1.1.18.1}$$

Used in Q.34.

1.1.19 Simple interest on a principal P at rate $r\%$ per annum for t years,

$$I = \frac{Prt}{100} \tag{1.1.19.1}$$

Used in Q.34.

1.1.20 Standard values and inverse sine,

$$\sin 60^\circ = \frac{\sqrt{3}}{2}, \quad \tan 30^\circ = \frac{1}{\sqrt{3}}, \quad \tan 45^\circ = 1 \tag{1.1.20.1}$$

The principal value $\sin^{-1}\left(\frac{\sqrt{3}}{2}\right) = 60^\circ$. Used in Q.5, Q.38.

1.1.21 Angle of depression. From a point at height h, a point on the ground at horizontal distance d is seen at an angle of depression α with

$$\tan\alpha = \frac{h}{d} \tag{1.1.21.1}$$

since the angle of depression equals the angle of elevation from the ground. Used in Q.38.

1.1.22 Probability with an indicator map. For a finite sample space S with outcome probabilities p_s and an event E with indicator $m(s) \in \{0, 1\}$,

$$\Pr(E) = \sum_{s \in S} m(s)\, p_s = \mathbf{m}^\top \mathbf{p} \tag{1.1.22.1}$$

For equally likely outcomes, $p_s = \frac{1}{|S|}$. Also $\Pr(E) + \Pr(E') = 1$. Used in Q.10, Q.27.

1.1.23 Grouped data. For a class with lower limit l, width h, frequency f, preceding cumulative frequency cf, and class mark x_i,

$$\bar{x} = \frac{\sum f_i x_i}{\sum f_i} \tag{1.1.23.1}$$

$$\text{Median} = l + \frac{\frac{N}{2} - cf}{f} \times h, \quad N = \sum f_i \tag{1.1.23.2}$$

The modal class is the class with the highest frequency.

B. Problems

1.2.1 If tangents PA and PB drawn from an external point P to a circle with centre O are inclined to each other at an angle of 80°, then $\angle POA$ is equal to (CBSE 2025 Set 2, Q.1)

a) 40° b) 50° c) 60° d) 80°




Fig. 1.2.1.1: Tangents PA and PB

Solution: A radius is perpendicular to the tangent at the point of contact. Hence,

$$\angle OAP = \angle OBP = 90^\circ \tag{1.2.1.1}$$

In the quadrilateral $AOBP$,

$$\angle AOB + \angle OAP + \angle APB + \angle OBP = 360^\circ \tag{1.2.1.2}$$

$$\implies \angle AOB = 360^\circ - 90^\circ - 80^\circ - 90^\circ = 100^\circ \tag{1.2.1.3}$$

Since $OA = OB$ (radii), $PA = PB$ (tangents from an external point) and OP is common, $\triangle OAP \cong \triangle OBP$, so OP bisects $\angle AOB$. Therefore,

$$\angle POA = \frac{\angle AOB}{2} = 50^\circ \quad \text{(Option B)} \tag{1.2.1.4}$$

The following code verifies the solution

codes/1.c

1.2.2 The volumes of two cubes are in the ratio $8 : 125$. The ratio of their surface areas is (CBSE 2025 Set 2, Q.3)

a) $8 : 125$ b) $4 : 25$ c) $2 : 5$ d) $16 : 25$

Solution: Let the sides of the cubes be a and b. Then

$$\frac{a^3}{b^3} = \frac{8}{125} \implies \frac{a}{b} = \sqrt[3]{\frac{8}{125}} = \frac{2}{5} \tag{1.2.2.1}$$

The surface area of a cube of side a is $6a^2$. Hence,

$$\frac{S_1}{S_2} = \frac{6a^2}{6b^2} = \left(\frac{a}{b}\right)^2 = \frac{4}{25} \implies S_1 : S_2 = 4 : 25 \quad \text{(Option B)} \tag{1.2.2.2}$$

The following code verifies the solution

codes/3.c

1.2.3 Two vertices of $\triangle PQR$ are $P(-1, 5)$ and $Q(5, 2)$. The coordinates of the point R on PQ which divides PQ in the ratio $2 : 1$ are (CBSE 2025 Set 2, Q.4)

a) $(3, -3)$ b) $(5, 5)$ c) $(3, 3)$ d) $(5, 1)$

Solution: Let $\mathbf{P} = \begin{pmatrix} -1 \\ 5 \end{pmatrix}$ and $\mathbf{Q} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$. The displacement from P to Q is

$$\mathbf{Q} - \mathbf{P} = \begin{pmatrix} 6 \\ -3 \end{pmatrix} \tag{1.2.3.1}$$

Since $PR : RQ = 2 : 1$, the point R lies two-thirds of the way from P to Q:

$$\mathbf{R} = \mathbf{P} + \frac{2}{3}(\mathbf{Q} - \mathbf{P}) = \begin{pmatrix} -1 \\ 5 \end{pmatrix} + \begin{pmatrix} 4 \\ -2 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \end{pmatrix} \quad \text{(Option C)} \tag{1.2.3.2}$$

The following code verifies the solution

codes/4.c

The following code generates Fig. 1.2.3.1.

codes/4plot.py




Fig. 1.2.3.1: Point R dividing PQ in the ratio $2 : 1$

1.2.4 If $\sin 4\theta = \frac{\sqrt{3}}{2}$, then $\frac{\theta}{3}$ is equal to (CBSE 2025 Set 2, Q.5)

a) 60° b) 20° c) 15° d) 5°

Solution: Taking the principal (acute) value,

$$4\theta = \sin^{-1}\left(\frac{\sqrt{3}}{2}\right) = 60^\circ \implies \theta = 15^\circ \implies \frac{\theta}{3} = 5^\circ \quad \text{(Option D)} \tag{1.2.4.1}$$

The following code verifies the solution

codes/5.c

1.2.5 The zeroes of the polynomial $p(y) = 7y^2 - \frac{11}{3}y - \frac{2}{3}$ are (CBSE 2025 Set 2, Q.6)

a) $-\frac{2}{3}, -\frac{1}{7}$ b) $-\frac{2}{7}, -\frac{1}{3}$ c) $\frac{2}{3}, \frac{1}{7}$ d) $\frac{2}{3}, -\frac{1}{7}$

Solution: Multiplying by 3, the zeroes of p are the zeroes of

$$g(y) = 21y^2 - 11y - 2, \quad g'(y) = 42y - 11 \tag{1.2.5.1}$$

The Newton–Raphson iteration is

$$y_{n+1} = y_n - \frac{g(y_n)}{g'(y_n)} = y_n - \frac{21y_n^2 - 11y_n - 2}{42y_n - 11} \tag{1.2.5.2}$$

Starting from $y_0 = 0.7$ and $y_0 = -0.2$, the iterations converge to

$$y = \frac{2}{3}, \quad y = -\frac{1}{7} \quad \text{(Option D)} \tag{1.2.5.3}$$

which agree with the factorisation $g(y) = (3y - 2)(7y + 1)$. The following code verifies the solution

codes/6.c

The following code generates Fig. 1.2.5.1.

codes/6plot.py




Fig. 1.2.5.1: Graph of $p\left(y\right)$ and its zeroes

1.2.6 $AOBC$ is a rectangle whose three vertices are $A\left(0,2\right)$, $O\left(0,0\right)$ and $B\left(4,0\right)$. The square of the length of its diagonal is (CBSE 2025 Set 2, Q.8)

a) 36 b) 20 c) 16 d) 4

Solution: The fourth vertex is $C\left(4,2\right)$ and the diagonal is AB. With $\mathbf{A} = \begin{pmatrix}0\\2\end{pmatrix}$, $\mathbf{B} = \begin{pmatrix}4\\0\end{pmatrix}$,

$$\mathbf{B}-\mathbf{A} = \begin{pmatrix}4\\-2\end{pmatrix}, \qquad \left\|\mathbf{B}-\mathbf{A}\right\|^2 = \left(\mathbf{B}-\mathbf{A}\right)^\top\left(\mathbf{B}-\mathbf{A}\right) = 16+4 = 20 \quad \text{(Option B)} \tag{1.2.6.1}$$

The following code verifies the solution

codes/8.c

The following code generates Fig. 1.2.6.1.

codes/8plot.py




Fig. 1.2.6.1: Rectangle $AOBC$ and its diagonals

1.2.7 If x is the LCM of $4, 6, 8$, y is the LCM of $3, 5, 7$ and p is the LCM of x and y, then (CBSE 2025 Set 2, Q.9)

a) $p = 35x$ b) $p = 4y$ c) $p = 8x$ d) $p = 16y$

Solution: The Euclidean algorithm gives $\gcd(a, b) = \gcd(b, a \bmod b)$, and

$$\operatorname{lcm}(a, b) = \frac{ab}{\gcd(a, b)} \tag{1.2.7.1}$$

Applying this repeatedly,

$$x = \operatorname{lcm}(4, 6, 8) = 24, \quad y = \operatorname{lcm}(3, 5, 7) = 105, \quad p = \operatorname{lcm}(24, 105) = 840 \tag{1.2.7.2}$$

Since $35x = 35 \times 24 = 840 = p$, the correct option is $p = 35x$ (Option A). The following code verifies the solution

codes/9.c

1.2.8 A die is thrown once. The probability of getting a number which is not a factor of 36 is (CBSE 2025 Set 2, Q.10)

a) $\frac{1}{2}$ b) $\frac{2}{3}$ c) $\frac{1}{6}$ d) $\frac{5}{6}$

Solution: The sample space is $S = \{1, 2, 3, 4, 5, 6\}$ and each outcome has probability $\frac{1}{6}$. Define the indicator map

$$m(s) = \begin{cases} 1 & s \text{ is not a factor of } 36 \\ 0 & s \text{ is a factor of } 36 \end{cases} \tag{1.2.8.1}$$

s	1	2	3	4	5	6
$m(s)$	0	0	0	0	1	0

TABLE 1.2.8: Mapping of outcomes

The required probability is


$$\Pr(E) = \sum_{s \in S} m(s) \cdot \frac{1}{6} = \frac{1}{6} \quad \text{(Option C)} \tag{1.2.8.2}$$

The following code verifies the solution

codes/10.c

1.2.9 In the figure, $\angle BAC = 90^\circ$ and $AD \perp BC$. If $BC = 16$ cm and $DC = 4$ cm, then $AC = x$ is equal to (CBSE 2025 Set 2, Q.11)

a) 4 cm b) 5 cm c) 8 cm d) 3 cm



$BC = 16$ cm

Fig. 1.2.9.1: Right triangle ABC with altitude AD

Solution: Take $\mathbf{B} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\mathbf{C} = \begin{pmatrix} 16 \\ 0 \end{pmatrix}$. Then $BD = 16 - 4 = 12$, so $\mathbf{D} = \begin{pmatrix} 12 \\ 0 \end{pmatrix}$ and, since $AD \perp BC$, $\mathbf{A} = \begin{pmatrix} 12 \\ h \end{pmatrix}$. As $\angle BAC = 90^\circ$,

$$(\mathbf{B} - \mathbf{A})^\top (\mathbf{C} - \mathbf{A}) = 0 \tag{1.2.9.1}$$

$$\begin{pmatrix} -12 & -h \end{pmatrix} \begin{pmatrix} 4 \\ -h \end{pmatrix} = -48 + h^2 = 0 \implies h^2 = 48 \tag{1.2.9.2}$$

Hence,

$$x^2 = \|\mathbf{C} - \mathbf{A}\|^2 = 4^2 + h^2 = 64 \implies x = 8 \text{ cm} \quad \text{(Option C)} \tag{1.2.9.3}$$

The following code verifies the solution

codes/11.c

1.2.10 The mid-point of the line segment joining $(a, 4)$ and $(2, 2b)$ is $(2, 6)$. Then $a + b$ is equal to (CBSE 2025 Set 2, Q.15)

a) 6 b) 7 c) 8 d) 16

Solution: The mid-point is the average of the position vectors:

$$\begin{pmatrix} 2 \\ 6 \end{pmatrix} = \frac{1}{2}\left(\begin{pmatrix} a \\ 4 \end{pmatrix} + \begin{pmatrix} 2 \\ 2b \end{pmatrix}\right) \implies \frac{a+2}{2} = 2, \quad \frac{4+2b}{2} = 6 \implies a = 2, \quad b = 4 \tag{1.2.10.1}$$

Therefore $a + b = 6$ (Option A). The following code verifies the solution

codes/15.c

The following code generates Fig. 1.2.10.1.

codes/15plot.py




Fig. 1.2.10.1: Segment PQ and its mid-point M

1.2.11 The area of a sector of a circle of radius 36 cm is 54π cm^2. The length of the corresponding arc of the sector is (CBSE 2025 Set 2, Q.16)

a) 8π cm b) 6π cm c) 4π cm d) 3π cm

Solution: For a sector of radius r and arc length L,

$$A = \frac{1}{2}rL \implies L = \frac{2A}{r} = \frac{2\,(54\pi)}{36} = 3\pi \text{ cm} \quad \text{(Option D)} \tag{1.2.11.1}$$

The central angle is $\theta = \frac{L}{r} = \frac{\pi}{12} = 15^\circ$. The following code verifies the solution

codes/16.c

The following code generates Fig. **??**.

codes/16plot.py

1.2.12 $\angle M = \angle B = 130^\circ$ and $\angle N = 28^\circ$. If $LM = 45$ cm, $MN = 63$ cm, $LN = 72$ cm and $AB = 5$ cm, then AC is equal to (CBSE 2025 Set 2, Q.17)

a) 16 cm b) 7 cm c) 8 cm d) 4 cm




Fig. 1.2.12.1: Triangles ABC and LMN (not to scale)

Solution: In $\triangle LMN$, $\angle L = 180^\circ - 130^\circ - 28^\circ = 22^\circ = \angle A$. Since $\angle A = \angle L$ and $\angle B = \angle M$, by AA similarity

$$\triangle LMN \sim \triangle ABC \quad (L \leftrightarrow A,\ M \leftrightarrow B,\ N \leftrightarrow C) \tag{1.2.12.1}$$

The scale factor is $k = \frac{AB}{LM} = \frac{5}{45} = \frac{1}{9}$, so

$$AC = k \cdot LN = \frac{72}{9} = 8 \text{ cm} \quad \text{(Option C)} \tag{1.2.12.2}$$

The following code verifies the solution

codes/17.c

1.2.13 $\triangle DEF$ and $\triangle PQR$ are such that $\angle D = \angle Q$ and $\angle E = \angle R$. Which of the following proportions is <u>not</u> true? (CBSE 2025 Set 2, Q.18)

a) $\frac{DE}{QR} = \frac{DF}{PQ}$ b) $\frac{EF}{RP} = \frac{DF}{PQ}$ c) $\frac{EF}{RP} = \frac{DE}{QR}$ d) $\frac{DE}{PQ} = \frac{EF}{RP}$

Solution: By AA similarity, $\triangle DEF \sim \triangle QRP$ with $D \leftrightarrow Q$, $E \leftrightarrow R$, $F \leftrightarrow P$. Corresponding sides are proportional:

$$\frac{DE}{QR} = \frac{EF}{RP} = \frac{DF}{QP} \tag{1.2.13.1}$$

Options A, B and C compare corresponding sides. Option D compares DE with PQ, which are not corresponding sides. Hence the proportion that is not true is Option D. The following code checks all four options on a pair of similar triangles

codes/18.c

1.2.14 If α and β are the zeroes of the polynomial $x^2 - 2x - 1$, find the value of $\frac{1}{2\alpha} + \frac{1}{2\beta} + 3\alpha\beta$. (CBSE 2025 Set 2, Q.21(b))

Solution: Let $f(x) = x^2 - 2x - 1$, so $f'(x) = 2x - 2$. The Newton–Raphson iteration is

$$x_{n+1} = x_n - \frac{x_n^2 - 2x_n - 1}{2x_n - 2} \tag{1.2.14.1}$$

Starting from $x_0 = 2.5$ and $x_0 = -0.5$ gives

$$\alpha = 1 + \sqrt{2}, \quad \beta = 1 - \sqrt{2} \tag{1.2.14.2}$$

From the sum and product of zeroes, $\alpha + \beta = 2$ and $\alpha\beta = -1$, so

$$\frac{1}{2\alpha} + \frac{1}{2\beta} + 3\alpha\beta = \frac{\alpha + \beta}{2\alpha\beta} + 3\alpha\beta = \frac{2}{-2} - 3 = -4 \tag{1.2.14.3}$$

The following code verifies the solution

codes/21b.c

1.2.15 Find the smallest number which is divisible by both 644 and 462. (CBSE 2025 Set 2, Q.23(a))

Solution: The required number is $\text{lcm}(644, 462)$. Using $644 = 2^2 \cdot 7 \cdot 23$ and $462 = 2 \cdot 3 \cdot 7 \cdot 11$,

$$\gcd(644, 462) = 14, \quad \text{lcm}(644, 462) = \frac{644 \times 462}{14} = 21252 \tag{1.2.15.1}$$

The following code verifies the solution (both parts of Q.23)

codes/23.c

1.2.16 Two numbers are in the ratio $4 : 5$ and their HCF is 11. Find their LCM. (CBSE 2025 Set 2, Q.23(b))

Solution: Let the numbers be $4k$ and $5k$. Since $\gcd(4, 5) = 1$, the HCF is k, so

$k = 11$ and the numbers are 44 and 55. Then

$$\text{lcm}\,(44, 55) = \frac{44 \times 55}{11} = 220 \tag{1.2.16.1}$$

This is verified by the code for Q.23(a) above.

1.2.17 In a school, 65% of the students have black eyes, 25% have brown eyes and the rest have blue eyes. A student is chosen at random. Find the probability that the student has (a) blue eyes, (b) brown or black eyes. (CBSE 2025 Set 2, Q.27)

Solution: The probability of blue eyes is $1-0.65-0.25 = 0.10$. Order the outcomes as (Black, Brown, Blue) with probabilities $\mathbf{p} = \begin{pmatrix}0.65 & 0.25 & 0.10\end{pmatrix}^\top$, and map each event to 0 or 1:

Event	Black	Brown	Blue
(a) Blue	0	0	1
(b) Brown or black	1	1	0

TABLE 1.2.17: Mapping of eye colours to events

For an event with mapping $\mathbf{m}$,

$$\Pr(E) = \mathbf{m}^\top \mathbf{p} \implies \text{(a) } 0.10, \quad \text{(b) } 0.65 + 0.25 = 0.90 \tag{1.2.17.1}$$

The following code verifies the solution

codes/27.c

1.2.18 In the figure, PC is a tangent to the circle at C and AB is a diameter which, when extended, meets the tangent at P. If $\angle PCA = 110°$, find $\angle CBA$ and $\angle BCO$. (CBSE 2025 Set 2, Q.29)




Fig. 1.2.18.1: Tangent PC and diameter AB

Solution: Since AB is a diameter, $\angle ACB = 90°$. As P lies beyond B,

$$\angle PCA = \angle PCB + \angle BCA \implies \angle PCB = 110° - 90° = 20° \tag{1.2.18.1}$$

By the tangent–chord theorem, $\angle CAB = \angle PCB = 20^\circ$. In $\triangle ABC$,

$$\angle CBA = 90^\circ - 20^\circ = 70^\circ \tag{1.2.18.2}$$

Since $OB = OC$, $\triangle OBC$ is isosceles and

$$\angle BCO = \angle CBO = 70^\circ \tag{1.2.18.3}$$

The following code verifies the solution

codes/29.c

1.2.19 Find the coordinates of the points which divide the line segment joining $A\,(-2, 2)$ and $B\,(2, 8)$ into four equal parts. (CBSE 2025 Set 2, Q.31(b))

Solution: With $\mathbf{A} = \begin{pmatrix} -2 \\ 2 \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} 2 \\ 8 \end{pmatrix}$, $\mathbf{B} - \mathbf{A} = \begin{pmatrix} 4 \\ 6 \end{pmatrix}$. The division points are $\mathbf{A} + \frac{k}{4}(\mathbf{B} - \mathbf{A})$ for $k = 1, 2, 3$:

$$\mathbf{P} = \begin{pmatrix} -1 \\ 3.5 \end{pmatrix}, \quad \mathbf{Q} = \begin{pmatrix} 0 \\ 5 \end{pmatrix}, \quad \mathbf{R} = \begin{pmatrix} 1 \\ 6.5 \end{pmatrix} \tag{1.2.19.1}$$

The following code verifies the solution

codes/31b.c

The following code generates Fig. 1.2.19.1.

codes/31bplot.py




Fig. 1.2.19.1: Division of AB into four equal parts

1.2.20 The nth term of an AP is 4, its common difference is 2 and the sum of its first n terms is -14. Find n, the first term, and the sum of the first 20 terms. (CBSE 2025 Set 2, Q.32(a))

Solution: Let the first term be a. Then $a_n = a + (n-1)\,d = 4$ gives $a = 6 - 2n$. Using $S_n = \frac{n}{2}\,(a + a_n)$,

$$-14 = \frac{n}{2}\,(6 - 2n + 4) = n\,(5 - n) \implies n^2 - 5n - 14 = 0 \implies (n-7)\,(n+2) = 0 \tag{1.2.20.1}$$

Since $n > 0$, $n = 7$ and $a = 6 - 14 = -8$. Then

$$S_{20} = \frac{20}{2}\,[2\,(-8) + 19\,(2)] = 220 \tag{1.2.20.2}$$

The following code verifies the solution

codes/32a.c

1.2.21 A vessel in the form of an inverted cone of height 8 cm and radius 5 cm is filled with water to the brim. Lead shots, each a sphere of radius 0.5 cm, are dropped into the vessel until one-fourth of the water flows out. Find the number of lead shots dropped. (CBSE 2025 Set 2, Q.33)

Solution: The volume of water in the cone is

$$V_c = \frac{1}{3}\pi R^2 h = \frac{1}{3}\pi\,(5)^2\,(8) = \frac{200\pi}{3}\ \text{cm}^3 \tag{1.2.21.1}$$

The water that flows out equals the total volume of the shots, so the volume displaced is $\frac{V_c}{4} = \frac{50\pi}{3}$ cm^3. Each shot has volume

$$V_s = \frac{4}{3}\pi r^3 = \frac{4}{3}\pi\,(0.5)^3 = \frac{\pi}{6}\ \text{cm}^3 \tag{1.2.21.2}$$

Therefore the number of shots is

$$N = \frac{50\pi/3}{\pi/6} = 100 \tag{1.2.21.3}$$

The following code verifies the solution

codes/33.c

1.2.22 A man lent part of his money at 10% per annum and the rest at 15% per annum. His yearly interest was ₹1900. Had he interchanged the two rates, he would have earned ₹200 more. Find the two amounts. (CBSE 2025 Set 2, Q.34)

Solution: Let x and y be the amounts lent at 10% and 15%. Then

$$0.10x + 0.15y = 1900 \tag{1.2.22.1}$$

$$0.15x + 0.10y = 2100 \tag{1.2.22.2}$$

In matrix form,

$$\begin{pmatrix} 0.10 & 0.15 \\ 0.15 & 0.10 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1900 \\ 2100 \end{pmatrix} \tag{1.2.22.3}$$

The determinant is $0.10^2 - 0.15^2 = -0.0125$, and by Cramer's rule

$$x = \frac{1900\,(0.10) - 0.15\,(2100)}{-0.0125} = 10000, \quad y = \frac{0.10\,(2100) - 0.15\,(1900)}{-0.0125} = 6000 \tag{1.2.22.4}$$

Hence the amounts are ₹10000 at 10% and ₹6000 at 15%. The following code verifies the solution

codes/34.c

1.2.23 **Case study (rainfall).** The distribution of the number of meteorological subdivisions by annual rainfall is given below. (CBSE 2025 Set 2, Q.36)

Rainfall (mm)	200–400	400–600	600–800	800–1000	1000–1200	1200–1400
Subdivisions	3	4	7	4	3	3

TABLE 1.2.23: Rainfall distribution, total 24 subdivisions

(i) Find the modal class. (ii) Find the median rainfall **or** the mean rainfall. (iii) Find the number of subdivisions with rainfall of at least 800 mm.

Solution: (i) The highest frequency is 7, so the modal class is 600–800 mm.

(ii) Here $N = 24$ and $\frac{N}{2} = 12$. The cumulative frequencies are $3, 7, 14, 18, 21, 24$, so the median class is 600–800 with $l = 600$, $cf = 7$, $f = 7$, $h = 200$:

$$\text{Median} = l + \frac{\frac{N}{2} - cf}{f} \times h = 600 + \frac{12 - 7}{7} \times 200 \approx 742.86 \text{ mm} \tag{1.2.23.1}$$

For the mean, with class marks $x_i = 300, 500, \ldots, 1300$,

$$\bar{x} = \frac{\sum f_i x_i}{\sum f_i} = \frac{18600}{24} = 775 \text{ mm} \tag{1.2.23.2}$$

(iii) The classes from 800 mm upward have $4 + 3 + 3 = 10$ subdivisions.

The following code verifies the solution

codes/36.c

1.2.24 **Case study (lighthouse).** A lighthouse AB is 50 m high. From its top B, the angles of depression of a ship at P and, after some time, at Q are 30° and 45° respectively, with P, Q, A on the same line. (CBSE 2025 Set 2, Q.38)




Fig. 1.2.24.1: Lighthouse AB and ship positions P, Q

Solution: **(i)** In $\triangle QAB$, $\tan 45^\circ = \frac{AB}{QA}$, so

$$QA = \frac{50}{\tan 45^\circ} = 50 \text{ m} \tag{1.2.24.1}$$

(iii)(a) In $\triangle PAB$, $\tan 30^\circ = \frac{AB}{PA}$, so $PA = 50\sqrt{3}$ m, and the distance moved from P to Q is

$$PQ = PA - QA = 50\left(\sqrt{3} - 1\right) \approx 36.60 \text{ m} \tag{1.2.24.2}$$

(iii)(b) The ship covers $QA = 50$ m $= 0.05$ km in 10 minutes $= \frac{1}{6}$ h, so

$$\text{Speed} = \frac{0.05}{1/6} = 0.3 \text{ km/h} \tag{1.2.24.3}$$

The following code verifies the solution

codes/38.c