

14. (2.75 points)

Given the following output from a generalized linear model fitted to a triangle of loss development data:

<u>Standardized Pearson Residuals</u>						
Accident	12	24	36	48	60	72
<u>Year</u>	<u>Months</u>	<u>Months</u>	<u>Months</u>	<u>Months</u>	<u>Months</u>	<u>Months</u>
2010	1.68	2.65	-4.70	-2.78	3.37	0.00
2011	7.74	-1.34	-7.33	-1.91	-3.61	
2012	5.72	1.94	-6.91	4.98		
2013	-1.22	0.12	-1.94			
2014	1.67	-1.87				
2015	0.00					

<u>Standard Deviations of Standardized Pearson Residuals</u>			
Accident	Standard	Accident	Standard
<u>Year</u>	<u>Deviation</u>	<u>Year Range</u>	<u>Deviation</u>
2010	3.571	2010 to 2011	4.463
2011	5.563	2012 to 2013	4.345
2012	5.797	2014 to 2015	2.503
2013	1.045	2010 to 2012	4.741
2014	2.503	2013 to 2015	1.537

<u>Fitted Cumulative Losses</u>		
Accident	12	24
<u>Year</u>	<u>Months</u>	<u>Months</u>
2012	3,500	7,612
2013	4,210	8,749
2014	5,400	10,654

CONTINUED ON NEXT PAGE

(14 continued)

An actuary reviewing the output notices heteroscedasticity in the residuals. The actuary decides to adjust for this by calculating variance parameters before running the sampling algorithm.

Given the following residual index values from one iteration of the sampling algorithm:

Accident <u>Year</u>	Sample Residual Index <u>(row, column)</u>
2012	5, 1
2013	1, 3
2014	4, 3

a. (2.25 points)

Calculate the actuary's sampled incremental losses for accident years 2012-2014 between 12 and 24 months for the sample under consideration.

b. (0.5 point)

Explain why heteroscedastic residuals might cause issues when using a bootstrapping technique to estimate variance of unpaid claim estimates.

CONTINUED ON NEXT PAGE

EXAM 7 SPRING 2016 SAMPLE ANSWERS AND EXAMINER'S REPORT

Sample hetero adjusted residual

$$2012 \quad 1.67 \times \frac{4.741}{1.537} = 5.15$$

$$2013 \quad -4.7 \times \frac{1.537}{4.741} = -1.524$$

$$2014 \quad -1.94 \times \frac{1.537}{1.537} = -1.94$$

	<u>Fitted Incremental Losses</u>	<u>Resampled Incremental Losses</u>
2012	4,112	$5.15 \times \sqrt{4,112} + 4,112 = 4,442$
2013	4,539	$-1.52 \times \sqrt{4,539} + 4,539 = 4,436$
2014	5,254	$-1.94 \times \sqrt{5,254} + 5,254 = 5,113$

Sample Answer 3

Accident years 2010 to 2012 have similar standard deviations, so group together. Other accident years do not have similar standard deviation. Selecting groups as follows:

AY	Std Dev	Hetero Group	Adj Factor
2010	3.571	1	$4.471/4.471 = 1.00$
2011	5.563	1	1.00
2012	5.797	1	1.00
2013	1.045	2	$4.471/1.045 = 4.537$
2014	2.503	3	$4.471/2.503 = 1.894$

AY	Adj Residual (r*)	Adj Residual (r*/h)
2012	$1.67 \times 1.894 = 3.163$	$3.163 / 1.00 = 3.163$
2013	$-4.70 \times 1.000 = -4.700$	$-4.70 / 4.537 = -1.036$
2014	$-1.94 \times 4.537 = -8.802$	$-8.802 / 1.894 = -4.648$

AY	Sampled Loss
2012	$(7,612 - 3,500) + 3.163(7,612 - 3,500)^{1/2} = 4,315$
2013	$(8,749 - 4,210) - 1.036(8,749 - 4,210)^{1/2} = 4,469$
2014	$(10,654 - 5,400) - 4.648(10,654 - 5,400)^{1/2} = 4,917$

Part b: 0.5 point

Sample Answer 1

The bootstrapping process assumes we can sample residuals from anywhere in the triangle. If the variance of residuals differs then our assumption of independent residuals is

EXAM 7 SPRING 2016 SAMPLE ANSWERS AND EXAMINER'S REPORT

not valid. Adjusting for this keeps us from having overstated or understated estimated incremental losses during each iteration (depending on how the variance in that cell relates to other cells) and keeps the bootstrap variance of loss estimate from being artificially distorted.

Sample Answer 2

Bootstrapping technique requires the residuals to be IID, then we can sample residuals from any place of the triangle. Heteroscedastic residuals violates this requirement.

EXAMINER'S REPORT

Very few candidates earned full credit for both parts of this question.

Part a

- Candidates were expected to know how to select hetero-groups based on the standard deviations provided. Solutions with different hetero-groups were given credit as long as the candidate provided justification.
- Candidates needed to recognize that sampled residuals should be adjusted for heteroscedasticity.
- Candidates had difficulty earning full credit for this part.
- Common errors include the following:
 - No justification provided for selected hetero-groups.
 - Hetero-adjustment factors were calculated using the wrong standard deviations
 - Entirely skipped hetero-adjustments.
 - Applied hetero-adjustment factors to the wrong sampled residuals.
 - Candidates failed to divide by the hetero-adjustment factors to bring the adjusted residuals back to the original distribution.
 - Some candidates did not use incremental fitted losses and applied the process to cumulative losses.

Part b

- Candidates were expected to know that residuals are independent and identically distributed in order to sample residuals from the whole triangle.
- Generally, candidates did not earn full credit for this part.
- Common errors include the following:
 - Noting that bootstrapping requires residuals that are independent and identically distributed, but did not fully explain the issue when the assumption is violated.
 - Other candidates used terms such as “the variance”, “variance of incremental losses”, or “losses are iid” without mentioning that it’s the residuals that have to be IID.
 - Many candidates mentioned that it would “increase the variance of the unpaid claim estimate” which is not always the case.
 - Full or partial credit was given to candidates who explained the variance of the unpaid claims would increase or decrease due to heteroscedastic residuals.