

# AI1110 ASSIGNMENT-6

DASARI SRINITH (CS21BTECH11015)

**Abstract**—This document contains the solution for Assignment 6 (NCERT GRADE 10 CHAPTER 15 Exercise 15.2 Example 4)

## EXAMPLE 4 :

One card is drawn from a well-shuffled deck of 52 cards. Calculate the probability that the card will

- (i) be an ace,
- (ii) not be an ace.

## SOLUTION :

- 1) A deck of 52 cards have 4 suits namely "Clubs , Spades , Diamonds and Hearts".
- 2) Clubs and Spades are of black colour ,while Hearts and Diamonds are of red colour.
- 3) Each suit have 13 cards, they are King ,Queen ,Jack ,10 ,9 ,8 ,7 ,6 ,5 ,4 ,3 ,2 and ace.

Given that , A card is drawn from a well-shuffled deck of 52 cards; well-shuffling ensures "Equally Likely" outcomes.

Let  $X$  be a random variable and  $X$  maps to the following set of real number ,  $X \in \{0, 1\}$  , as defined below in Table I

Event	Description of event
$X = 0$	The drawn card is not an ace
$X = 1$	The drawn card is an ace

TABLE I

We know that there are 4 aces in a deck of 52 cards. So ,

- (i) The Probability that the drawn card is an ace is same as ,

$$\Pr(X = 1) = \frac{\text{Number of ace cards}}{\text{Total Number of cards}} \quad (1)$$

$$\Pr(X = 1) = \frac{4}{52} = \frac{1}{13} \quad (2)$$

- (ii) The Probability that the drawn card is not an ace is same as ,

$$\Pr(X = 0) = \frac{\text{Number of non-ace cards}}{\text{Total Number of cards}} \quad (3)$$

$$\Pr(X = 0) = \frac{52 - 4}{52} = \frac{48}{52} = \frac{12}{13} \quad (4)$$

One can also say that , events "the drawn card being an ace" and "the drawn card not being an ace" are complementary events . So , we can write that,

$$\Pr(X = 0) = 1 - \Pr(X = 1) \quad (5)$$

$$= 1 - \frac{1}{13} = \frac{12}{13} \quad (6)$$

which gives the same result.