Linear Grammars

Grammars with
at most one variable at the right side
of a production

Examples:

$$S \to aSb$$
$$S \to \lambda$$

$$S \to Ab$$
$$A \to aAb$$
$$A \to \lambda$$

A Non-Linear Grammar

Grammar G :

$$S \to SS$$

$$S \to \lambda$$

$$S \to aSb$$

$$S \to bSa$$

$$L(G) = \{w: \quad n_a(w) = n_b(w)\}$$

Number of a in string w

Another Linear Grammar

Grammar G :

$$S \to A$$

$$A \to aB \mid \lambda$$

$$B \to Ab$$

$$L(G) = \{a^n b^n : n \geq 0\}$$

Right-Linear Grammars

All productions have form: $A \to xB$

or

$A \to x$

string of terminals

Example: $S \to abS$

$S \to a$

Left-Linear Grammars

All productions have form: $$A \to Bx$$

or

$$A \to x$$

string of terminals

Example:

$$S \to Aab$$

$$A \to Aab \mid B$$

$$B \to a$$

Regular Grammars

Regular Grammars

A regular grammar is any
right-linear or left-linear grammar

Examples:

G_1

$$S \to abS$$
$$S \to a$$

G_2

$$S \to Aab$$
$$A \to Aab \mid B$$
$$B \to a$$

Observation

Regular grammars generate regular languages

Examples:

G_1

$$S \rightarrow abS$$

$$S \rightarrow a$$

$$L(G_1) = (ab)^* a$$

G_2

$$S \rightarrow Aab$$

$$A \rightarrow Aab \mid B$$

$$B \rightarrow a$$

$$L(G_2) = aab(ab)^*$$

Regular Grammars Generate Regular Languages

Theorem

$$\left\{\begin{array}{l}\text{Languages} \\ \text{Generated by} \\ \text{Regular Grammars}\end{array}\right\} = \left\{\begin{array}{l}\text{Regular} \\ \text{Languages}\end{array}\right\}$$

Theorem - Part 1

$$\left\{ \begin{array}{l} \text{Languages} \\ \text{Generated by} \\ \text{Regular Grammars} \end{array} \right\} \subseteq \left\{ \begin{array}{l} \text{Regular} \\ \text{Languages} \end{array} \right\}$$

Any regular grammar generates a regular language

Theorem - Part 2

$$\left\{\begin{array}{l}\text{Languages}\\ \text{Generated by}\\ \text{Regular Grammars}\end{array}\right\} \supseteq \left\{\begin{array}{l}\text{Regular}\\ \text{Languages}\end{array}\right\}$$

Any regular language is generated by a regular grammar

Proof - Part 1

$$\left\{\begin{array}{l}\text{Languages} \\ \text{Generated by} \\ \text{Regular Grammars}\end{array}\right\} \subseteq \left\{\begin{array}{l}\text{Regular} \\ \text{Languages}\end{array}\right\}$$

The language $L(G)$ generated by any regular grammar G is regular

The case of Right-Linear Grammars

Let G be a right-linear grammar

We will prove: $L(G)$ is regular

Proof idea: We will construct NFA M with $L(M) = L(G)$

Grammar G is right-linear

Example:

$$S \to aA \mid B$$

$$A \to aa\ B$$

$$B \to b\ B \mid a$$

Construct NFA M such that every state is a grammar variable:



special final state

$$S \to aA \mid B$$

$$A \to aa\ B$$

$$B \to b\ B \mid a$$

Add edges for each production:



$$S \to aA$$



$$S \to aA \mid B$$



$$S \rightarrow aA \mid B$$

$$A \rightarrow aa\ B$$



$$S \to aA \mid B$$

$$A \to aa\ B$$

$$B \to bB$$



$$S \rightarrow aA \mid B$$

$$A \rightarrow aa\ B$$

$$B \rightarrow bB \mid a$$



$$S \Rightarrow aA \Rightarrow aaaB \Rightarrow aaabB \Rightarrow aaaba$$

NFA M



Grammar

G

$$S \rightarrow aA \mid B$$

$$A \rightarrow aa\ B$$

$$B \rightarrow bB \mid a$$

$$L(M) = L(G) =$$

$$aaab^*a + b^*a$$

In General

A right-linear grammar G

has variables: $V_0, V_1, V_2, \ldots$

and productions: $V_i \rightarrow a_1 a_2 \cdots a_m V_j$

or

$V_i \rightarrow a_1 a_2 \cdots a_m$

We construct the NFA M such that:

each variable V_i corresponds to a node:



special final state

For each production: $V_i \rightarrow a_1 a_2 \cdots a_m V_j$

we add transitions and intermediate nodes



For each production: $V_i \to a_1 a_2 \cdots a_m$

we add transitions and intermediate nodes



Resulting NFA M looks like this:



It holds that: $L(G) = L(M)$

The case of Left-Linear Grammars

Let G be a left-linear grammar

We will prove: $L(G)$ is regular

Proof idea:

We will construct a right-linear grammar G' with $L(G) = L(G')^R$

Since G is left-linear grammar the productions look like:

$$A \to B a_1 a_2 \cdots a_k$$

$$A \to a_1 a_2 \cdots a_k$$

Construct right-linear grammar G'

Left linear G

$$A \to Ba_1a_2 \cdots a_k$$

$$A \to Bv$$

Right linear G'

$$A \to a_k \cdots a_2a_1B$$

$$A \to v^R B$$

Construct right-linear grammar G'

Left linear G

$$A \to a_1 a_2 \cdots a_k$$

$$A \to v$$

Right linear G'

$$A \to a_k \cdots a_2 a_1$$

$$A \to v^R$$

It is easy to see that: $L(G) = L(G')^R$

Since G' is right-linear, we have:

$L(G')$ $\Rightarrow$ $L(G')^R$ $\Rightarrow$ $L(G)$

$L(G')$		$L(G')^R$		$L(G)$
Regular Language		Regular Language		Regular Language

Proof - Part 2

$$\left\{\begin{array}{l}\text{Languages}\\ \text{Generated by}\\ \text{Regular Grammars}\end{array}\right\} \supseteq \left\{\begin{array}{l}\text{Regular}\\ \text{Languages}\end{array}\right\}$$

Any regular language L is generated by some regular grammar G

Any regular language L is generated by some regular grammar G

Proof idea:

Let M be the NFA with $L = L(M)$.

Construct from M a regular grammar G such that $L(M) = L(G)$

Since L is regular

there is an NFA M such that $L = L(M)$

Example:



$$L = ab^*ab(b^*ab)^*$$

$$L = L(M)$$

Convert M to a right-linear grammar



$$q_0 \to aq_1$$



$$q_0 \rightarrow aq_1$$

$$q_1 \rightarrow bq_1$$

$$q_1 \rightarrow aq_2$$

$$q_0 \to aq_1$$

$$q_1 \to bq_1$$

$$q_1 \to aq_2$$

$$q_2 \to bq_3$$



$$L(G) = L(M) = L$$

G

$$q_0 \to aq_1$$

$$q_1 \to bq_1$$

$$q_1 \to aq_2$$

$$q_2 \to bq_3$$

$$q_3 \to q_1$$

$$q_3 \to \lambda$$



In General

For any transition:



Add production:

$$q \rightarrow ap$$

variable terminal variable

For any final state:



Add production: $$q_f \to \lambda$$

Since G is right-linear grammar

G is also a regular grammar

with $L(G) = L(M) = L$